

Table 2. Differences between the treatment and control groups, both uncontrolled and controlled for initial differences. Analyses were carried out using a linear regression. Reduction in scores means improvement, except for GAF scores where an increase indicates improvement.

	Differences uncontrolled for baseline			P*	Differences controlled for baseline	
	TAU (Control Group) Mean (SD)	Intervention +TAU (Experimental Group) Mean (SD)	Mean difference (CI)		Mean difference (CI)	P*
BPRS	44.8 (16.8)	36.4 (11.1)	-8.4 (-14.5, -2.2)	0.008	-9.5 (-15.4, -3.7)	0.002
PANSS Positive	18.4 (8.1)	13.4 (5.9)	-5.0 (-8.0, -1.8)	0.002	-5.3 (-8.0, -2.4)	0.000
PANSS Negative	16.4 (6.7)	14.3 (6.5)	-2.1 (-5.0, 0.8)	0.160	-2.5 (-5.3, 0.3)	0.079
PANSS General	42.3 (16.0)	34.4 (12.2)	-7.9 (-14.1, -1.7)	0.013	-8.8 (-14.5, -3.0)	0.003
PANSS Total	86.1 (30.7)	66.6 (21.1)	-19.5 (-30.9, -8.1)	0.001	-22.6 (-33.0, -12.1)	0.000
GAF	47.0 (16.7)	56.2 (13.7)	9.2 (2.4, 15.8)	0.008	10.6 (5.0, 16.1)	0.000

*p values were obtained using linear regression, to compare the differences between two groups at follow up, both with and without controlling for the baseline differences

BPRS= Brief Psychotic Rating Scale

PANSS= Positive And Negative Syndrome Scale

GAF= Global Assessment of Functioning