



Synthesis of the Beryllium 3131Å Spectral Region

Johanna F. Ashwell¹, R.D. Jeffries¹, and Barry Smalley¹

Astrophysics Group, School of Chemistry and Physics, Keele University, Staffordshire, ST5 5BG, UK e-mail: jfa@astro.keele.ac.uk

Abstract. The Beryllium spectral region of the Sun, Procyon and 4 stars in the open cluster NGC6633 up to $T_{\text{eff}} = 7500\text{K}$ have been synthesised using ATLAS9 model atmospheres and the MOOG spectral synthesis program.

The line list used for these syntheses has been modified from the ATLAS9 line list to improve the quality of the fits in light of the improved opacities in the new version of the MOOG code. Significant changes have been made to the Mn I line at ATLAS9 wavelength 3131.037Å and an OH line has been added at 3131.358Å. In addition there are a number of minor changes to gf -values throughout the synthesised region thus improving the fit for the spectra across the temperature range considered.

Key words. Line: profiles - stars: abundances - stars: atmospheres - stars: chemically peculiar

1. Introduction

The synthesis of the Beryllium region presented in this paper was carried out with the objective of determining the Be abundance of 4 stars in the open cluster NGC6633 (one of which exhibits chemical peculiarities) using new CCD data from the UV Visual Echelle Spectrograph (UVES) on the Very Large Telescope (VLT) (see Ashwell et al. (2005)).

As both components of the Be II doublet are blended with other lines it is necessary to synthesise this region of the spectrum. This synthesis has been carried out using an updated version of the LTE analysis code MOOG (Snedden 2002) and atmospheres interpolated for the individual stellar parameters from the Kurucz ATLAS9 grids.

Send offprint requests to: Johanna F. Ashwell

2. Line List Selection

The 3131Å spectral region which contains the Be II resonance doublet is rich with strong atomic and molecular lines in solar type stars. This results in substantial line absorption and a deficit of true continuum regions making normalisation difficult.

Though, spectrum synthesis can reduce normalisation problems by taking into account possible blending features, laboratory studies of the identification, precise wavelengths and oscillator strengths of many features in this spectral region are limited.

The initial line list used consists of selected atomic and molecular lines from the ATLAS9/Kurucz CDROMs with the lines selected on the basis of having an excitation potential less than 10.0 eV and $\log gf$ greater than 10.0.




Fig. 1. Spectral synthesis of the Solar Be II doublet (3130.4Å and 3131.1Å) region from the NOAO Solar atlas (upper) and Procyon (lower). The upper portion of each plot shows the spectrum (solid line) and synthesis (dashed line). The lower portion of each plot shows the residuals.

3. Line List Calibration

Throughout the 3129.5 - 3132.5Å region synthesised the oscillator strengths for a number of lines were altered from their ATLAS9 values to better fit both the NOAO Solar Atlas (Kurucz et al. 1984) and a spectrum of Procyon (F5 IV) obtained with UVES by Bagnulo et al. (2003) (see Figure 1).

Possible arbitrary changes were avoided by adopting the philosophy that gf -values which needed the least adjustment should be chosen where a number of lines affected the same feature. The resulting solar abundance, $A(\text{Be}) = 1.15$, is in excellent agreement with the photospheric abundance quoted in Anders & Grevesse (1989), whilst Procyon is confirmed as very Be-depleted, $A(\text{Be}) \leq -0.5$.

To simultaneously match the Procyon spectrum, with $T_{\text{eff}} = 6700\text{K}$, $\log g = 4.05$ (Lemke et al. 1993); and the Sun a few major changes were made.

4. Line List Modifications

Of these major changes made to the ATLAS9 line list only two affect the Be II lines directly. The first is an alteration of both the wavelength and gf -value of an Mn I line and the second is the addition an OH molecular line where a line is clearly missing in the ATLAS9 line list.

The wavelength of the Mn I in the ATLAS9 list is stated as 3131.037Å with a $\log gf$ -value of 1.725 and gives the fit shown in the left plot of Figure 2 using MOOG. The synthesis (dashed line) is clearly discrepant from the spectrum (solid line) appearing to be positioned at too longer wavelength.

However, a remedy to this problem is suggested by King et al. (1997). They suggest that shifting the Mn I line to 3131.017Å (a shift of 0.02Å is not unreasonable when considering errors) could account for the discrepancy. Therefore, by shifting this line and increasing its gf -value by +1.56 dex the fit shown in the right hand plot of Figure 2 is achieved.

The addition of an OH molecular line as seen in the right hand plot of figure 3 corrects for the apparently missing line at 3131.350Å. The added line with a wavelength of 3131.358Å, excitation potential of 1.941 and $\log gf$ -value 1.347 was used in the line list for various King papers including King et al. (1997) however, the gf -value used here has been increased by 0.110 dex.

5. Conclusions

Via a combination of small changes to the gf -values of a number of lines across the spectral region synthesised and a few more major changes in the immediate vicinity of the Be II doublet lines an accurate fit to the spectra of both the Sun and Procyon have been achieved.

Furthermore, the same line list is capable of synthesising the high resolution VLT/UVES spectra of four stars in NGC6633 with tem-




Fig. 2. A magnified section of Figure 1 showing the changes made to the Mn I line at 3131.037Å. The left plot shows the original ATLAS9 line list parameters and the right plot shows the fit achieved by altering these parameters (see section 4).




Fig. 3. A magnified section of Figure 1 showing the changes made redward of the Be II doublet. The left plot shows the original ATLAS9 line list parameters and the right plot shows the fit achieved by adding an OH molecular line (see section 4).

peratures ranging from 6300K to 7600K (Ashwell et al. 2005).

Acknowledgements. JFA acknowledges the financial support of the UK Particle Physics and Astronomy Research Council (PPARC).

References

- Anders, E., & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197
 Ashwell, J.F., Jeffries, R.D., Smalley, B., Deliyannis, C.P., Steinhauer, A., & King, J.R. 2005, accepted by MNRAS
 Bagnulo, S., Jehin, E., Ledoux, C., Cabanac, R., Melo, C., Gilmozzi, R., & The ESO Paranal Science Operations Team 2003, The

- Messenger, 114, 10
- King, J. R., Deliyannis, C. P., & Boesgaard, A. M. 1997, ApJ, 478, 778
- Kurucz, R. L., Furenlid, I., & Brault, J. T. L. 1984, National Solar Observatory Atlas, Sunspot, New Mexico: National Solar Observatory, 1984,
- Lemke, M., Lambert, D. L., & Edvardsson, B. 1993, PASP, 105, 468
- Snedden, C. 2002, moog An LTE Stellar Line Analysis Program