

EDITORIAL

# Quantitative Flow Ratio and Virtual Percutaneous Coronary Intervention for Serial Coronary Stenoses: Attractive Technology, But Still Crawling

Wael Sumaya  , MD, MRCP, PhD; Mamas A. Mamas  , BMBCh, DPhil; Rodrigo Bagur  , MD, PhD,

**C**oronary physiology assessment is an important field in which emerging technologies have been successfully used in facilitating decision-making in interventional cardiology. This started with the advent of coronary wires mounted with pressure sensors to measure pressure distal to the lesion and compare it with aortic pressure. This allowed us to derive fractional flow reserve (FFR) to determine whether a lesion is flow-limiting or not after hyperemic stimulation with adenosine.<sup>1–3</sup>

**See Article by Guan et al.**

Subsequently, resting pressure indices (also known as nonhyperemic pressure ratio) were derived including instantaneous wave-free ratio,<sup>4,5</sup> diastolic pressure ratio,<sup>6,7</sup> and resting full-cycle ratio.<sup>7,8</sup> It is now well-established that physiology-guided percutaneous coronary intervention (PCI) leads to improved outcomes, guiding treatment in hemodynamically significant lesions and allowing us to safely defer lesions that are functionally **nonsignificant** (Table). Nevertheless, both instantaneous wave-free ratio and FFR still require an invasive wire-based approach.

More recently, it has now become possible to assess physiology “virtually” with reasonable accuracy, using 3-dimensional reconstruction and computational fluid dynamics based on coronary angiography alone, avoiding the need to instrument the vessel with coronary wires.<sup>9–11</sup> Quantitative flow ratio (QFR) is one of the studied virtual tools to assess coronary flow. In the FAVOR III China (Comparison of Quantitative Flow Ratio Guided and Angiography Guided Percutaneous Intervention in Patients With Coronary Artery Disease) trial, 3825 patients were randomized to QFR-guided PCI or angiography-guided PCI. QFR-guided PCI resulted in improved outcomes with regards to the primary end point of death, myocardial infarction, or unplanned revascularization compared with angiography-guided PCI.<sup>12</sup> Further iterations of QFR, Murray-fractal law-QFR (uQFR), enabled computation of QFR from a single angiographic view.<sup>13</sup>

In this issue of the *Journal of the American Heart Association (JAHA)*, Guan and colleagues<sup>14</sup> report the results of a prospective study assessing the utility of uQFR to virtually plan PCI. Participants presented with 2 serial lesions that were visually estimated between 30% and 90% in the same epicardial vessel and separated by at least 10mm of an angiographically-looking normal segment. Fifty-four patients with 61 vessels were evaluated

**Key Words:** Editorials ■ coronary stenosis ■ percutaneous coronary intervention ■ physiology ■ technology

Correspondence to: Rodrigo Bagur, MD, PhD, University Hospital, London Health Sciences Centre, 339 Windermere Road, N6A 5A5, London, Ontario, Canada. Email: [rodrigobagur@yahoo.com](mailto:rodrigobagur@yahoo.com)

For Disclosures, see page 3.

© 2022 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

JAHA is available at: [www.ahajournals.org/journal/jaha](http://www.ahajournals.org/journal/jaha)

**Table 1. Landmark Clinical Trials Establishing the Role of Coronary Physiology in Interventional Cardiology**

Author, study, and population	Intervention	Outcomes
Pijls et al, <sup>1</sup> DEFER, 325 patients with intermediate de novo stenosis	All patients had FFR measured. FFR $\leq 0.75$ : (n=144) PCI was performed; FFR $> 0.75$ (n=181), patients were randomized to PCI or medical management	5-y event-free survival: 80% vs 73%; $P=0.52$ (deferral group vs PCI group, respectively)
Tonino et al, <sup>2</sup> FAME, 1005 patients with multivessel CAD planned for PCI	Patients were randomized to angiography-guided or FFR-guided PCI. In the FFR group, PCI was performed if FFR was $\leq 0.80$	Death, MI, and repeat revascularization at 1 y: 13.2% vs 18.3%; $P=0.02$ (FFR-guided PCI vs angiography-guided PCI, respectively)
De Bruyne et al, <sup>3</sup> FAME II, 1220 patients with stable CAD	Patients were randomized to FFR-guided PCI or medical therapy alone. In the FFR group, PCI was performed if FFR was $\leq 0.80$	Death, MI, or urgent revascularization at 1 y: 4.3% vs 12.7%; $P<0.001$ (FFR-guided PCI+medical therapy vs medical therapy alone, respectively)
Gotberg et al, <sup>4</sup> iFR-SWEDEHEART, 2037 patients with stable angina or acute coronary syndrome with planned physiology-guided PCI	Patients were randomized to FFR-guided PCI or iFR-guided PCI	Death, MI, or unplanned revascularization at 1 y: 6.1% vs 6.7%; $P=0.53$ (FFR vs iFR)
Davies et al, <sup>5</sup> DEFINE-FLAIR, 2492 patients with intermediate coronary stenosis	Patients were randomized to FFR-guided PCI or iFR-guided PCI	Death, MI, or unplanned revascularization at 1 y: 7% vs 6.8%; $P=0.83$ (FFR vs iFR)

iFR is the only nonhyperemic pressure ratio method evaluated and validated against fractional flow reserve (FFR) on hard clinical end points. CAD indicates coronary artery disease; DEFER, Deferral Versus Performance of Percutaneous Coronary Intervention of Functionally Nonsignificant Coronary Stenosis; DEFINE-FLAIR, Functional Lesion Assessment of Intermediate Stenosis to Guide Revascularisation; FAME, Fractional Flow Reserve Versus Angiography in Multivessel Evaluation; FAME II, Fractional Flow Reserve Versus Angiography for Multivessel Evaluation II; iFR, instantaneous wave-free ratio; iFR-SWEDEHEART, Instantaneous Wave-Free Ratio Versus Fractional Flow Reserve in Patients With Stable Angina Pectoris or Acute Coronary Syndrome: A Multicenter, Prospective, Randomized Controlled Clinical Trial Based on the Swedish Angiography and Angioplasty Registry; MI, myocardial infarction; and PCI, percutaneous coronary intervention.

in the study. All vessels were assessed by maximal hyperemic FFR, and PCI was performed if FFR was  $\leq 0.80$  (44 vessels). PCI was first performed in lesions that were believed to be most hemodynamically significant. Repeat FFR was then performed and PCI to the remaining hemodynamically significant lesion was undertaken whenever residual FFR was  $\leq 0.80$  (14 lesions). uQFR was computed at an experienced central academic core laboratory with excellent intraobserver and interobserver variability ( $\pm 0.03$ ). Predicted uQFR was also calculated after “virtual PCI” of the most significant lesion.

The authors found an excellent correlation ( $r=0.97$ ,  $P<0.001$ ) and agreement between baseline uQFR and FFR (mean difference:  $0.00\pm 0.03$ ) for evaluation of the hemodynamic significance of serial lesions in the target vessel. Discordance between FFR and uQFR was only present in 2 cases where FFR was  $> 0.80$  while uQFR was  $\leq 0.80$ .

There was no difference between  $\Delta\mu\text{QFR}$  and  $\Delta\text{FFR}$  in the proximal lesion ( $0.15\pm 0.10$  and  $0.15\pm 0.10$ , respectively;  $P=0.71$ ), with good correlation ( $r=0.79$ ,  $P<0.001$ ; mean difference of  $0.00\pm 0.06$ ). The same pattern was observed in terms of the difference between  $\Delta\mu\text{QFR}$  and  $\Delta\text{FFR}$  of the distal lesion ( $0.13\pm 0.08$  and  $0.14\pm 0.08$ , respectively;  $P=0.56$ ). There was very good correlation between  $\Delta\mu\text{QFR}$  and  $\Delta\text{FFR}$  ( $r=0.82$ ,  $P<0.001$ ; mean difference of  $0.00\pm 0.06$ ).

uQFR and FFR were consistent in judging the primary lesion by comparing the gradient through the

proximal and distal lesion. However, 1 proximal lesion (4.0%) and 2 distal lesions (10.5%) showed discordance between  $\mu\text{QFR}$  and FFR, meaning that the primary lesion was misjudged in 1 in 7 cases.

After stenting the primary lesion, the accuracy of QFR for identifying the hemodynamic significance of the nonprimary lesion was  $\approx 90\%$  (area under the curve, 0.94) and predicted  $\mu\text{QFR}$  was higher than residual observed FFR ( $0.87\pm 0.06$  versus  $0.83\pm 0.07$ ,  $P<0.001$ ). Using the same cutoff value of  $\leq 0.80$  for the detection of hemodynamically significant stenosis, the predicted uQFR was shown to be 100% specific but only  $\approx 70\%$  sensitive.

The authors are to be congratulated for their efforts to simplify physiology assessment by avoiding wire-based hyperemic methods and further guide PCI optimization using an angiographic, computer-based reconstruction and fluid dynamics algorithm. The concept is attractive as it attempts to remove the “guess work” and apply a physiologic approach to help decision-making. This approach has been shown to alter interventional cardiologists’ treatment strategy and potentially affect chosen stent size.<sup>15</sup> However, does this mean that such a strategy is ready for prime time or does it simply represent the early days of an emerging technology? Several caveats need to be taken into consideration.

First, although the QFR-guided PCI strategy is superior to angiography-guided PCI, this has not been

compared with the “gold standard,” being FFR-guided PCI for intermediate lesions in a clinical outcome trial. Second, authors included a wide range of stenoses, namely 30% to 90%, while similar studies have looked at more intermediate lesions of at least 50% stenosis. Third, the performance of a QFR-guided PCI strategy has only modest performance in identifying functional significance of the nonprimary lesion once the primary lesion has been treated, with a sensitivity of only 70%, meaning that many unnecessary PCIs would be performed from false-positives. On the other hand, the high observed specificity is certainly because many lesions included in the study were relatively minor stenosis (ie, 30%–50%), which will tend to be negative, therefore accounting for the high specificity values. Fourth, it remains uncertain as to whether physiological assessment of 2 serial borderline stenoses is the best strategy to guide PCI. Intracoronary imaging may be better suited to precisely decide on stent sizing, landing zone, and plaque vulnerability. Fifth, in this study, uQFR was measured in a central core laboratory with minimal intraobserver and interobserver variability. It is yet to be determined how variable this tool will be once introduced to mainstream laboratories. Sixth, target vessel revascularization was previously shown to be higher in patients with residual FFR of  $<0.90$ .<sup>16</sup> This may be indicative of undersized stenting, which may account for target vessel failure. Hence, in the presence of serial stenoses, this issue becomes more challenging to judge. Is it an undersized stent causing suboptimal residual FFR or is it a remaining stenosis (ie, geographical miss)? It is well-known that FFR requires several steps to assess serial lesions, including the induction of maximal hyperemia with adenosine after the completion of the first lesion PCI. In this regard, nonhyperemic pressure ratio methods simplify the assessment of multiple lesions using pullback, because individual pressure indexes do not interact with each other.<sup>17,18</sup> Nonetheless, it should be highlighted that the only nonhyperemic pressure ratio method validated against FFR that looked at patient-important clinical outcomes is instantaneous wave-free ratio (Table).<sup>4,5</sup> Notably, the same comment also applies to the paper by Guan and colleagues.<sup>14</sup> Given the lack of clinical outcomes assessment, it substantially limits the knowledge translation into daily clinical practice. Finally, while the adoption of less invasive approaches is welcome, pressure wire-based measurements provide important diagnostic information for the workup of ischemia with nonobstructive coronary arteries (INOCA), providing evidence of impaired coronary microvascular function through impaired coronary flow reserve ( $<2.0$ )<sup>19</sup> or abnormal coronary microvascular resistance indices (such as indices index of microcirculatory resistance  $\geq 25$ ). Indeed, coronary microvascular resistance has been associated with 59.1% of the variability when

calculating virtual FFR estimates.<sup>20</sup> Importantly, the diagnostic workup of a patients with coronary artery disease does not merely stop at the decision to stent or not.

In summary, the study by Guan and colleagues<sup>14</sup> is informative and sheds light for further research. Once streamlined, QFR may serve as a useful tool to supplement coronary angiography, but further studies are needed to establish its role in predicting residual FFR following PCI to help procedural planning as well as broadening the role of noninvasive tools to assess microvascular function.

## ARTICLE INFORMATION

### Affiliations

Interventional Cardiology, Division of Cardiology, Queen Elizabeth II Health Sciences Centre, Dalhousie University, Halifax, Nova Scotia, Canada (W.S.); Keele Cardiovascular Research Group, Institute for Applied Clinical Science and Centre for Prognosis Research, Institute of Primary Care and Health Sciences, Keele University, Stoke-on-Trent, United Kingdom (M.A.M., R.B.); and Interventional Cardiology, Division of Cardiology, London Health Sciences Centre, Western University, London, Ontario, Canada (R.B.).

### Disclosures

None.

## REFERENCES

- Pijls NH, van Schaardenburgh P, Manoharan G, Boersma E, Bech JW, van't Veer M, Bar F, Hoorntje J, Koolen J, Wijns W, et al. Percutaneous coronary intervention of functionally nonsignificant stenosis: 5-year follow-up of the DEFER study. *J Am Coll Cardiol*. 2007;49:2105–2111. doi: 10.1016/j.jacc.2007.01.087
- Tonino PA, De Bruyne B, Pijls NH, Siebert U, Ikeno F, van't Veer M, Klauss V, Manoharan G, Engstrom T, Oldroyd KG, et al. Fractional flow reserve versus angiography for guiding percutaneous coronary intervention. *N Engl J Med*. 2009;360:213–224. doi: 10.1056/NEJMoa0807611
- De Bruyne B, Pijls NH, Kalesan B, Barbato E, Tonino PA, Piroth Z, Jagic N, Mobius-Winkler S, Rioufol G, Witt N, et al. Fractional flow reserve-guided pci versus medical therapy in stable coronary disease. *N Engl J Med*. 2012;367:991–1001. doi: 10.1056/NEJMoa1205361
- Gotberg M, Christiansen EH, Gudmundsdottir IJ, Sandhall L, Danielewicz M, Jakobsen L, Olsson SE, Ohagen P, Olsson H, Omerovic E, et al. Instantaneous wave-free ratio versus fractional flow reserve to guide PCI. *N Engl J Med*. 2017;376:1813–1823. doi: 10.1056/NEJMoa1616540
- Davies JE, Sen S, Dehbi HM, Al-Lamee R, Petraco R, Nijjer SS, Bhandi R, Lehman SJ, Walters D, Sapontis J, et al. Use of the instantaneous wave-free ratio or fractional flow reserve in PCI. *N Engl J Med*. 2017;376:1824–1834. doi: 10.1056/NEJMoa1700445
- Van't Veer M, Pijls NHJ, Hennigan B, Watkins S, Ali ZA, De Bruyne B, Zimmermann FM, van Nunen LX, Barbato E, Berry C, et al. Comparison of different diastolic resting indexes to iFR: are they all equal? *J Am Coll Cardiol*. 2017;70:3088–3096. doi: 10.1016/j.jacc.2017.10.066
- Lee JM, Choi KH, Park J, Hwang D, Rhee TM, Kim J, Park J, Kim HY, Jung HW, Cho YK, et al. Physiological and clinical assessment of resting physiological indexes. *Circulation*. 2019;139:889–900. doi: 10.1161/CIRCULATIONAHA.118.037021
- Svanerud J, Ahn JM, Jeremias A, van't Veer M, Gore A, Maehara A, Crowley A, Pijls NHJ, De Bruyne B, Johnson NP, et al. Validation of a novel non-hyperaemic index of coronary artery stenosis severity: the resting full-cycle ratio (VALIDATE FFR) study. *EuroIntervention*. 2018;14:806–814. doi: 10.4244/EIJ-D-18-00342
- Xu B, Tu S, Qiao S, Qu X, Chen Y, Yang J, Guo L, Sun Z, Li Z, Tian F, et al. Diagnostic accuracy of angiography-based quantitative flow ratio

- measurements for online assessment of coronary stenosis. *J Am Coll Cardiol*. 2017;70:3077–3087. doi: [10.1016/j.jacc.2017.10.035](https://doi.org/10.1016/j.jacc.2017.10.035)
10. Westra J, Andersen BK, Campo G, Matsuo H, Koltowski L, Eftekhari A, Liu T, Di Serafino L, Di Girolamo D, Escaned J, et al. Diagnostic performance of in-procedure angiography-derived quantitative flow reserve compared to pressure-derived fractional flow reserve: the FAVOR II Europe-Japan study. *J Am Heart Assoc*. 2018;7:e009603. doi: [10.1161/JAHA.118.009603](https://doi.org/10.1161/JAHA.118.009603)
  11. Fearon WF, Achenbach S, Engstrom T, Assali A, Shlofmitz R, Jeremias A, Fournier S, Kirtane AJ, Kornowski R, Greenberg G, et al. Accuracy of fractional flow reserve derived from coronary angiography. *Circulation*. 2019;139:477–484. doi: [10.1161/CIRCULATIONAHA.118.037350](https://doi.org/10.1161/CIRCULATIONAHA.118.037350)
  12. Xu B, Tu S, Song L, Jin Z, Yu B, Fu G, Zhou Y, Wang J, Chen Y, Pu J, et al. Angiographic quantitative flow ratio-guided coronary intervention (FAVOR III China): a multicentre, randomised, sham-controlled trial. *Lancet*. 2021;398:2149–2159. doi: [10.1016/S0140-6736\(21\)02248-0](https://doi.org/10.1016/S0140-6736(21)02248-0)
  13. Tu S, Ding D, Chang Y, Li C, Wijns W, Xu B. Diagnostic accuracy of quantitative flow ratio for assessment of coronary stenosis significance from a single angiographic view: a novel method based on bifurcation fractal law. *Catheter Cardiovasc Interv*. 2021;97:1040–1047. doi: [10.1002/ccd.29592](https://doi.org/10.1002/ccd.29592)
  14. Guan S, Gan Q, Han W, Zhai X, Wang M, Chen Y, Zhang L, Li T, Chang X, Liu H, et al. Feasibility of quantitative flow ratio virtual stenting for guidance of serial coronary lesions intervention. *J Am Heart Assoc*. 2022;11:e025663. doi: [10.1161/JAHA.122.025663](https://doi.org/10.1161/JAHA.122.025663)
  15. Gosling RC, Adam Z, Barmby DS, Iqbal J, Morgan KP, Richardson JD, Rothman AMK, Lawford PV, Hose DR, Gunn JP, et al. The impact of virtual fractional flow reserve and virtual coronary intervention on treatment decisions in the cardiac catheter laboratory. *Can J Cardiol*. 2021;37:1530–1538. doi: [10.1016/j.cjca.2021.06.004](https://doi.org/10.1016/j.cjca.2021.06.004)
  16. Diletti R, Masdjedi K, Daemen J, van Zandvoort LJC, Neleman T, Wilschut J, Den Dekker WK, van Bommel RJ, Lemmert M, Kardys I, et al. Impact of poststenting fractional flow reserve on long-term clinical outcomes: the FFR-Search study. *Circ Cardiovasc Interv*. 2021;14:e009681. doi: [10.1161/CIRCINTERVENTIONS.120.009681](https://doi.org/10.1161/CIRCINTERVENTIONS.120.009681)
  17. Omori H, Kawase Y, Mizukami T, Tanigaki T, Hirata T, Kikuchi J, Ota H, Sobue Y, Miyake T, Kawamura I, et al. Comparisons of nonhyperemic pressure ratios: predicting functional results of coronary revascularization using longitudinal vessel interrogation. *JACC Cardiovasc Interv*. 2020;13:2688–2698. doi: [10.1016/j.jcin.2020.06.060](https://doi.org/10.1016/j.jcin.2020.06.060)
  18. Zeitouni M, Kerneis M. Resting coronary physiology in stable coronary artery disease: still building evidence. *JACC Cardiovasc Interv*. 2020;13:2699–2701. doi: [10.1016/j.jcin.2020.07.033](https://doi.org/10.1016/j.jcin.2020.07.033)
  19. Kunadian V, Chieffo A, Camici PG, Berry C, Escaned J, Maas A, Prescott E, Karam N, Appelman Y, Fraccaro C, et al. An EAPCI Expert Consensus document on ischaemia with non-obstructive coronary arteries in collaboration with european society of cardiology working group on coronary pathophysiology & microcirculation endorsed by coronary vasomotor disorders international study group. *EuroIntervention*. 2021;16:1049–1069. doi: [10.4244/EIJY20M07\\_01](https://doi.org/10.4244/EIJY20M07_01)
  20. Morris PD, Silva Soto DA, Feher JFA, Rafiroiu D, Lungu A, Varma S, Lawford PV, Hose DR, Gunn JP. Fast virtual fractional flow reserve based upon steady-state computational fluid dynamics analysis: results from the virtu-fast study. *JACC Basic Transl Sci*. 2017;2:434–446. doi: [10.1016/j.jacbts.2017.04.003](https://doi.org/10.1016/j.jacbts.2017.04.003)