# **Eclipsing Binaries in Open Clusters**

## John Southworth

Department of Physics, University of Warwick, Coventry, CV4 7AL, UK email: jkt@astro.keele.ac.uk

**Abstract.** Detached eclipsing binaries are very useful objects for calibrating theoretical stellar models and checking their predictions. Detached eclipsing binaries in open clusters are particularly important because of the additional constraints on their age and chemical composition from their membership of the cluster. I compile a list containing absolute parameters of well-studied eclipsing binaries in open clusters, and present new observational data on the B-type systems V1481 Cyg and V2263 Cyg which are members of the young open cluster NGC 7128.

**Keywords.** stars: binaries: eclipsing, stars: binaries: close, stars: distances, stars: evolution, stars: abundances, open clusters and associations: general

## 1. The problem

Theoretical stellar evolution models are of huge importance to stellar and galactic astrophysics because they are basically the only way of deriving the age, internal structure and composition of stars and galaxies from simple observational data (e.g. Meynet et al. 2005; Nordström et al. 2004). The accuracy of the results depends on the predictive power of the models, but this is severely limited by the parameteric treatment of several effects, including convective core overshooting ( $\alpha_{\rm OV}$ ), the efficiency of convective energy transport ( $\alpha_{\rm MLT}$ ), the effects of rotation, magnetic fields and mass loss, and the need to use theoretically-calculated opacities.

The main offender is  $\alpha_{\rm OV}$ , changes in which cause major modifications to the predicted lifetimes, chemical yields and luminosities of massive stars (e.g. Maeder & Meynet 1989). This in turn has a large effect on the predicted properties and formation rates of evolved objects such as core-collapse supernovae (Eldridge & Tout 2004) and black holes. Also, changes in  $\alpha_{\rm MLT}$  can affect the derived ages of globular clusters by up to 2 Gyr (Chaboyer 1995), whilst reasonable variation in  $\alpha_{\rm OV}$  can alter the derived ages of open clusters by a factor of two (Bragaglia & Tosi 2006). It is clear that this must be improved.

Theoretical models can be calibrated with observational constraints derived using detached eclipsing binaries (dEBs) (e.g. Pols et al. 1997; VandenBerg et al. 2006). The study of these objects allows us to accurately measure the masses, radii and  $T_{\rm eff}$ s of two stars of the same age and chemical composition (Andersen 1991). Observational errors in mass and radius can be below 1% (e.g., Southworth et al. 2004e, 2005b). Accurate  $T_{\rm eff}$ s can also be found, allowing luminosities and distances to be directly measured.

dEBs have been used to constrain  $\alpha_{\rm OV}$  (Andersen et al. 1990) and its dependence on mass (Ribas et al. 2000);  $\alpha_{\rm MLT}$  was also studied by Ludwig & Salaris (1999). However, the stength of these results has been compromised by the lack of information available for each object. For a dEB, the radii and effective temperatures are known for two specific masses. When comparing to theory, two of these four datapoints (usually the radii) are required to deduce the age and metal abundance of the system. The remaining two are generally not accurate enough to do anything more than indicate the helium abundance or, if you assume a normal helium abundance, perhaps the extent of convective overshooting. Further information is needed to derive accurate constraints on  $\alpha_{\rm OV}$  or

#### John Southworth

**Table 1.** Parameters of well-studied dEBs in open clusters. Only systems with masses andradii known to within 5% are included, sorted by decreasing primary star mass.

dEB	Cluster	Masses (M_{\odot})	Radii (R_{\odot})	$T_{\rm eff} { m s}$ (K)	Reference
V1034 Sco	NGC $6231$	$16.8 {\pm} 0.48$	$7.45{\pm}0.07$	26200	Bouzid et al. (2005)
		$16.8 {\pm} 0.49$	$7.44{\pm}0.08$	20350	
V578 Mon	NGC 2244	$14.5 {\pm} 0.08$	$5.23 {\pm} 0.06$	$30000 \pm 500$	Hensberge et al. $(2000)$
		$10.3 {\pm} 0.06$	$4.32 {\pm} 0.07$	$26400 \pm 400$	
V453 Cyg	NGC 6871	$14.4 \pm 0.20$	$11.1 {\pm} 0.13$	$26600 \pm 500$	Southworth et al. (2004b)
		$8.55 {\pm} 0.06$	$5.49 {\pm} 0.06$	$25500 \pm 800$	
DW Car	Cr. 228	$11.2 \pm 0.10$	$4.54{\pm}0.05$	$27900 \pm 1000$	Southworth & Clausen (2006a)
		$10.5 {\pm} 0.12$	$4.28 {\pm} 0.05$	$26500 \pm 1000$	
V615 Per	NGC 869	$4.08 {\pm} 0.06$	$2.29 {\pm} 0.14$	$15000 \pm 500$	Southworth et al. (2004a)
		$3.18 {\pm} 0.05$	$1.90 {\pm} 0.09$	$11000 \pm 1000$	
V906 Sco	NGC 6475	$3.25 {\pm} 0.07$	$3.52{\pm}0.03$	$10700 \pm 500$	Alencar et al. (1997)
		$3.38 {\pm} 0.07$	$4.52 {\pm} 0.04$	$10400 \pm 500$	
GV Car	NGC 3532	$2.51 {\pm} 0.03$	$2.57 {\pm} 0.05$	$10200 \pm 500$	Southworth & Clausen (2006b)
		$1.54{\pm}0.02$	$1.43 {\pm} 0.06$	$8000 \pm 1000$	(provisional)
V618 Per	NGC 869	$2.33 {\pm} 0.03$	$1.64{\pm}0.07$	$11000 \pm 1000$	Southworth et al. (2004a)
		$1.56 {\pm} 0.03$	$1.32 {\pm} 0.07$	$8000 \pm 1000$	
HD 23642	Pleiades	$2.19 {\pm} 0.02$	$1.83 {\pm} 0.03$	$9750 {\pm} 250$	Southworth et al. (2005a)
		$1.55 {\pm} 0.02$	$1.55 {\pm} 0.04$	$7600 \pm 400$	× /
V392 Car	NGC 2516	$1.90 {\pm} 0.02$	$1.85 {\pm} 0.02$	$8850 \pm 200$	Debernardi & North (2001)
		$1.85 {\pm} 0.02$	$1.60 {\pm} 0.03$	$8650 \pm 200$	
V818 Tau	Hyades	$1.06 {\pm} 0.01$	$0.76 {\pm} 0.01$	$5530 {\pm} 100$	Torres & Ribas (2002)
	~	$0.90{\pm}0.02$	$0.77{\pm}0.01$	$4220{\pm}150$	

 $\alpha_{\text{MLT}}$ . One possibility is to measure the surface metal abundances of the stars from highdispersion spectroscopy, but this is rarely done. Another possibility is to study dEBs which display apsidal motion (rotation of the oritentation of an eccentric orbit due to tidal and relativistic effects; Claret & Willems 2002) but these are less common and require observations over a long period of time.

### 2. Solution: eclipsing binaries in open clusters

The study of eclipsing binaries in stellar clusters allows more constraints to be placed on theoretical models (e.g. Thompson et al. 2000; Lebreton et al. 2001). As the cluster stars, including both components of the dEB, were born in the same place at the same time, theoretical predictions must be able to match their observed properties for a single age and chemical composition. As a useful byproduct, the age and composition of the cluster can be derived from the theoretical models (e.g. Southworth et al. 2004ad) and its distance can be measured empirically from the properties of the dEB (e.g. Southworth et al. 2005ac). Fundamentally, this method requires a simultaneous fit of the morphology of the cluster in colour-magnitude diagrams and the properties of the dEB in mass-radiustemperature diagrams.

We are currently undertaking a research project to study dEBs which are members of open clusters with a range of ages and metallicities, in order to study how  $\alpha_{\rm OV}$  and other theoretical model parameters vary with mass and composition. Whilst results so far have been very encouraging, it has still not been possible to match a well-studied dEB to a well-studied cluster; work to remedy this is in progress. One concept which has, however, come to light is that studying several dEBs in open cluster allows you to put four or more stars with the same age and composition but with differing masses into one mass-radius or mass-temperature plot (Southworth et al. 2004ac). This raises the possibility of



Figure 1. Preliminary (no debiassing or flat-fielding) Strömgren uvby light curves of the detached eclipsing binary V1481 Cyg in NGC 7128 (y is at the top and u at the bottom).

obtaining "standard sequences" of eclipsing stars in one cluster with accurately known masses, radii,  $T_{\rm eff}$ s and surface gravities. To this end we have obtained photometry and spectroscopy of the eclipsing binaries V1481 Cyg, V2261 Cyg and V2263 Cyg in the young open cluster NGC 7128. Preliminary light curves for these are shown in Figs. 1 and 2.

## References

Alencar, S. H. P., Vaz, L. P. R., Helt, B. E., 1997, A&A, 326, 709 Andersen, J., Clausen, J. V., Nordström, B., 1990, ApJ, 363, L33 Bouzid, M. Y., Sterken, C., Pribulla, T., 2005, A&A, 437, 769 Bragaglia, A., Tosi, M., 2006, AJ, 131, 1544 Chaboyer, B., 1995, ApJ, 444, L9 Claret, A., Willems, B., 2002, A&A, 388, 518 Debernardi, Y., North, P., 2001, A&A, 374, 204 Eldridge, J. J., Tout, C. A., 2004, MNRAS, 353, 87 Hensberge, H., Pavlovski, K., Verschueren, W., 2000, A&A, 358, 553 Lebreton, Y., Fernandes, J., Lejeune, T., 2001, A&A, 374, 540 Ludwig, H.-G., Salaris, M., 1999, in ASP Conf. Ser. 173, p. 229 Maeder, A., Meynet, G., 1989, A&A, 210, 155 Meynet, G., Maeder, A., Ekström, S., 2005, in ASP Conf. Ser. 332, p. 228 Nordström, B., et al., 2004, A&A, 418, 989 Pols, O. R., Tout, C. A., Schroder, K.-P., Eggleton, P. P., Manners, J., 1997, MNRAS, 289, 869 Ribas, I., Jordi, C., Giménez, Á., 2000, MNRAS, 318, L55 Southworth, J., Clausen, J. V., 2006a, A&A, accepted Southworth, J., Clausen, J. V., 2006b, Ap&SS, in press Southworth, J., Maxted, P. F. L., Smalley, B., 2004a, MNRAS, 349, 547 Southworth, J., Maxted, P. F. L., Smalley, B., 2004b, MNRAS, 351, 1277

John Southworth



Figure 2. Preliminary (no debiassing or flat-fielding) Strömgren uvby light curve of the eclipsing binary V2263 Cyg in NGC 7128 (y is at the top and u at the bottom). As it is semi-detached it cannot be used to constrain single-star evolutionary theory, but it can provide an accurate distance to the cluster.

Southworth, J., Zucker, S., Maxted, P. F. L., Smalley, B., 2004c, MNRAS, 355, 986
Southworth, J., Maxted, P. F. L., Smalley, B., 2004d, in ASP Conf. Ser. 318, p. 218
Southworth, J., Smalley, B., Maxted, P. F. L., Etzel, P. B., 2004e, in IAU Symp. 224, p. 548
Southworth, J., Maxted, P. F. L., Smalley, B., 2005a, A&A, 429, 645
Southworth, J., et al., 2005b, MNRAS, 363, 529
Southworth, J., Maxted, P. F. L., Smalley, B., 2005c, in IAU Colloq. 196, p. 361
Thompson, I. B., et al., 2001, AJ, 121, 3089
Torres, G., Ribas, I., 2002, ApJ, 567, 1140
VandenBerg, D. A., Bergbusch, P. A., Dowler, P. D., 2006, ApJS, 162, 375