

Patients' experience of shoulder disorders: a systematic review of qualitative studies for the OMERACT Shoulder Core Domain Set

Matthew J Page¹, Denise A O'Connor^{2,3}, Mary Malek⁴, Romi Haas^{2,3}, Dorcas Beaton⁵, Hsiaomin Huang⁶, Sofia Ramiro^{7,8}, Pamela Richards⁹, Marieke JH Voshaar¹⁰, Beverley Shea^{11,12}, Arianne P Verhagen¹³, Samuel L Whittle¹⁴, Danielle A van der Windt¹⁵, Joel J Gagnier^{6,16}, Rachelle Buchbinder^{2,3},
for the OMERACT Shoulder Core Set Working Group

1. School of Public Health and Preventive Medicine, Monash University, Melbourne, Victoria, Australia
2. Monash Department of Clinical Epidemiology, Cabrini Institute, Malvern, Victoria, Australia
3. Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Monash University, Melbourne, Victoria, Australia
4. Monash University, Clayton, Victoria, Australia
5. Institute of Health & Work and the University of Toronto, Toronto, Ontario, Canada
6. Department of Orthopaedic Surgery, University of Michigan, Ann Arbor, Michigan, US
7. Department of Rheumatology, Leiden University Medical Center, Leiden, The Netherlands
8. Zuyderland Medical Center, Heerlen, The Netherlands
9. University of Bristol, Academic Rheumatology Bristol, Bristol, UK
10. Department of Psychology, Health and Technology, University of Twente, Enschede, The Netherlands
11. Ottawa Hospital Research Institute, Clinical Epidemiology Program, Ottawa, Ontario, Canada
12. School of Epidemiology and Public Health, Faculty of Medicine, University of Ottawa, Ottawa, Ontario, Canada
13. University of Technology Sydney, Sydney, New South Wales, Australia
14. Rheumatology Unit, The Queen Elizabeth Hospital, Adelaide, South Australia, Australia

15. Arthritis Research UK Primary Care Centre, Institute for Primary Care and Health Sciences, Keele University, Keele, UK

16. Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, Michigan, US

Correspondence to: Dr. Matthew Page, School of Public Health and Preventive Medicine, Monash University, 553 St Kilda Road, Melbourne, Victoria, 3004, Australia. Email address: matthew.page@monash.edu

Word count: 3,431 (abstract 248)

ABSTRACT

Objectives: To describe the experiences (including symptoms and perceived impacts on daily living) of people with a shoulder disorder.

Methods: Systematic review of qualitative studies. We searched for eligible qualitative studies indexed in Ovid MEDLINE, Ovid Embase, CINAHL (EBSCO), SportDiscus (EBSCO) and Ovid PsycINFO up until November 2017. Two authors independently screened studies for inclusion, appraised their methodological quality using the CASP checklist, used thematic synthesis methods to generate themes describing the experiences reported by participants, and assessed the confidence in the findings using the GRADE-CERQual approach.

Results: The inclusion criteria were met by eight studies, which included 133 participants (49 females and 84 males) with either rotator cuff disease, adhesive capsulitis, proximal humeral fracture, shoulder instability or unspecified shoulder pain. We generated seven themes to describe what people in the included studies reported experiencing: pain; physical function/activity limitations; participation restriction; sleep disruption; cognitive dysfunction; emotional distress; and

other pathophysiological manifestations (other than pain). There were interactions between the themes, with particular experiences impacting on others (e.g. pain leading to reduced activities and sleep disruption). Following grading of the evidence, we considered it likely that most of the review findings were a reasonable representation of the experiences of people with shoulder disorders.

Conclusion: Patients with shoulder disorders contend with considerable disruption to their life. The experiences described should be considered by researchers seeking to select the most appropriate outcomes to measure in clinical trials and other research studies in people with shoulder disorders.

Systematic review registration: PROSPERO CRD42017082628

Keywords: Shoulder pain; Qualitative research; Systematic review; Qualitative evidence synthesis; Outcome assessment

KEY MESSAGES

- People with shoulder disorders contend with several disruptive experiences.
- Some experiences appear to impact on others (e.g. pain can lead to avoidance of particular activities and sleep disruption).
- Identifying the problems most important to people with shoulder disorders may improve patient-centred care.

BACKGROUND

Shoulder disorders, including rotator cuff disease, adhesive capsulitis, glenohumeral osteoarthritis and other conditions such as instability and humeral head fractures, affect a considerable proportion of the general population, with an estimated point prevalence ranging from 7 to 26% (1, 2). Common experiences of people with shoulder disorders include pain that can lead to problems with performing daily activities, such as dressing and bathing (3, 4). Such experiences should be reflected in outcome measures used to investigate the effectiveness of treatment options for shoulder

disorders. However, the outcomes used to evaluate interventions for people with shoulder disorders are not consistently measured in randomised trials. For example, in our previous analysis of shoulder trials published between 1954 and 2015, we found that 90% measured pain, 71% measured performance of activities of daily living, and 6% measured participation in work (5). This diversity in outcome measurement limits our ability to compare findings between studies and synthesise data in meta-analyses.

The Outcome Measures in Rheumatology (OMERACT) Shoulder Core Set Working Group was established in 2015 to develop a core domain set for clinical trials of interventions for shoulder disorders (6). A core domain set is an agreed minimum set of outcome domains (i.e. constructs such as pain or function) that should be measured and reported in all clinical trials for a particular health condition, which can help reduce the diversity in outcome measurement in future trials (7). We have used several approaches to inform the development of a core domain set for shoulder disorders. As noted above, we examined what outcome domains (such as pain) have been measured in previous shoulder trials, to generate a list of potentially important domains (5, 8). We also conducted an international Delphi study, in which patients, clinicians and researchers were asked to consider the domains identified from our previous literature review, judge the importance of each domain, and identify any domains missing from the list (4). Findings of this research were discussed at face-to-face stakeholder meetings prior to and during the OMERACT 2016 conference, where we sought consensus on a preliminary core domain set (9).

Since the OMERACT 2016 conference, greater emphasis has been placed on the value of qualitative research to inform the development of core domain sets (10, 11). Using qualitative methods to explore the lived experience and perspectives of people with shoulder disorders may identify additional important outcome domains that have not been measured in existing trials or considered within the Delphi process. It may also highlight areas that clinicians should consider, which may

improve patient-centred care (12). We are aware of several qualitative studies that have investigated the lived experience of people with shoulder disorders, but to our knowledge, there has been no attempt to synthesise the findings of these studies. Therefore, we conducted a systematic review of qualitative studies to address the following question: what are the experiences (including symptoms and perceived impacts on daily living) of people with a shoulder disorder?

METHODS

We registered our systematic review in PROSPERO in November 2017 (CRD42017082628; a full protocol for the review was uploaded at the same time). We reported our systematic review according to the Enhancing Transparency in Reporting the synthesis of Qualitative research (ENTREQ) statement (13).

Eligibility criteria

We included any study in which the authors used qualitative methods (e.g. focus groups, interviews, nominal group techniques, participant observation) to explore the experiences and perceptions of people living with a shoulder disorder. Eligible shoulder disorders included rotator cuff disease (an umbrella term to classify disorders of the rotator cuff, including subacromial impingement syndrome, rotator cuff tendinopathy or tendinitis, partial or full thickness rotator cuff tear, calcific tendinitis and subacromial bursitis (14)), adhesive capsulitis, dislocation or shoulder instability, glenohumeral or acromioclavicular osteoarthritis, glenoid labrum pathologies, proximal humeral or humeral head fractures, or unspecified shoulder pain. We included full articles (i.e. not conference abstracts) written in English, Dutch, French, Chinese or German (languages spoken by the authors), which were published in peer-reviewed journals. We included mixed-methods studies (i.e. those that reported both quantitative and qualitative data) but only if the qualitative data could be separated from the quantitative data. We also included mixed-participant studies (i.e. those that included participants with a shoulder disorder or another musculoskeletal condition, such as back pain or

neck pain) only if the data on participants with shoulder disorders could be separated from data on participants with other conditions. The primary outcomes of our review included the symptoms of people with shoulder disorders and the perceived impact of these symptoms on their daily lives, and the outcome(s) of most importance to patients, as elicited by qualitative research methods.

We excluded qualitative studies that enrolled participants with systemic inflammatory conditions, such as rheumatoid arthritis, hemiplegia causing secondary shoulder pain, or pain in the shoulder region as part of a complex myofascial neck/shoulder/arm pain condition (e.g. complex regional pain syndrome). We excluded qualitative studies of health professionals who treat people with shoulder disorders. We also excluded studies which have used quantitative instruments to measure patient-reported outcomes (e.g. surveys asking dichotomous or Likert-style questions).

Search methods

We conducted a comprehensive search to seek all available studies, rather than seeking all available concepts until theoretical saturation was achieved. We searched the following bibliographic databases, all from inception until November 2017: Ovid MEDLINE, Ovid Embase, CINAHL (EBSCO), SportDiscus (EBSCO) and Ovid PsycINFO (see all search strategies in Table S1 in the Supplementary Data). To capture any papers that may have been missed by the searches, one author (MJP) examined the references of included articles.

Selection of studies

Two review authors (MJP and MM) independently screened all titles and abstracts yielded from the searches, and all full text articles considered relevant. All disagreements were resolved via discussion or by consultation with a third review author (RB) when necessary. We summarised the selection of studies using a Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) flow diagram (15).

Data collection

Pairs of review authors (MJP and DAO or MM) independently extracted the following data from each included study using a standardised data collection form: lead author, year of publication, country, research question, sample size, participant characteristics (age and sex/gender, type(s) of shoulder disorders), data collection method (e.g. focus group, interview), data analysis method (e.g. thematic analysis), themes and sub-themes relating to individual symptoms and their perceived impacts on daily living that were generated by the study authors, along with any supporting quotes and explanations of each theme. Disagreements were resolved via discussion or through inclusion of a third review author (RB) when necessary.

Quality assessment

Pairs of review authors (MJP and RH or MM) independently assessed the quality of the included studies using the Critical Appraisal Skills Programme (CASP) checklist for qualitative studies (16). All disagreements were resolved via discussion. The CASP checklist consists of 10 items, each of which includes multiple signalling questions to help users interpret the item (29 signalling questions in total). Following the method employed by Slade et al. (17), we constructed a summary table detailing the frequency of responses ('Yes' or 'No') to each signalling question. A narrative summary of the quality of the included studies is also provided. Summary scores or quality ratings were not generated as the CASP checklist does not have a scoring matrix and a cut-off point has not been established for ratings of the quality of qualitative studies.

Data analysis

We analysed study data using the methods of thematic synthesis outlined by Thomas and Harden (18). We used an inductive approach to coding. In the first stage, we read each line of extracted text and derived codes based on the meaning and content of each extract (e.g. the line "...the pain is

really quite excruciating when I go to bed...” was coded as ‘pain severe at night’). As each new study was coded, existing codes were reviewed and revised, and new codes were added, when necessary. Once all studies had been coded, all text which had a given code was reviewed to check for consistency of coding across the studies, thus ensuring the translation of concepts from one study to another. One review author (MJP) coded all text line-by-line, and codes were verified by another review author (DAO). Any disagreements were resolved via discussion.

In the next stage, similarities and differences between the codes were reviewed in order to generate themes describing the experiences of people with a shoulder disorder. For example, all codes relating to the intensity or severity of pain were grouped under a theme labelled ‘pain’. The generation of themes was based closely on what was reported in the included studies (i.e. quotes from patients and interpretations offered by the study authors). Two review authors (MJP and DAO) generated themes independently, then discussed and finalised the themes in consultation with another review author (RB).

We drafted ‘review findings’ to summarise each theme describing the experiences of people with a shoulder disorder (19). Two review authors (MJP and RH) then independently assessed the confidence in each of the review findings using the GRADE-CERQual approach (19). This approach requires assessors to consider four components: the methodological limitations of the qualitative studies contributing to each review finding (based on the CASP assessments) (20), the coherence of data contributing to each review finding (i.e. how clear and cogent the fit is between the data from the primary studies and a review finding that synthesises the data) (21), the adequacy of data supporting each review finding (i.e. the degree of richness and the quantity of data supporting a review finding) (22), and the relevance of the studies contributing to each review finding (i.e. the extent to which the body of data from the primary studies supporting a review finding is applicable to the context specified in the review question) (23). After considering the four components, both

review authors independently judged the overall confidence in each review finding (i.e. for each theme generated) as: *high* (highly likely that the review finding is a reasonable representation of the phenomenon of interest); *moderate* (likely that the review finding is a reasonable representation of the phenomenon of interest); *low* (possible that the review finding is a reasonable representation of the phenomenon of interest); or *very low* (unclear whether the review finding is a reasonable representation of the phenomenon of interest) (19). Disagreements in confidence ratings were resolved via discussion. We presented the review findings, our confidence judgement for each finding and an explanation of the judgement in a Summary of Qualitative Findings table.

Patient involvement

Two patient research partners (PR and MJHV) were involved in the interpretation of the results and writing up of the manuscript.

RESULTS

In total, 3,664 records were identified from the searches, of which we retrieved 61 for full text screening (Figure 1). Inclusion criteria were met by eight studies (24-31). Most (n=41) of the 53 excluded studies were excluded because they were not qualitative studies. We excluded eight qualitative studies which included either participants with ineligible conditions, or mixed populations, where data on participants with shoulder disorders could not be separated from data on other participants (32-39) (Table S2 in the Supplementary Data). We also excluded four qualitative studies which explored patients' views about a particular intervention (e.g. manual therapy), rather than the symptoms and perceived impact of their shoulder disorder on daily living (40-43) (Table S2 in the Supplementary Data).

Across the eight included studies there were 133 participants (Table 1). Participants had diagnoses of rotator cuff disease (three studies), adhesive capsulitis (two studies), proximal humeral fracture,

shoulder instability or unspecified shoulder pain (one study each). The studies were conducted in the UK (four studies), Canada (two studies), Finland or New Zealand (one study each). Most participants were at least 40 years old, whereas one study focused on younger athletes with shoulder instability (mean age 27 years). There were 49 females and 84 males. Data were gathered using semi-structured interviews in seven of the studies, whereas focus groups were conducted in one study.

The methodological quality of the included studies varied

(




Figure 2; full details available in Table S3 in the Supplementary Data). Strengths which were observed in all studies included clearly stated objectives; clearly stated data collection methods, approval sought from an ethics committee; and sufficient, explicit data presented to support the findings. At least one negative response to a CASP signalling question was recorded in all but one study (30). The quality of one study (24) was particularly problematic, as responses to 18 of the 29 CASP signalling questions were negative. The following limitations were identified in more than half of the studies: no critical examination of the researchers’ own role, potential bias and influence during formulation of the research question, data collection, analysis and selection of data for presentation; no explanation as to why patients approached for participation were the most appropriate to provide access to the type of knowledge sought by the study; no justification for the setting of data collection; and no description of issues around informed consent or confidentiality




Figure 2. Critical Appraisal Skills Programme (CASP) methodological assessments




Figure 2).

We generated seven themes to describe what people in the included studies reported experiencing; these included 1) pain; 2) physical function/activity limitations; 3) participation restriction; 4) sleep disruption; 5) cognitive dysfunction; 6) emotional distress; and 7) other pathophysiological manifestations (other than pain). Table 2 provides a summary of each review finding and Table 3 presents illustrative quotes reflecting each finding. Not all participants reported having all of the above experiences. For example, while pain, physical function/activity limitations and sleep disruption were experienced by most participants in the studies, fewer participants reported experiencing cognitive dysfunction and other pathophysiological manifestations (e.g. poor muscle strength).

We generated sub-themes for three themes that addressed multiple concepts (Table 2). For example, the 'participation restriction' theme consists of sub-themes on 'work disruption', 'limited recreation' and 'limited social interactions', reflecting the finding that some participants may experience problems in one or multiple aspects of life participation. In addition, we generated sub-themes for 'emotional distress' to reflect the various emotional experiences that can accompany a shoulder disorder. Many people in the included studies reported experiencing frustration because of their shoulder disorder, whereas a smaller subset reported symptoms of depression, anxiety, and a perception that their suffering was hidden from the casual observer given the lack of outward signs of their shoulder disorder.

There were interactions between the themes describing the experiences of people with a shoulder disorder

(



Figure 3). Pain emerged as the predominant symptom that was perceived to “affect everything”, that is, lead to difficulties with performing activities of daily living, engaging in work and recreation, and getting to and staying asleep. Each of these experiences were perceived by participants to have several follow-on effects. For example, many patients claimed that work disruption caused their emotional distress, including frustration and anxiety about their recovery. Also, some people perceived that sleep disruption resulted in them having less energy to perform activities of daily living and recreational activities, and aggravated their pain, the following day

(



Figure 3).

We considered it likely that most of the review findings were a reasonable representation of the experiences of people with shoulder disorders, i.e. we had ‘moderate confidence’ in most of the review findings (Table 2; a more detailed GRADE-CERQual Qualitative Evidence Profile is available in Table S4 in the Supplementary Data). Our judgement of ‘moderate’ rather than ‘high’ confidence was driven mostly by the minor concerns we had about methodological limitations of the contributing studies, and minor concerns about the relevance of each contributing study to the review question (given that all data came from only four high-income countries). In contrast, we had very low confidence in the finding that cognitive dysfunction is experienced by people with shoulder disorders, as this finding was based on limited data from two studies, one of which had several methodological limitations. We also had low confidence in the review findings regarding the perception of hidden suffering, and other pathophysiological manifestations, because of concerns about methodological limitations, adequacy of the data and relevance of studies to the review question.

DISCUSSION

Our synthesis suggests that patients with shoulder disorders contend with considerable disruption to their life. They can experience difficulties in performing self-care activities, work, and leisure, which can cause considerable distress (particularly frustration and anxiety). Some experiences (e.g. pain, physical function/activity limitations and sleep disruption) are more common than others (e.g. cognitive dysfunction and other pathophysiological manifestations). Also, some experiences appear to impact on others (e.g. pain can lead to avoidance of particular activities and to sleep disruption), which suggests that addressing one may alleviate the other(s) in some patients. We considered it likely that most of the review findings were a reasonable representation of the experiences of people with shoulder disorders.

There are several strengths of our review. We pre-specified our methods and registered our review prior to formal screening of search results against eligibility criteria. Errors in selection, extraction, appraisal and coding of studies were minimised by the involvement of at least two authors in each process. We were able to obtain rich, unpublished data for one of the included studies (30). We included studies regardless of the type of shoulder disorder, setting or country, which enhances the generalisability of our findings. Unlike most systematic reviews of qualitative research evidence, we formally assessed our confidence in each of the review findings using the recently developed GRADE-CERQual framework (19).

The review findings should be considered in light of some limitations. Our previous research to identify important outcomes for shoulder disorders (4, 5, 9) may have influenced our analysis and selection of data for presentation. We attempted to minimise this influence by having a second researcher not involved in our prior work (DAO) code the studies. Also, we have uploaded our data collection and coding form to the Open Science Framework (<https://osf.io/xnmyd/>), so that readers can see the text and quotes that underlie each theme generated. Another limitation is that we were unable to assess the risk of reporting biases on our review findings, given the lack of suitable methods for qualitative evidence syntheses (44). We were also unable to include a study of people who had undergone rotator cuff repair as the report was not written in English (43). Finally, our findings may not reflect the experiences of people with glenohumeral or acromioclavicular osteoarthritis, glenoid labrum pathologies, or dislocation of the shoulder, given that no such people were included in the studies we identified.

It is important that clinicians ask patients which problems are arising because of their shoulder disorder and which they would like to focus on. If a mismatch exists between what clinicians and patients consider important in terms of progress and recovery, then patients may not receive the

care most appropriate for them. For example, addressing strength or range of movement may not be most helpful to a patient whose main concerns might be night pain or anxiety about certain movements causing further damage, or how to modify activities so that they can continue to work. Addressing treatment expectations and supporting psychosocial needs may help improve outcomes for people with shoulder disorders.

There are several avenues for further research in this area. It would be useful to explore the experiences of people with shoulder disorders in low- and middle-income countries, as these may differ from those documented in this review, which is based on data from high-income countries only. Such research could also be conducted on people with shoulder disorders that were not represented in this review (e.g. glenohumeral osteoarthritis), or on people underrepresented in this review (e.g. only 11% of participants with rotator cuff tear or instability were women). Finally, more methodological research could be conducted to evaluate the contribution of qualitative evidence syntheses to the development of core domain sets. For example, when we mapped the outcome domains arising from the current review of qualitative studies against the domains included in our 2016 preliminary core domain set for shoulder disorders (9), we uncovered one potentially missing domain – cognitive dysfunction. This and other findings of our review informed discussions at the OMERACT 2018 conference, where we sought endorsement from the wider OMERACT community on our core domain set for shoulder disorders (45).

Our thematic synthesis revealed that patients with shoulder disorders may need to contend with several disruptive experiences, including pain, physical function/activity limitations, participation restriction, sleep disruption, cognitive dysfunction, emotional distress, and other pathophysiological manifestations such as loss of muscle strength or reduced range of shoulder movement. A better understanding of patients' experiences is useful for clinicians who treat them and can also inform

selection of the most appropriate outcomes to measure in clinical trials and other research studies for shoulder disorders.

Acknowledgments

We thank Carol Payne (Norfolk and Norwich University Hospital) for providing us with unpublished data from her study.

Competing Interests

We have read the journal's policy and the authors of this manuscript have the following competing interests: DB is an author of one of the studies included in this systematic review, but was not involved in eligibility assessment, data collection, or quality assessment.

Funding

There was no direct funding for this study. MJP is supported by an Australian National Health and Medical Research Council (NHMRC) Early Career Fellowship. RB is supported by an Australian NHMRC Senior Principal Research Fellowship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

All authors declare to meet the ICMJE conditions for authorship. RB conceived the study design. MJP, DAO, DB, DVDW, JIG and RB provided input into the study design. MJP and MM selected studies for inclusion. MJP, DAO and MM collected data from studies. MJP, RH and MM assessed the quality of studies. MJP and RH assessed the confidence in review findings. MJP, DAO performed data coding and thematic synthesis, with contribution from RB. MJP wrote the first draft of the article. All authors contributed to revisions of the article. All authors approved the final version of the article.

Data availability

The study protocol, data collection form, and coding for this study are available on the Open Science Framework: <https://osf.io/vq4mz/>

References

1. Linsell L, Dawson J, Zondervan K, Rose P, Randall T, Fitzpatrick R, et al. Prevalence and incidence of adults consulting for shoulder conditions in UK primary care; patterns of diagnosis and referral. *Rheumatology*. 2006;45:215-21.
2. Ostor AJ, Richards CA, Prevost AT, Speed CA, Hazleman BL. Diagnosis and relation to general health of shoulder disorders presenting to primary care. *Rheumatology (Oxford)*. 2005;44(6):800-5.
3. Largacha M, Parsons Iv IM, Campbell B, Titelman RM, Smith KL, Matsen Iii F. Deficits in shoulder function and general health associated with sixteen common shoulder diagnoses: A study of 2674 patients. *Journal of Shoulder and Elbow Surgery*. 2006;15(1):30-9.
4. Page MJ, Huang H, Verhagen AP, Buchbinder R, Gagnier JJ. Identifying a core set of outcome domains to measure in clinical trials for shoulder disorders: a modified Delphi study. *RMD open*. 2016;2(2):e000380.

5. Page MJ, Huang H, Verhagen AP, Gagnier JJ, Buchbinder R. Outcome reporting in randomized trials for shoulder disorders: Literature review to inform the development of a core outcome set. *Arthritis care & research*. 2018;70(2):252-9.
6. Gagnier JJ, Page MJ, Huang H, Verhagen AP, Buchbinder R. Creation of a core outcome set for clinical trials of people with shoulder pain: a study protocol. *Trials*. 2017;18(1):336.
7. Boers M, Kirwan JR, Wells G, Beaton D, Gossec L, d'Agostino M-A, et al. Developing Core Outcome Measurement Sets for Clinical Trials: OMERACT Filter 2.0. *Journal of Clinical Epidemiology*. 2014;67(7):745-53.
8. Page MJ, McKenzie JE, Green SE, Beaton DE, Jain NB, Lenza M, et al. Core domain and outcome measurement sets for shoulder pain trials are needed: systematic review of physical therapy trials. *J Clin Epidemiol*. 2015;68(11):1270-81.
9. Buchbinder R, Page MJ, Huang H, Verhagen AP, Beaton D, Kopkow C, et al. A Preliminary Core Domain Set for Clinical Trials of Shoulder Disorders: A Report from the OMERACT 2016 Shoulder Core Outcome Set Special Interest Group. *The Journal of rheumatology*. 2017.
10. Boers M, Kirwan JR, Tugwell P, Beaton D, Bingham CO 3rd, Conaghan PG, et al. The OMERACT Handbook, Updated Jan 19 2018: OMERACT.
11. Williamson PR, Altman DG, Bagley H, Barnes KL, Blazeby JM, Brookes ST, et al. The COMET Handbook: version 1.0. *Trials*. 2017;18(Suppl 3):280.
12. Keeley T, Williamson P, Callery P, Jones LL, Mathers J, Jones J, et al. The use of qualitative methods to inform Delphi surveys in core outcome set development. *Trials*. 2016;17(1):230.
13. Tong A, Flemming K, McInnes E, Oliver S, Craig J. Enhancing transparency in reporting the synthesis of qualitative research: ENTREQ. *BMC medical research methodology*. 2012;12:181.
14. Whittle S, Buchbinder R. In the clinic. Rotator cuff disease. *Ann Intern Med*. 2015;162(1):Itc1-15.

15. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine / Public Library of Science*. 2009;6(7):e1000097.
16. Critical Appraisal Skills Programme (CASP). *CASP Qualitative Research Checklist: 10 questions to help you make sense of qualitative research*. Oxford: Public Health Resource Unit; UK: Milton Keynes Primary Care Trust, 2002.
17. Slade SC, Kent P, Bucknall T, Molloy E, Patel S, Buchbinder R. Barriers to primary care clinician adherence to clinical guidelines for the management of low back pain: protocol of a systematic review and meta-synthesis of qualitative studies. *BMJ Open*. 2015;5(4):e007265.
18. Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC medical research methodology*. 2008;8:45.
19. Lewin S, Bohren M, Rashidian A, Munthe-Kaas H, Glenton C, Colvin CJ, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings-paper 2: how to make an overall CERQual assessment of confidence and create a Summary of Qualitative Findings table. *Implementation science : IS*. 2018;13(Suppl 1):10.
20. Munthe-Kaas H, Bohren MA, Glenton C, Lewin S, Noyes J, Tuncalp O, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings-paper 3: how to assess methodological limitations. *Implementation science : IS*. 2018;13(Suppl 1):9.
21. Colvin CJ, Garside R, Wainwright M, Munthe-Kaas H, Glenton C, Bohren MA, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings-paper 4: how to assess coherence. *Implementation science : IS*. 2018;13(Suppl 1):13.
22. Glenton C, Carlsen B, Lewin S, Munthe-Kaas H, Colvin CJ, Tuncalp O, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings-paper 5: how to assess adequacy of data. *Implementation science : IS*. 2018;13(Suppl 1):14.

23. Noyes J, Booth A, Lewin S, Carlsen B, Glenton C, Colvin CJ, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings-paper 6: how to assess relevance of the data. *Implementation science* : IS. 2018;13(Suppl 1):4.
24. Carter B. Clients' experiences of frozen shoulder and its treatment with Bowen technique. *Complementary therapies in nursing & midwifery*. 2002;8(4):204-10.
25. Gillespie MA, A MC, Wassinger CA, Sole G. Rotator cuff-related pain: Patients' understanding and experiences. *Musculoskeletal science & practice*. 2017;30:64-71.
26. Jones S, Hanchard N, Hamilton S, Rangan A. A qualitative study of patients' perceptions and priorities when living with primary frozen shoulder. *BMJ Open*. 2013;3(9):e003452.
27. Minns Lowe CJ, Moser J, Barker K. Living with a symptomatic rotator cuff tear 'bad days, bad nights': a qualitative study. *BMC Musculoskelet Disord*. 2014;15:228.
28. Nyman P, Palenius K, Panula H, Malkia E. Patients' experiences of shoulder problems prior to and following intervention. *Physiotherapy theory and practice*. 2012;28(3):221-31.
29. O'Hara NN, Garibaldi A, Sprague S, Jackson J, Kwok AK, Beaton D, et al. Rehabilitation, not injury or treatment details, dominate proximal humeral fracture patient concerns: a thematic analysis. *European Journal for Person Centered Healthcare*. 2017;5(3):351-6.
30. Payne C. *The assessment of outcome for musculoskeletal shoulder pain: a mixed methods study*. Norwich, UK: University of Anglia; 2010.
31. Tjong VK, Devitt BM, Murnaghan ML, Ogilvie-Harris DJ, Theodoropoulos JS. A Qualitative Investigation of Return to Sport After Arthroscopic Bankart Repair: Beyond Stability. *The American journal of sports medicine*. 2015;43(8):2005-11.
32. Bernhardsson S, Larsson MEH, Johansson K, Oberg B. "In the physio we trust": A qualitative study on patients' preferences for physiotherapy. *Physiotherapy theory and practice*. 2017;33(7):535-49.

33. Calner T, Isaksson G, Michaelson P. "I know what I want but I'm not sure how to get it"- Expectations of physiotherapy treatment of persons with persistent pain. *Physiotherapy theory and practice*. 2017;33(3):198-205.
34. Eriksson L, Lindstrom B, Ekenberg L. Patients' experiences of telerehabilitation at home after shoulder joint replacement. *Journal of telemedicine and telecare*. 2011;17(1):25-30.
35. Harding G, Parsons S, Rahman A, Underwood M. "It struck me that they didn't understand pain": the specialist pain clinic experience of patients with chronic musculoskeletal pain. *Arthritis Rheum*. 2005;53(5):691-6.
36. Hunhammar C, Nilsson-Wikmar L, Lofgren M. Striving to master variable pain: an interview study in primary care patients with non-specific long-term neck/shoulder pain. *J Rehabil Med*. 2009;41(9):768-74.
37. Hutting N, Heerkens YF, Engels JA, Staal JB, Nijhuis-van der Sanden MW. Experiences of employees with arm, neck or shoulder complaints: a focus group study. *BMC Musculoskeletal Disord*. 2014;15:141.
38. Hutting N, Detaille SI, Heerkens YF, Engels JA, Staal JB, Nijhuis-van der Sanden MW. Experiences of Participants in a Self-Management Program for Employees with Complaints of the Arm, Neck or Shoulder (CANS): A Mixed Methods Study. *Journal of occupational rehabilitation*. 2017;27(1):35-48.
39. Witaavaara B, Brulin C, Barnekow-Bergkvist M. When the body makes itself heard - The experience of bodily illness among people with neck-shoulder problems *Advances in Physiotherapy*. 2008;10:85-94.
40. Littlewood C, Malliaras P, Mawson S, May S, Walters S. Patients with rotator cuff tendinopathy can successfully self-manage, but with certain caveats: a qualitative study. *Physiotherapy*. 2014;100(1):80-5.
41. Cuff A, Littlewood C. Subacromial impingement syndrome - What does this mean to and for the patient? A qualitative study. *Man Ther*. 2018;33:24-8.

42. Sandford FM, Sanders TAB, Lewis JS. Exploring experiences, barriers, and enablers to home- and class-based exercise in rotator cuff tendinopathy: A qualitative study. *Journal of hand therapy : official journal of the American Society of Hand Therapists*. 2017;30(2):193-9.
43. Sundberg G. Patients' experiences of the rehabilitation period and long-term effects of rehabilitation after rotator cuff repair *Nordisk Fysioterapi*. 2004;8:183-9.
44. Booth A, Lewin S, Glenton C, Munthe-Kaas H, Toews I, Noyes J, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings-paper 7: understanding the potential impacts of dissemination bias. *Implementation science : IS*. 2018;13(Suppl 1):12.
45. Ramiro S, Page MJ, Whittle SL, Huang H, Verhagen AP, Beaton D, et al. The OMERACT core domain set for clinical trials of shoulder disorders (submitted). 2018.

Table 1. Characteristics of included studies

Study ID	Shoulder disorder	Country	Sample size	Gender	Age	Qualitative method
Carter 2002 (24)	Adhesive capsulitis	UK	20	10 F, 10 M	75% were aged over 50 years	Semi-structured interview
Gillespie 2017 (25)	Rotator cuff-related pain	New Zealand	10	5 F, 5 M	Median 58 years (range 47-68)	Semi-structured interview
Jones 2013 (26)	Adhesive capsulitis	UK	11	6 F, 5 M	All at least 40 years, most (n=7) in their 50s	Semi-structured interview
Minns Lowe 2014 (27)	Rotator cuff tear	UK	20	4 F, 16 M	Mean 67 years (SD 6, range 54-75)	Semi-structured interview
Nyman 2012 (28)	Supraspinatus tendinitis	Finland	21	9 F, 12 M	Mean 53 years (range 43-63)	Focus group
O'Hara 2017 (29)	Proximal humeral fracture	Canada	11	7 F, 4 M	Median 68 years (range 60-87)	Semi-structured interview
Payne 2010 (30)	Unspecified shoulder pain	UK	15	7 F, 8 M	Mean 48 years (SD 12, range 26-69)	Semi-structured interview
Tjong 2015 (31)	Shoulder instability	Canada	25	1 F, 24 M	Mean 27 years (SD 7)	Semi-structured interview

F = female; M = male; SD = standard deviation

Table 2. GRADE-CERQual Summary of Qualitative Findings table

Objective: To identify, appraise and synthesise qualitative research evidence on the experiences (including symptoms and perceived impacts on daily living) of people with a shoulder disorder

Perspective: Experiences and perspectives of people with any shoulder disorder living in any country

Summary of review finding	Studies contributing to the review finding	CERQual assessment of confidence in the evidence ^a
<p>1. Pain: Pain was the predominant symptom experienced by people with rotator cuff disease, adhesive capsulitis, proximal humeral fracture, or unspecified shoulder pain. Many people reported severe pain. Some people experienced worse pain at night or intense pain only on certain movements.</p>	(24-30)	Moderate confidence
<p>2. Physical function/activity limitations:</p>		
<p>a. Difficulty performing activities of daily living: Most people with rotator cuff disease, adhesive capsulitis, proximal humeral fracture, or unspecified shoulder pain reported that shoulder pain affected their ability to carry out activities of daily living. These activities include dressing, bathing/grooming, cooking, hanging washing, shopping, lifting/carrying, and driving. In some cases, activity limitations were attributed to difficulties with purposeful movements (e.g. overhead use, reaching, pushing/pulling, prolonged use in one position).</p>	(24-30)	Moderate confidence
<p>b. Avoiding movements: Many people with rotator cuff disease, adhesive capsulitis, shoulder instability, or unspecified shoulder pain reported avoiding movements that aggravated the pain and changing the way they performed particular activities. Some avoided certain movements or activities due to fear of re-injury.</p>	(24-28, 30, 31)	Moderate confidence
<p>3. Participation restriction:</p>		
<p>a. Work disruption: Pain and difficulties with purposeful movements affected the work performance of many people with rotator cuff disease, adhesive capsulitis, or unspecified shoulder pain. Some people reported experiencing problems undertaking physically demanding work tasks. A few people described problems with concentration due to pain, which negatively impacted work performance.</p>	(24-28, 30)	Moderate confidence
<p>b. Limited recreation: In many people with rotator cuff disease, adhesive capsulitis, proximal humeral fracture, shoulder instability, or unspecified shoulder pain, pain and difficulties with purposeful movements also affected the</p>	(24, 25, 27-31)	Moderate confidence

Summary of review finding	Studies contributing to the review finding	CERQual assessment of confidence in the evidence ^a
ability to engage in recreational/leisure activities, such as gardening and sports requiring heavier hand and arm use (e.g. tennis, climbing).		
c. Limited social interactions: Some people with rotator cuff disease or unspecified shoulder pain reported that shoulder pain restricted their social interactions (e.g. they needed someone to drive them to events, avoided crowded places due to fear of exacerbating their pain if bumped into). In some cases interpersonal relationships were adversely affected (e.g. changed role within family).	(27, 28, 30)	Moderate confidence
4. Sleep disruption: Most people with rotator cuff disease, adhesive capsulitis, proximal humeral fracture, or unspecified shoulder pain reported that pain at night disrupted their sleep. The type of sleep disruption varied; people either had trouble getting to sleep, were woken by pain, struggled to get back to sleep due to constant ‘nagging’ pain, experienced problems when lying on their affected side or experienced difficulties turning over in bed. Some reported that sleep deprivation negatively affected their cognition (e.g. concentration, memory), emotions (i.e. made them irritable), and ability to perform daily activities and participate in work. A few people perceived their shoulder pain to be aggravated by tiredness.	(24-30)	Moderate confidence
5. Cognitive dysfunction: A few people with rotator cuff disease or adhesive capsulitis reported that their shoulder pain was so severe that it prevented them from being able to concentrate on anything else. In some cases sleep deprivation due to shoulder pain affected people’s concentration and memory, which in turn affected their work-performance.	(24, 25)	Very low confidence
6. Emotional distress:		
a. Frustration: Many people with rotator cuff disease, adhesive capsulitis, shoulder instability, or unspecified shoulder pain reported feeling frustrated and annoyed by their pain. Some people reported feeling frustrated by their inability to complete daily activities, recreational activities, or physically demanding work tasks. Some people found that the sleep deprivation associated with their shoulder pain made them irritable.	(24, 25, 27, 30, 31)	Moderate confidence
b. Anxiety: A few people with rotator cuff disease, adhesive capsulitis, shoulder instability, or unspecified shoulder pain reported feeling worried about what could be the cause of their pain. Some were fearful of certain movements, activities or social interactions that might exacerbate the pain or cause re-injury.	(26, 28, 30, 31)	Moderate confidence

Summary of review finding	Studies contributing to the review finding	CERQual assessment of confidence in the evidence ^a
<p>c. Depression: Some people with rotator cuff disease, shoulder instability, or unspecified shoulder pain reported that their pain and reduced activities made them feel depressed.</p>	(27, 30, 31)	Moderate confidence
<p>d. Hidden suffering: A few people with rotator cuff disease or adhesive capsulitis reported that pain and impairments in activities of daily living occurred without obvious outward signs, so that much of the suffering involved was hidden from the casual observer. This led to the perception of much less sympathy from others than would be the case with many other conditions.</p>	(26, 28)	Low confidence
<p>7. Other pathophysiological manifestations: Some people with rotator cuff disease, adhesive capsulitis, or unspecified shoulder pain reported other pathophysiological manifestations, including reduced range of motion, loss of strength (including lack of grip), increased muscle tension, loss of automatic protective reactions, instability and problems when weight-bearing through the affected limb. A few people attributed their difficulties with certain activities (e.g. lifting, overhead use) to poor muscle strength rather than pain.</p>	(24, 26, 27, 30)	Low confidence

^aExplanations of each CERQual assessment are provided in the evidence profile presented in Table S4 in the Supplementary Data. ‘High confidence’ = highly likely that the review finding is a reasonable representation of the phenomenon of interest; ‘Moderate confidence’ = likely that the review finding is a reasonable representation of the phenomenon of interest; ‘Low confidence’ = possible that the review finding is a reasonable representation of the phenomenon of interest; ‘Very low confidence’ = unclear whether the review finding is a reasonable representation of the phenomenon of interest (19).

Table 3. Illustrative quotes reflecting each theme describing the experiences of people with a shoulder disorder

Theme	Illustrative quotes ^a
1. Pain	<p data-bbox="600 316 2018 384">“It was only maybe a few weeks after that I began to find my shoulder getting really, really sore...Well, it gradually just got worse and worse. Eventually it was really extremely painful” (24).</p> <p data-bbox="600 403 2018 469">“I have had [a] back operation, I have fallen out of trees but this pain was far worse than any other pain I have ever had in my life” (27).</p>
2. Physical function/activity limitations	<p data-bbox="241 577 2018 646">a. Difficulty performing activities of daily living “Try getting out the bath ’cause you can’t find a way up. You just cannot find a way to get out the bath! We’ve got bars on the side of the bath anyway but I couldn’t find a way. I just had to snake over the side sometimes” (26).</p> <p data-bbox="600 665 1973 694">“Shopping was more difficult, food shopping, that sort of thing, not being able to drive or carry the stuff back” (30).</p> <p data-bbox="241 716 1682 801">b. Avoiding movements “I will have to do it with the other arm, because I don’t want to hurt it (my shoulder)” (27). “My fear of dislocating it again has made me just more apprehensive about playing...so I stopped” (31).</p>
3. Participation restriction	<p data-bbox="241 874 2018 992">a. Work disruption “I could’na work at all well, I could’na work with my tools the way I usually did...I had to know my limits...It was hard to work but I had to...It was there all the time through the day...” (24). “I need to work more hours and...I physically can’t” (27).</p> <p data-bbox="241 1015 1720 1099">b. Limited recreation “I can’t lift the [grand]children. I don’t walk my dogs for fear of that (...) pulling” (25). “I couldn’t play tennis, absolutely out of the question, I couldn’t even lift the racquet up” (27).</p> <p data-bbox="241 1121 1861 1200">c. Limited social interactions “It is not fun to go and sit somewhere a long time”; “In a large crowd someone could bump into you” (28). “It has been a big upheaval and I’m lucky that I’ve got some good friends to take me out nights” (30).</p>
4. Sleep disruption	<p data-bbox="600 1222 2018 1291">“At night particularly, if I lay on it and turned over it would wake me up and I’d give a little yelp of pain, annoy the wife, roll over again...I spent the night spinning round” (24).</p> <p data-bbox="600 1310 1868 1340">“...trying to find a comfortable position to sleep, terrible...then turn over suddenly and it was painful” (27).</p>

Theme	Illustrative quotes ^a
5. Cognitive dysfunction	<p>“In my particular case it gave really violent pain to the point of not being able to concentrate on anything else” (24).</p> <p>“One night, two nights I can do, three nights I'm starting to get really tired (...) memory getting really ditsy at work” (25).</p>
6. Emotional distress	
a. Frustration	<p>“It made me more moody and lose my temper with people...my family definitely noticed a difference, I was more snappy...” (24).</p> <p>“[The pain] would annoy me more than anything, rather than worry me. Because I get frustrated if I can't do everything” (25).</p>
b. Anxiety	<p>“Even now in the bath I feel a bit anxious because you can feel a bit trapped” (30).</p> <p>“I'm frightened I'm going to suddenly get that sharp pain” (30).</p>
c. Depression	<p>“It (shoulder) just wears you down...I just got so down about it. It was awful. So she (GP) put me on amitriptyline...it was not a good time in my life” (27).</p> <p>“I remember feeling hopeless...that my shoulder would never recover” (31).</p>
d. Hidden suffering	<p>“...you don't get any sympathy from people because it is not something that stops you doing things but in yourself you feel so frustrated, because when I reached up to do anything, we do a lot of overhead stuff, there was no strength in the arm, it was very painful but nobody would say, “How is it?” You felt that they didn't understand, because you are at work with it and that was the hardest thing, and they still expected you to do things because you didn't have your arm strapped up” (26).</p> <p>“Others do not understand when it is not visible” (28).</p>
7. Other pathophysiological manifestations	<p>“I would have to use the good arm to lift it, I've no strength at all in that arm, above about that high” (27).</p> <p>“I just have no movement in me arm at all. I can sort of get it half way up in front of me and half way to the side of me...but I can't raise me arm above me head” (30).</p>

^aAll quotes reported in the included studies are presented in the data collection form, available at <https://osf.io/xnmyd/>

FIGURES

Figure 1. Flow diagram of identification, screening and inclusion of studies.




Figure 2. Critical Appraisal Skills Programme (CASP) methodological assessment.



Figure 3. Interrelationship between themes describing the experiences of people with shoulder disorders.

