BMC Molecular and Cell Biology

An interaction of heart disease-associated proteins POPDC1/2 with XIRP1 in transverse tubules and intercalated discs

Manuscript Number:	CEBI-D-20-00243R3	
Full Title:	An interaction of heart disease-associated p transverse tubules and intercalated discs	proteins POPDC1/2 with XIRP1 in
Article Type:	Research article	
Section/Category:	Molecular Basis of Disease	
Funding Information:	British Heart Foundation (PG/16/68/31991)	Prof Glenn E Morris
	British Heart Foundation (PG/19/13/34247)	Prof. Thomas Brand
	Orthopaedic Institute Ltd., Oswestry (RPG141 and RPG189)	Dr. Ian Holt
Abstract:	Background Popeye domain-containing proteins 1 and 2 transmembrane proteins involved in cyclic A required for normal cardiac pacemaking and protein interaction partners, POPDC1 and 2 compared by proteomic analysis with contro- cultured human skeletal myotubes. Results There were highly-significant interactions of (Xin actin binding repeat-containing protein A5. In adult human skeletal muscle, both XI sarcolemma and in T-tubules. The interaction adult rat heart extracts. Using new monoclor POPDC2, both proteins, together with XIRF but also at T-tubules in adult rat and human Conclusions Mutations in human POPDC1, POPDC2 pathological cardiac arrhythmias, suggestin interaction in normal cardiac conduction.	2 (POPDC1 and POPDC2) are AMP-mediated signalling processes and are d conduction. In order to identify novel 2 proteins were attached to beads and of beads in the pull-down of proteins from 5 both POPDC1 and POPDC2 with XIRP1 1), actin and, to a lesser degree, annexin RP1 and POPDC1/2 were present at the on of POPDC1 with XIRP1 was confirmed in nal antibodies specific for POPDC1 and 21, were found mainly at intercalated discs a heart. and in human XIRP1, all cause g a possible role for POPDC1/2 and XIRP1
Corresponding Author:		
	Ian Holt, Ph.D. Robert Jones and Agnes Hunt Orthopaedic Oswestry, Shropshire UNITED KINGDOM	Hospital
Corresponding Author E-Mail:	Ian Holt, Ph.D. Robert Jones and Agnes Hunt Orthopaedic Oswestry, Shropshire UNITED KINGDOM ian.holt@nhs.net	Hospital
Corresponding Author E-Mail: Corresponding Author Secondary Information:	Ian Holt, Ph.D. Robert Jones and Agnes Hunt Orthopaedic Oswestry, Shropshire UNITED KINGDOM ian.holt@nhs.net	Hospital
Corresponding Author E-Mail: Corresponding Author Secondary Information: Corresponding Author's Institution:	Robert Jones and Agnes Hunt Orthopaedic Oswestry, Shropshire UNITED KINGDOM ian.holt@nhs.net Robert Jones and Agnes Hunt Orthopaedic	Hospital Hospital
Corresponding Author E-Mail: Corresponding Author Secondary Information: Corresponding Author's Institution: Corresponding Author's Secondary Institution:	Robert Jones and Agnes Hunt Orthopaedic Oswestry, Shropshire UNITED KINGDOM ian.holt@nhs.net Robert Jones and Agnes Hunt Orthopaedic	Hospital Hospital
Corresponding Author E-Mail: Corresponding Author Secondary Information: Corresponding Author's Institution: Corresponding Author's Secondary Institution: First Author:	Ian Holt, Ph.D. Robert Jones and Agnes Hunt Orthopaedic Oswestry, Shropshire UNITED KINGDOM ian.holt@nhs.net Robert Jones and Agnes Hunt Orthopaedic	Hospital Hospital
Corresponding Author E-Mail: Corresponding Author Secondary Information: Corresponding Author's Institution: Corresponding Author's Secondary Institution: First Author: First Author Secondary Information:	Ian Holt, Ph.D. Robert Jones and Agnes Hunt Orthopaedic Oswestry, Shropshire UNITED KINGDOM ian.holt@nhs.net Robert Jones and Agnes Hunt Orthopaedic Ian Holt, Ph.D.	Hospital Hospital
Corresponding Author E-Mail: Corresponding Author Secondary Information: Corresponding Author's Institution: Corresponding Author's Secondary Institution: First Author: First Author Secondary Information: Order of Authors:	Ian Holt, Ph.D. Robert Jones and Agnes Hunt Orthopaedic Oswestry, Shropshire UNITED KINGDOM ian.holt@nhs.net Robert Jones and Agnes Hunt Orthopaedic Ian Holt, Ph.D. Ian Holt, Ph.D.	Hospital Hospital

	Roland FR Schindler
	Sally L Shirran
	Thomas Brand
	Glenn E Morris
Order of Authors Secondary Information:	
Response to Reviewers:	Re: CEBI-D-20-00243 An interaction of heart disease-associated proteins POPDC1/2 with XIRP1 in transverse tubules and intercalated discs Ian Holt; Heidi R Fuller; Roland FR Schindler; Sally L Shirran; Thomas Brand; Glenn E Morris
	Dear Dr. Latto and The Editorial Staff, Thank you for the editorial comments. We have made the necessary corrections as follows:
	1. Availability of Data and Materials:
	Thank you for providing links in your 'Availability of Data and Materials' statement. However, we note that the link is not found. Please update links within the manuscript:
	Brackets have been removed from around the link so that it now works.
	2. Gels:
	We note that some of the gel/blot images are not accurately delineated. The grouping of gels/blots cropped from different parts of the same gel, or from different gels, fields, or exposures must be made explicit (e.g., using clear delineation either with dividing lines or white space) and described in the figure legend. Additionally, we note that the full images for the first line of each image (Figure 1A, B, C and D) are missing from the Additional File. Please provide all full images for in-text images within the Additional File, clearly labelled:
	Thank you for drawing our attention to the problems with Figure 1, which have now been corrected. The figure legend has been altered to describe accurately that molecular weight markers are colourimetric images and antibody blots are chemiluminescence. Where lanes are developed for different exposure times or where lanes have been moved from non-adjacent positions, these are delineated with black lines and described in the legend. Additional file 2 has been altered so that it now contains the original colourimetric images, which show the molecular weight markers that were used for the first lanes in Figure 1. The lanes from the original blots that were used in Figure 1 are clearly labelled in Additional File 2.
	3. Rats:
	In the methods section, please clarify the original source of the rat heart used in your study. If required, please also detail the euthanization methods used in your study.
	In the "Methods" section we have included the following sentence: "The heart was removed from a five month old Sprague Dawley rat that had been euthanized humanely with an overdose of pentobarbitone anesthetic (0.5 mL/100g) via intraperitoneal injection." "Ethics approval and consent to participate" contains the following sentence: "All animal procedures were performed with approval of the Keele University Animal Welfare and Ethical Review Body and conducted under the licensed authority of the UK Home Office."
	We hope that the revised manuscript is now suitable for publication in BMC Molecular and Cell Biology.

	Thank you. Best regards, Ian Holt
Additional Information:	
Question	Response
Has this manuscript been submitted before to this journal or another journal in the BMC series?</a 	No

1	1	An interaction of heart disease-associated proteins POPDC1/2 with XIRP1 in transverse
23	2	tubules and intercalated discs
4 5 6	3	
7 8 9	4	Ian Holt ^{*1,2} , Heidi R Fuller ^{1,2} , Roland FR Schindler ³ , Sally L Shirran ⁴ , Thomas Brand ³ and
10 11	5	Glenn E Morris ^{1,2}
12 13 14	6	
15 16	7	¹ Wolfson Centre for Inherited Neuromuscular Disease, RJAH Orthopaedic
17 18 19	8	Hospital, Oswestry, SY10 7AG, UK.
20 21 22	9	² Institute for Science and Technology in Medicine, Keele University, Keele ST5 5BG, UK;
23 24	10	³ National Heart and Lung Institute, 4th Floor, Imperial Centre of Translational and
25 26 27	11	Experimental Medicine, Imperial College, Du Cane Road, London W12 0NN, UK
28 29 30	12	⁴ BSRC Mass Spectrometry and Proteomics Facility, University of St Andrews, North Haugh,
31 32	13	St Andrews, Fife, KY16 9ST, UK.
33 34 35	14	*Correspondence: ian.holt@nhs.net
36 37	15	
38 39 40	16	
41 42 43		
43 44 45		
46		
47 48		
49		
50 51		
52		
53 54		
55		
56		
57		
58 59		
60		
61		
62 63		1
64		
65		

Abstract

Background: Popeye domain-containing proteins 1 and 2 (POPDC1 and POPDC2) are transmembrane proteins involved in cyclic AMP-mediated signalling processes and are required for normal cardiac pacemaking and conduction. In order to identify novel protein interaction partners, POPDC1 and 2 proteins were attached to beads and compared by proteomic analysis with control beads in the pull-down of proteins from cultured human skeletal myotubes. **Results:** There were highly-significant interactions of both POPDC1 and POPDC2 with XIRP1 (Xin actin binding repeat-containing protein 1), actin and, to a lesser degree, annexin A5. In adult human skeletal muscle, both XIRP1 and POPDC1/2 were present at the sarcolemma and in T-tubules. The interaction of POPDC1 with XIRP1 was confirmed in adult rat heart extracts. Using new monoclonal antibodies specific for POPDC1 and POPDC2, both proteins, together with XIRP1, were found mainly at intercalated discs but also at T-tubules in adult rat and human heart. Conclusions: Mutations in human POPDC1, POPDC2 and in human XIRP1, all cause pathological cardiac arrhythmias, suggesting a possible role for POPDC1/2 and XIRP1 interaction in normal cardiac conduction. **Keywords** Popeye domain-containing, Xin actin binding repeat-containing, Cardiac conduction,

21 Intercalated discs, Transverse tubules.

Background

Popeye domain containing protein 1 (POPDC1), also known as blood vessel epicardial substance (BVES), was first found in chicken heart by subtractive hybridisation [1,2]. *Popdc1* and two related gene family members, *Popdc2* and *Popdc3*, were identified in mammals and shown to be developmentally regulated and preferentially expressed in cardiac and skeletal muscle [2]. Human *POPDC1* is found on chromosome 6q21 along with *POPDC3* in tandem array, whereas *POPDC2* is found on human chromosome 3q13.33. The POPDC proteins are highly conserved throughout the animal kingdom, suggesting that they play an essential role [3].

POPDC proteins consist of a short extracellular N-terminal sequence which is glycosylated, three transmembrane domains, a conserved intracellular Popeye domain and a variable C-terminal domain which is isoform-specific, contains regions of low complexity and may be phosphorylated [4]. POPDC1 exists at the plasma membrane as a homodimer, which is stabilised by disulphide bonds [5,6]. The predicted secondary structure of the Popeye domain contains a cyclic nucleotide binding domain, which binds the second messenger cyclic adenosine 3',5'-monophosphate (cAMP) with high affinity [7]. Interaction between POPDC proteins and the potassium two pore domain channel subfamily K member 2 (KCNK2, also known as TREK-1) has been demonstrated, which leads to an increase in KCNK2 current in isolated mouse sinus node myocytes, and this activity was reduced by an increase in cAMP levels [7]. A number of other membrane proteins have been reported to interact with POPDC proteins, including caveolin-3 (CAV3) in mouse cardiomyocytes, which is a major component of caveolae in striated muscle membranes [8].

A homozygous missense variant in *POPDC1* has been found in a family with cardiac
 arrhythmia and limb-girdle muscular dystrophy (LGMD). This autosomal recessive mutation

in POPDC1 is associated with reduced cAMP affinity [9]. More recently, three homozygous loss-of-function mutations in POPDC1 were identified in three families with LGMD and cardiac conduction abnormalities [10] and a missense mutation in POPDC1 was observed in a patient with contractures and possible mild cardiac involvement [11]. A heterozygous nucleotide substitution in POPDC2 has been associated with severe atrioventricular block [12] and homozygous missense variants in POPDC3 have been associated with limb girdle muscular dystrophy in the absence of a cardiac phenotype [13]. POPDC1 protein was down-regulated with abnormal immunolocalisation in failing human hearts and POPDC1 and POPDC3 mRNA levels were reduced in the left ventricles of end-stage failing hearts [14]. Popdc1 null mice showed impaired skeletal muscle regeneration [15] and increased sensitivity towards ischemia reperfusion [8]. Moreover, mice with null-mutations in Popdc1 or Popdc2 developed a stress-induced sinus node bradycardia due to pacemaker dysfunction [7,16]. Knockdown of *popdc2* in zebrafish by injecting embryos with morpholino oligonucleotides resulted in the aberrant development of skeletal muscle and heart. A reduction in oligonucleotide concentration lead to an improvement in the skeletal muscle pathology, but abnormalities in the cardiac conduction system remained, resulting in cardiac arrhythmia and a reduction in heart rate [17]. Immunolocalization studies with polyclonal antibodies have shown that POPDC1 and POPDC2 generally localise to the sarcolemma of control skeletal muscle, but this membrane localisation was greatly reduced in muscle tissue from patients with pathogenic mutations in POPDC1 [9,10]. In the heart, POPDC1 and POPDC2 were found at the plasma membrane of cardiomyocytes, with high levels in the cardiac conduction system [7,18].

In addition to the essential roles that POPDC proteins play in the maintenance of
 structure and function of skeletal muscles and in cardiac pacemaking and conduction,

POPDC1 may play a role in tumor formation [19]. POPDC1 is thought to have a tumor suppressor function and decreased POPDC1 expression due to DNA methylation occurs in the early stages of a number of cancers [reviewed: 20,21].

Here we report a proteomic study to identify novel POPDC1/2-interacting proteins.
The most significant hit with both POPDC1 and POPDC2 was XIRP1 (Xin actin binding repeat
containing 1, also known as Xin or cardiomyopathy-associated protein 1 (CMYA1)). Pulldown and co-immunoprecipitation experiments confirmed the interaction of XIRP1 with
POPDC1 and POPDC2. POPDC1/2 and XIRP1 displayed colocalization at the intercalated disk
and in T-tubules. Mutations for both genes in model organisms [7,17,22,23] and patients
[9,10,24] have been associated with severe cardiac conduction defects and impaired
skeletal muscle regeneration was reported for both *Popdc1* [15] and *Xirp1* [25,26] null
mutants. This suggests some overlapping function between POPDC1 and XIRP1, which could
be based in part on the interaction reported here.

XIRP1 and cardiac actin are the highest-scoring binding partners for both POPDC1 and

With the goal of identifying novel protein interaction partners for POPDC proteins in
skeletal muscle, POPDC proteins available as GST-fusion "bait" proteins (mouse POPDC1 or
POPDC2) were attached to glutathione beads and used to pull-down proteins from RIPA
extracts of human skeletal myotubes. Proteins pulled down by the beads were identified by
mass spectrometry (unfiltered results shown in: Additional file 1). Cardiac actin and XIRP1

had the highest scores for a number of significant peptides identified by mass spectrometry (Table 1) for both mouse POPDC1 and mouse POPDC2, after eliminating non-specific binding of proteins to control beads (no fusion protein attached). Annexin A5 was also present in both pull-downs. For further study, we selected from Table 1 those proteins with known involvement in cardiac conduction and / or cardiomyopathy: XIRP1 and annexin A5. Cardiac actin is the principal actin isoform during early stages of skeletal myogenesis and is only gradually replaced by skeletal muscle actin in adult skeletal muscle [27], so it is not surprising that the cardiac isoform was detected in myotube extracts. TABLE 1 TO BE INSERTED HERE Confirmation by western blotting of POPDC1 interaction with XIRP1 in extracts of both human myotube cultures and adult rat heart The presence of XIRP1 in POPDC1/2 pull downs was confirmed by western blotting with a commercial anti- XIRP1 antibody (Fig. 1A upper blot). Figure 1A lower blot is a loading control showing antibody against the GST "bait" protein. XIRP1 was also immunoprecipitated from the same human myotube extracts by mAb 1C12 against POPDC1 (Fig. 1B). Western blots are representative of pull-downs repeated using three separate extracts of the cultured myotube cell line. Recombinant POPDC1 attached to glutathione beads also pulled-down XIRP1 from a RIPA extract of adult rat heart ventricle (Fig. 1D). The pull down of annexin A5 by recombinant POPDC1 from myotubes was also confirmed by western blotting (Fig. 1C). (Uncropped blots shown in: Additional file 2) New monoclonal antibodies against POPDC1 and POPDC2

Monoclonal antibodies were prepared against recombinant protein preparations and selected for their recognition of human POPDC1 or POPDC2 proteins using techniques described previously [28]. Immunofluorescence specificity of the new POPDC1 1C12 (Fig. 2A) and POPDC2 12G12 (Fig. 2B) mAbs was shown by the staining of COS7 cells transfected with recombinant POPDC1 or POPDC2 respectively. POPDC1 1C12 did not recognise transfected POPDC2 and mAb POPDC2 12G12 did not recognise transfected POPDC1. Specificity of the mAbs was further demonstrated by competition experiments. Pre-incubation of mAb POPDC1 1C12 with recombinant POPDC1, but not with recombinant POPDC2, inhibited localisation of the mAb to the sarcolemma of human skeletal muscle (Fig. 2C). Similarly, pre-incubation of mAb POPDC2 12G12 with recombinant POPDC2, but not with recombinant POPDC1, inhibited sarcolemmal localisation (Fig. 2D). XIRP1 and POPDC1/2 are found, together with annexin A5, at intercalated discs and transverse tubules in rat or human heart sections In rat heart, XIRP1 and POPDC2 were both present at intercalated discs and near the Z-line of myofibrils (Fig. 3A,B). Unlike XIRP1, however, POPDC2 was also present at the cardiac sarcolemma (Fig. 3B). A similar staining pattern in heart was reported by Alcalay et al. [8] for POPDC1 and Soni et al. [29] for POPDC2. Intercalated discs were identified using a connexin 43 antibody, which stains neither sarcolemma nor the Z-line (Fig. 3A). Our human-specific POPDC1 mAb does not recognize the rat protein, but a commercial POPDC1

antibody (BVES pAb STJ22848) gave similar results on rat heart to our POPDC2 mAb (Fig.

3C). Fig 3C also shows the location of POPDC1 near the Z-line which lies at the centre of the

23 actin I-band stained by phalloidin (ALEXA 488 - phalloidin). This staining appears to

24 correspond to transverse tubules, since an antibody against amphiphysin II, a marker for T-

tubules, gave a similar staining pattern as POPDC2 (Fig. 3E) and POPDC proteins have transmembrane domains. Annexin A5 (ANXA5) was also present in both POPDC1 and POPDC2 pull downs (Table 1) and antibodies against this protein also located it close to the Z-line (Fig. 3D). Annexin A5 is a known component of the sarcolemma, transverse tubules and intercalated discs in the heart [30]. In human heart, mAbs POPDC1 1C12 (Fig. 4A) and POPDC2 12G12 (Fig. 4B) stained the intercalated discs and the cardiac sarcolemma whereas XIRP1 was present at the intercalated discs only. There were similar staining patterns in rat and human hearts, but cross striations were difficult to resolve in the human heart sections. In low power microscopy (40x) of human heart, T-tubules were only faintly-stained (Fig. 4) compared with the higher power microscopy (63x) of rat heart (Fig.3), suggesting that intercalated discs may be the major site of POPDC-XIRP1 interaction in the heart. Figures 3 and 4 show representative images of results that were obtained with sections from two different rat hearts and confirmed with the sections from a human heart. POPDC1, POPDC2, XIRP1 and annexin A5 are also found in T-tubules in longitudinal sections of human skeletal muscle. Figure 5A-D shows the co-localization of POPDC1 and 2, XIRP1 and annexin A5 at Z-lines/T-tubules in longitudinal sections of human skeletal muscle, using a monoclonal antibody against alpha-actinin to locate Z-lines (Fig. 5C). POPDC1, POPDC2 and XIRP1 are present at the sarcolemma in transverse sections of human skeletal muscle

POPDC1 (Fig. 6A) and POPDC2 (Fig. 6B) are present at the sarcolemma in transverse sections of skeletal muscle. XIRP1 is also present at the sarcolemma, but does not co-localise exactly with POPDC proteins. Figures 5 and 6 show representative images of sections from five different human muscle biopsies which gave similar results in these experiments. Discussion Our initial proteomic study of pull-downs from human myotube extracts using recombinant POPDC1 or POPDC2 as the bait proteins showed that XIRP1 and cardiac actin were the highest-scoring hits (Table 1). XIRP1 has been located to myotendinous junctions in skeletal muscle [22] and to intercalated discs in heart [31,32]. We examined the interacting POPDC and XIRP1 proteins further by co-localization studies, bearing in mind that POPDC proteins (via zonula occludens-1, ZO-1 [33]) and XIRP1 [34] are known actin-binding proteins. Proteomic studies only give clues to potential interaction partners in a single tissue; Table 1 should not be expected to include all interaction partners of POPDC proteins and all potential partners revealed by proteomics require biochemical confirmation. XIRP1 is a scaffolding protein which stabilizes and maintains the integrity of the intercalated disc. Overexpression of XIRP1 leads to increased phosphorylation of gap junction protein alpha 1 (GJA1), also known as connexin-43 (Cx43), and down-regulation of the activity of gap junctions in cardiomyocytes [35]). Knockout Xirp1 (or Xin α) mice develop

a late-onset cardiomyopathy with conduction defects [22]. Missense mutations in *XIRP1*

cause cardiac arrhythmias [24] and a number of mutations in XIRP1 that are classed as

deleterious have been found in patients with conditions including myalgia, proximal weakness, arrhythmia, contractures and cardiac conduction defects [11]. XIRP proteins have binding domains for both β -catenin and p120-catenin, through which they attach to Ncadherin in the intercalated disc [36]. Intercalated discs provide electromechanical coupling between adjacent cardiomyocytes. Depolarising current is transmitted via gap junctions in the intercalated discs from cell to cell across the heart. Defects in intercalated disc structure and function are common features of cardiomyopathy, arrhythmia and heart failure (Reviewed: [32]).

Annexin A5 was also a significant interaction partner for POPDC1/2 (Table 1). Annexin A5 is upregulated in failing human hearts [37] and has been linked to plasma membrane repair in skeletal muscle [38]), a property shared with caveolin-3 [39] and XIRP1 [40,41]. There is some evidence of a role for POPDC1 in the maintenance of plasma membrane integrity, as discontinuities in the plasma membrane have been observed in the grandfather of a family carrying a pathogenic homozygous missense mutation (c.602C>T, p.S201F) in POPDC1 [9]. This mutation gives rise to cardiac arrhythmia and LGMD [9].

Using new monoclonal antibodies (mAbs) specific for either POPDC1 (1C12) or POPDC2 (12G12), together with a commercial polyclonal antibody against POPDC1 (BVES), we have shown that both proteins are localized at the sarcolemma and transverse tubules in both human skeletal muscle and rat heart muscle. Partial co-localization of both XIRP1 and annexin A5 with POPDC1/2 was found in T-tubules in both cardiac and skeletal muscles. Ttubules are invaginations in the plasma membrane of muscle cells, they are rich in ion channels and relay excitation signals from the external membrane to the interior of cells and are very sensitive to eccentric stretch (reviewed in [42]). In heart, POPDC1 and 2 are mainly concentrated at intercalated discs. The presence of POPDC2 at intercalated discs was also

shown by Soni *et al.* 2016 [29], while co-localization of XIRP1 with Connexin-43 at the intercalated discs of human hearts was shown by Xie *et al.* 2017 [35]. There is considerable functional overlap between POPDC proteins and XIRP proteins. In null mutants in the mouse for both gene families, a cardiac arrhythmia phenotype was reported. In the case of *Popdc1* and *Popdc2* null mutants, a stress-induced bradycardia phenotype was observed, while at baseline the mutants showed a normal rhythmic beating heart [7]. In contrast, the *Xirp1* null mutant phenotype was characterized by a late-onset cardiomyopathy with conduction defects [22]. In patients, *POPDC1* has also been linked to prolonged QT interval, increased QRS duration and AV-block [43]. Interestingly, in contrast to the mouse, in zebrafish, an AV-block was seen for both *popdc1* and *popdc2* null mutants [9,17,43]. Functional overlap is also seen in *Popdc1* null mutants and *Xirp1* null mutants with regard to skeletal muscle regeneration [25,26].

Although the pull-downs clearly show association between POPDC1/2, XIRP1 and actin, it is not clear whether the interactions between them are direct or indirect. It is possible that they are members of a larger complex, some components of which were not detected by proteomics for technical reasons. Earlier studies identified different binding partners for POPDC proteins, including dystrophin, dysferlin, VAMP2, VAMP3, GEFT, NDRG4, ZO-1, TREK-1; c-MYC and PR61-alpha, as well as caveolin-3 [7,8,9,33,44, ,45,46,47]. None of these are among the 340 proteins that appeared in our pull-downs, although this may be either because their interaction with POPDC is tissue-specific and not occurring in myotube cultures, or because they are present at very low levels in RIPA buffer extracts, or because they showed some non-specific binding to our control beads or because the interaction does not withstand the extraction conditions used. With numerous other interaction partners having been linked to POPDC1 and POPDC2 [4], it is not surprising that the loss-of-

function phenotype of *POPDC1* and *XIRP1* would show only partial overlap. In this context it
will also be interesting to define all components of the protein complex of which XIRP1 and
POPDC1 are part in cardiac and skeletal muscle.

5 Conclusions

POPDC1/2 proteins interact with XIRP1 in skeletal myotubes and adult heart. Determining
the functions of POPDC and XIRP1 proteins at the t-tubules and intercalated discs will help
to understand their roles in normal cardiac and skeletal muscle function and what can go
wrong when pathogenic mutations occur.

11 Methods

Popdc "bait" proteins

GST fusion proteins were generated by PCR from cDNA for mouse Popdc1 (Bves, GenBank Accession No: NM 024285.3) (aa 114 to 357) and for mouse Popdc2 (GenBank Accession No: NM_001081984.2) (aa 98 to 339) and cloned into pGEX bacterial expression vectors. pGEX constructs were transformed into E. coli BL21(DE3) and induced by IPTG to give GST-fusion proteins. After incubation the cells were washed with TNE buffer and sonicated sequentially with TNE, 2M urea, 4M urea, 6M urea and 8M urea. A Coomassie Blue-stained gel showed that most of the recombinant proteins were extracted in the 6M urea fraction and these were used in subsequent work. Resultant proteins contained GST-tags followed by the intracellular C-terminal tails of mPopdc1 and mPopdc2, with molecular weights of 58.3 and 57.1 kDa respectively.

Muscle cell culture and extraction

A clonal immortalized human myoblast cell line from a 25-year-old control donor without neuromuscular disease was immortalized by transduction with human telomerase reverse transcriptase (hTERT) and cyclin-dependent kinase-4 (Cdk4) containing retroviral vectors, at the Institut de Myologie, Paris, as described previously [48]. Myoblasts were cultured in skeletal muscle cell growth medium (Cat No: C-23060; PromoCell GmbH, Heidelberg, Germany) containing supplement mix (Cat No: C-39365; PromoCell) with 20% Fetal Bovine Serum (Cat No: 10270; Gibco; ThermoFisher Scientific, Paisley, UK). At around 80% confluency, adherent myoblasts were washed in medium lacking serum and differentiation was induced by culturing in DMEM (Cat No: 31966-021; Gibco; ThermoFisher Scientific) supplemented with Insulin (1721nM), Transferrin (68.7nM), Selenium (38.7nM) (ITS-X; Cat No: 51500-056; Gibco; ThermoFisher Scientific) and Penicillin-Streptomycin (Cat No: DE17-603E; Lonza; Verviers, Belgium). Cell culture was continued for a further 4 days, after which, over 80% of the cells had fused into myotubes. Cells cultured for pull-down experiments were extracted in RIPA buffer (1% NP-40; 0.25% sodium deoxycholate; 1mM EDTA; 150mM NaCl; 50mM Tris-HCl, pH7.4), in the presence of protease inhibitor cocktail (Sigma P8340; plus 1mM PMSF), by sonication followed by centrifugation.

Pull-down and proteomic analysis

Glutathione high capacity magnetic agarose beads (Cat No: G0924; Sigma) were loaded with GST-tagged "Bait" proteins (mouse Popdc1 or mouse Popdc2 or No Bait control) and then incubated with "Prey" (cultured human myotube extract in RIPA buffer). Captured proteins were digested by overnight incubation with 1% v/v trypsin at 37°C and analysed by nanoLC MSMS as described previously [49]. The MS/MS data file was analysed using the

Mascot search algorithm (Matrix Science), against the SwissProt database (May 2018) restricting the search to Homo sapiens (557,491 sequences), with trypsin as the cleavage enzyme and methionine oxidation as a variable modification. The peptide mass tolerance was set to 20 ppm and the MSMS mass tolerance to \pm 0.1 Da. This analysis was used to identify medium-to-high abundance proteins that interact with Popdc proteins.

7 Extraction and pull-down with rat heart

The heart was removed from a five month old Sprague Dawley rat that had been euthanized humanely with an overdose of pentobarbitone anesthetic (0.5 mL/100g) via intraperitoneal injection. The heart was cut in half, washed with PBS, weighed, snap frozen in liquid nitrogen, transferred to a mortar and ground to a fine powder with a pestle. The liquid nitrogen was then allowed to evaporate and the powdered heart extracted with RIPA buffer with inhibitors (around 1mL buffer per 100 µg heart), sonicated on ice and centrifuged. The heart extract was used instead of myotube extract as "prey", with mouse GST- mouse Popdc1 on glutathione beads as "bait", as described earlier.

17 Hybridoma production

Human *BVES* (*POPDC1*) cDNA (GenBank Accession No: NM_007073.4) was a gift from D. Bader [33]. A GST fusion protein was generated by PCR from the C-terminal tail of (aa 115-347) and cloning into the pGEX bacterial expression vector. This protein contains the GST tag followed by the intracellular C-terminal tail of POPDC1. The GST-mouse Popdc2 (amino acids 98 to 339) fusion protein preparation described earlier for pull-down experiments was also used for hybridoma production. Induction and extraction of the GSTfusion proteins was performed as described earlier. Fusion protein in the 6M urea fraction

was used in the protocol for immunization of BALB/c mice for production of mAbs [28]. ELISA plates were coated with either the GST-POPDC fusion proteins used for immunization or with an unrelated GST-fusion protein, in order to eliminate those mAbs reacting with the GST fusion tag. Hybridoma supernatants were additionally screened for colocalisation with transfected human POPDC proteins as follows: COS-7 cells were transfected (Lipofectamine) with either hPOPDC1-YFP or hPOPDC2-FLAG tag on glass coverslips and the cells fixed and permeabilized with 50% acetone : 50% methanol. Transfected cells were incubated with hybridoma supernatants. hPOPDC1-YFP transfectants were then incubated with 5µg/ml goat anti-mouse ALEXA 546 (Cat No: A11030, Molecular Probes, Eugene, Oregon, USA). hPOPDC2-FLAG transfectants were incubated with rabbit oligoclonal anti-FLAG (Cat No: 740001, Thermo Fisher Scientific), followed by 5µg/ml goat anti-mouse ALEXA 546 (Cat No: A11030, Molecular Probes) and 5µg/ml goat anti-rabbit ALEXA 488 (Cat No: A11034, Molecular Probes). DAPI was added for the final 10 minutes of incubation to counterstain nuclei and confocal microscopy performed as described below. Monoclonal antibodies POPDC1-1C12 and POPDC2-12G12 were selected because they recognized the corresponding transfected POPDC1 or POPDC2 by immunofluorescence microscopy (Fig.2). SDS-polyacrylamide gel electrophoresis and Western blotting Samples were mixed with SDS buffer (125mM Tris pH 6.8; 2% SDS; 5% 2-beta mercaptoethanol; 5% glycerol; with bromophenol blue), boiled and subjected to SDS-PAGE using 10% polyacrylamide gels and transferred to nitrocellulose membranes (Protan BA85, Whatman). Non-specific sites were blocked with 5% skimmed milk protein, membranes

23 washed with PBS and then incubated with primary monoclonal antibodies against: XIRP1,

24 (Xin-alpha (D-8); sc-166658; 1/100; Santa Cruz Biotechnology, Insight Biotechnology Ltd.,

Wembly, UK); or GST, mAb 17A10 (1/100). This was followed by washing in PBS and incubation with secondary antibody (peroxidase-labelled rabbit anti-mouse immunoglobulins, P0260; 1/1000; Dako, Denmark). Alternatively, for the detection of annexin A5, pAb ANXA5 (Abcam; ab14196; 1.4 μ g/mL) primary antibody followed by goat anti-rabbit Ig HRP (P0448; Dako; 1/1000) secondary antibody. All antibodies were diluted in PBS containing 0.05% Triton X, 0.1% BSA, 1% horse serum and 1% fetal bovine serum. XIRP1 and annexin A5 antibody reacting bands were detected with SuperSignal West Femto chemiluminescent reagent (Cat No: 34094; ThermoFisher Scientific) and GST antibody reacting bands were detected with SuperSignal West Pico Plus chemiluminescent reagent (Cat No: 34580; ThermoFisher Scientific) and visualized with a ChemiDoc Touch imaging system (BioRad Ltd.). Immunoprecipitation Superparamagnetic polystyrene beads coated with monoclonal human anti-mouse IgG antibodies (Dynabeads Pan Mouse IgG; Cat No: 11041; Invitrogen) were washed with 4% bovine serum albumin in PBS and then incubated with mAb against POPDC1 (1C12) or PBS alone (control) with gentle rolling for 1 hour. Beads were then washed with PBS and incubated with cultured human myotube extract in RIPA buffer for 1 hour with rolling. Beads were thoroughly washed with RIPA buffer and then with PBS and the bound material extracted by incubating the beads with SDS sample buffer, heating to 90°C for 3 minutes and then saving the eluate. Immunofluorescence microscopy

1	Immunohistochemistry was performed on unfixed cryostat sections of human
2	skeletal muscle or rat cardiac ventricle muscle. Monoclonal antibodies against: POPDC1
3	(1C12; 1:2 dilution); POPDC2 (12G12 1:2 dilution); XIRP1 (Xin-alpha (D-8); sc-166658; 1/100);
4	Actinin (MANDYS 141, 4A12, 1/50 [50]); were diluted and incubated on specimens for 1
5	hour. Specimens were then washed and incubated with rabbit polyclonal antibodies:
6	Connexin 43 (GJA1, ab11370; Abcam; 1/1000); XIRP1 (Cat No: HPA016750; Sigma; 1/150);
7	Annexin A5 (ANXA5, ab14196; Abcam; 1/50); POPDC1/BVES (Cat No: STJ22848; St. John's
8	Laboratory Ltd., London, E16 2RD; 1/100); Amphiphysin (ab244375; Abcam; 1/50). Following
9	antibody incubation, specimens were washed with PBS and then incubated with 5 μ g/ml of
10	each secondary antibody which was either: goat anti-mouse ALEXA 488 (Cat No: A11029,
11	Molecular Probes, Eugene, Oregon, USA) and goat anti-rabbit ALEXA 546 (Cat No: A11010),
12	or with goat anti-mouse ALEXA 546 (Cat No: A11030) and goat anti-rabbit ALEXA 488 (Cat
13	No: A11034). Secondary antibodies were diluted in PBS containing 1% horse serum, 1% fetal
14	bovine serum and 0.1% BSA, and incubated on the sections for 1 hour. Alternatively, for
15	some immunofluorescent labelling of rat cardiac sections, a single primary (polyclonal)
16	antibody then single goat anti-rabbit ALEXA 546 antibody were used, followed by ALEXA-488
17	Phalloidin (Cat No: A12383; Molecular Probes) to label filamentous actin. Finally, DAPI (at
18	200 ng/ml) was added for the final 10 minutes of incubation to counterstain nuclei before
19	washing and mounting the sections (Hydromount; Merck). For protein competition
20	experiments, monoclonal antibodies were pre-incubated with 1mg/mL recombinant protein
21	preparation for 1 h at room temperature. Images were acquired by sequential scanning with
22	a Leica TCS SP5 spectral confocal microscope (63x/1.4 oil) (Leica Microsystems, Milton
23	Keynes, UK), except the human heart images which were acquired with a Zeiss LSM780
24	confocal microscope (40x/1.3 oil) (Carl Zeiss Microscopy, Cambridge, UK.

1	1		
1 2 3	2	Abbreviation	5
4 5 6	3		
7 8 9	4	ACTC1	- Actin Alpha Cardiac Muscle 1
10 11	5	ANXA5	- Annexin A5
12 13 14	6	AV	- Atrioventricular
15 16 17	7	BIN1	- Bridging Integrator 1 (also known as Amphiphysin II)
18 19	8	BSA	- Bovine serum albumin
20 21 22	9	BVES	- Blood Vessel Epicardial Substance (also known as POPDC1)
23 24 25	10	cAMP	- Cyclic adenosine 3',5'-monophosphate
25 26 27	11	CAV3	- Caveolin 3
28 29 30	12	Cdk4	- Cyclin-dependent kinase-4
31 32	13	CMYA1	- Cardiomyopathy-associated protein 1 (also known as XIRP1)
33 34 35	14	с-Мус	- MYC Proto-Oncogene
36 37	15	COS-7	- African green monkey kidney fibroblast-like cell line
38 39 40	16	Cx43	- Connexin-43 (Gap Junction Protein Alpha 1, GJA1)
41 42 43	17	DAPI	- 4',6-diamidino-2-phenylindole
44 45	18	FLAG	- DYKDDDDK-tag
46 47 48	19	FRET	- Fluorescence resonance energy transfer
49 50	20	GEFT	- ARHGEF25 or Rho Guanine Nucleotide Exchange Factor 25
51 52 53	21	GST	- Glutathione S-transferase
54 55 56	22	HEK293	- Human embryonic kidney 293 cells
57 58	23	HRP	- Horseradish peroxidase
59 60 61	24	hTERT	- Human telomerase reverse transcriptase
62 63 64			18

1	1	IPTG	- Isopropyl beta-D-1-thiogalactopyranoside
2 3	2	KCNK2	- Potassium Two Pore Domain Channel Subfamily K Member 2 (TREK-1)
4 5 6	3	LGMD	- Limb-girdle muscular dystrophy
7 8 9	4	mAb	- Monoclonal antibody
10 11	5	Nano LC MSM	1S - Nanoscale liquid chromatography tandem mass spectrometry
12 13 14	6	NDRG4	- NDRG Family Member 4 (or N-Myc Downstream-Regulated Gene 4 Protein)
15 16 17	7	pAb	- Polyclonal antibody
18 19	8	PBS	- Phosphate-buffered saline
20 21 22	9	PCR	- Polymerase chain reaction
23 24	10	PMSF	- Phenylmethylsulfonyl fluoride (serine protease inhibitor)
25 26 27	11	POPDC1	- Popeye Domain Containing 1 (also known as BVES)
28 29 20	12	POPDC2	- Popeye Domain Containing 2
31 32	13	POPDC3	- Popeye Domain Containing 3
33 34 35	14	PR61-alpha	- Protein Phosphatase 2 Regulatory Subunit B'Alpha (PPP2R5A)
36 37	15	RIPA buffer	- Radioimmunoprecipitation assay buffer
38 39 40	16	SDS	- Sodium dodecyl sulfate
41 42	17	TNE buffer	- Tris-HCl, NaCl, EDTA buffer
44 45	18	TREK-1	- Potassium Two Pore Domain Channel Subfamily K Member 2 (KCNK2)
46 47 48	19	VAMP2	- Vesicle Associated Membrane Protein 2
49 50	20	VAMP3	- Vesicle Associated Membrane Protein 3
51 52 53	21	XIRP1	- Xin Actin Binding Repeat Containing 1 (also known as XIN or CMYA1 -
54 55	22		cardiomyopathy-associated protein 1)
56 57 58	23	YFP	- Yellow fluorescent protein
59 60	24	ZO-1	- Zonula Occludens Protein 1 (Tight Junction Protein 1)
o⊥ 62 63			19
54 55			

Declarations: б Ethics approval and consent to participate This study has been approved by the RJAH Orthopaedic Hospital Research Committee. All human tissue studies complied with the UK Human Tissue Act (2006). Human biopsies were obtained following written informed consent using protocols approved by RJAH Orthopaedic Hospital, Oswestry and Imperial College, London. Muscle biopsy for immortalisation of myoblasts was obtained with written informed consent in accordance with European recommendations and French legislation. All animal procedures were performed with approval of the Keele University Animal Welfare and Ethical Review Body and conducted under the licensed authority of the UK Home Office. **Consent for publication** Not applicable. Availability of data and materials Mass spectrometry data generated and analysed during this study are available in the supplementary information file, "Additional file 1". Monoclonal antibodies are available from the MDA monoclonal antibody resource, Oswestry, UK: http://www.glennmorris.org.uk/mabs/WCIND.htm POPDC nucleotide sequence data is available from GenBank with the following accession numbers: Mouse Bves/Popdc1 (NM 024285.3); Mouse Popdc2 (NM 001081984.2); Human BVES/POPDC1 (NM 007073.4); Human POPDC2 (NM 022135.4).

Competing interests

The authors declare that they have no competing interests

Funding

This work was supported by the British Heart Foundation [PG/16/68/31991; to IH and GEM], [PG/19/13/34247; to TB] and the Orthopaedic Institute Ltd., RJAH Orthopaedic Hospital, Oswestry UK [RPG141 and RPG189; to IH and GEM]. Infrastructure support for this research was provided by the NIHR Imperial Biomedical Research Centre (BRC) and ICHTB. The funding bodies played no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Authors' contributions

Conceived and designed the experiments: IH, HRF, TB, GEM; Hybridoma production: IH, GEM; Mass spectrometry: HRF, SLS; Other experimental work: IH, HRF, RFRS, GEM; Analysed the data: IH, HRF, RFRS, SLS, TB, GEM; Wrote the manuscript: IH, HRF, TB, GEM. All authors read and approved the final manuscript.

Acknowledgements

We thank The Platform for Immortalization of Human Cells, Institute of Myology, Paris, France, for the immortalized human myoblasts. We thank Ursula Herbort-Brand for expert technical assistance. The donation of the POPDC1-GST construct by David Bader is gratefully acknowledged.

1	
2	References
3	
4	1/ Reese, D.E., Zavaljevski, M., Streiff, N.L. and Bader, D. Bves: A novel gene expressed
5	during coronary blood vessel development. Dev. Biol. 1999; 209 ,159-171.
6	
7	2/ Andrée, B., Hillemann, T., Kessler-Icekson, G., Schmitt-John, T., Jockusch, H., Arnold,
8	H.H. and Brand, T. Isolation and characterization of the novel popeye gene family expressed
9	in skeletal muscle and heart. Dev. Biol. 2000; 223 ,371-382.
10	
11	3/ Brand, T., Simrick, S.L., Poon, K.L. and Schindler, R.F. The cAMP-binding Popdc
12	proteins have a redundant function in the heart. Biochem. Soc. Trans. 2014;42,295-301.
13	
14	4/ Brand, T. POPDC proteins and cardiac function. Biochem. Soc. Trans. 2019;47,1393-
15	1404.
16	
17	5/ Knight, R.F., Bader, D.M. and Backstrom, J.R. Membrane topology of Bves/Pop1A, a
18	cell adhesion molecule that displays dynamic changes in cellular distribution during
19	development. J. Biol. Chem. 2003; 278 ,32872-32879.
20	
21	6/ Vasavada, T.K., DiAngelo, J.R. and Duncan, M.K. Developmental expression of
22	Pop1/Bves. J. Histochem. Cytochem. 2004; 52 ,371-377.
23	

1	7/ Froese, A., Breher, S.S., Waldeyer, C., Schindler, R.F., Nikolaev, V.O., Rinné, S.,
2	Wischmeyer, E., Schlueter, J., Becher, J., Simrick, S., et al. Popeye domain containing
3	proteins are essential for stress-mediated modulation of cardiac pacemaking in mice. J. Clin.
4	Invest. 2012; 122 ,1119-1130.
5	
6	8/ Alcalay, Y., Hochhauser, E., Kliminski, V., Dick, J., Zahalka, M.A., Parnes, D.,
7	Schlesinger, H., Abassi, Z., Shainberg, A., Schindler, R.F., et al. Popeye domain containing 1
8	(Popdc1/Bves) is a caveolae-associated protein involved in ischemia tolerance. PLoS One
9	2013; 8 ,e71100.
10	
11	9/ Schindler, R.F., Scotton, C., Zhang, J., Passarelli, C., Ortiz-Bonnin, B., Simrick, S.,
12	Schwerte, T., Poon, K.L., Fang, M., Rinné, S., et al. POPDC1 (S201F) causes muscular
13	dystrophy and arrhythmia by affecting protein trafficking. J. Clin. Invest. 2016; 126 ,239-253.
14	
15	10/ De Ridder, W., Nelson, I., Asselbergh, B., De Paepe, B., Beuvin, M., Ben Yaou, R.,
16	Masson, C., Boland, A., Deleuze, J.F., Maisonobe, T., et al. Muscular dystrophy with
17	arrhythmia caused by loss-of-function mutations in BVES. Neurol. Genet. 2019; 5 ,e321.
18	
19	11/ Meinke, P., Kerr, A.R.W., Czapiewski, R., de Las Heras, J.I., Dixon, C.R., Harris, E.,
20	Kölbel, H., Muntoni, F., Schara, U., Straub, V., et al. A multistage sequencing strategy
21	pinpoints novel candidate alleles for Emery-Dreifuss muscular dystrophy and supports gene
22	misregulation as its pathomechanism. EBioMedicine 2020; 51 ,102587.
23	
	23

12/ Rinné, S., Ortiz-Bonnin, B., Stallmeyer, B., Schindler, R.F.R., Kiper, A.K., Dittmann, S., Friedrich, C., Zumhagen, S., Simrick, S.L., González, W., et al. Conduction disorder caused by a mutation in POPDC2, a novel modulator of the cardiac sodium channel SCN5A. Acta. Physiol. 2016;216(S707),0S03-1 (Abstract). Vissing, J., Johnson, K., Töpf, A., Nafissi, S., Díaz-Manera, J., French, V.M., Schindler, 13/ R.F., Sarathchandra, P., Løkken, N., Rinné, S., et al. POPDC3 Gene Variants Associate with a New Form of Limb Girdle Muscular Dystrophy. Ann. Neurol. 2019;86,832-843. 14/ Gingold-Belfer, R., Bergman, M., Alcalay, Y., Schlesinger, H., Aravot, D., Berman, M., Salman, H., Brand, T. and Kessler-Icekson, G. Popeye domain-containing 1 is down-regulated in failing human hearts. Int. J. Mol. Med. 2011;27,25-31. 15/ Andrée, B., Fleige, A., Arnold, H.H. and Brand, T. Mouse Pop1 is required for muscle regeneration in adult skeletal muscle. Mol. Cell. Biol. 2002;22,1504-1512. 16/ Boukens, B.J. and Christoffels, V.M. Popeye proteins: muscle for the aging sinus node. J. Clin. Invest. 2012;122,810-813. 17/ Kirchmaier, B.C., Poon, K.L., Schwerte, T., Huisken, J., Winkler, C., Jungblut, B., Stainier, D.Y. and Brand, T. The Popeye domain containing 2 (popdc2) gene in zebrafish is required for heart and skeletal muscle development. Dev. Biol. 2012;363,438-450. Erratum in: Dev. Biol. **366**,433.

18/ Schindler, R.F., Scotton, C., French, V., Ferlini, A. and Brand, T. The Popeye Domain Containing Genes and their Function in Striated Muscle. J. Cardiovasc. Dev. Dis. 2016;3,22. 19/ Parang, B., Thompson, J.J. and Williams, C.S. Blood Vessel Epicardial Substance (BVES) in junctional signaling and cancer. Tissue Barriers 2018;6,1-12. 20/ Han, P., Lei, Y., Li, D., Liu, J., Yan, W. and Tian, D. Ten years of research on the role of BVES/ POPDC1 in human disease: a review. Onco. Targets Ther. 2019;12,1279-1291. Amunjela, J.N., Swan, A.H. and Brand, T. The role of the Popeye domain containing 21/ gene family in organ homeostasis. Cells 2019;8,1594. 22/ Feng, H.Z., Wang, Q., Reiter, R.S., Lin, J.L., Lin, J.J. and Jin, J.P. Localization and function of Xinα in mouse skeletal muscle. Am. J. Physiol. Cell. Physiol. 2013;**304**,C1002-1012. 23/ Wang, Q., Lin, J.L., Chan, S.Y. and Lin, J.J. The Xin repeat-containing protein, mXinβ, initiates the maturation of the intercalated discs during postnatal heart development. Dev. Biol. 2013;374,264-280. Huang, L., Wu, K.H., Zhang, L., Wang, Q., Tang, S., Wu, Q., Jiang, P.H., Lin, J.J., Guo, J., 24/ Wang, L., et al. Critical Roles of Xirp Proteins in Cardiac Conduction and Their Rare Variants Identified in Sudden Unexplained Nocturnal Death Syndrome and Brugada Syndrome in Chinese Han Population. J. Am. Heart Assoc. 2018;7,e006320.

2	25/ Al-Sajee, D., Nissar, A.A., Coleman, S.K., Rebalka, I.A., Chiang, A., Wathra, R., van de	r
3	Ven, P.F., Orfanos, Z. and Hawke, T.J. Xin-deficient mice display myopathy, impaired	
4	contractility, attenuated muscle repair and altered satellite cell functionality. Acta. Physiol	•
5	(Oxf). 2015; 214 ,248-260.	
6		
7	26/ Nissar, A.A., Zemanek, B., Labatia, R., Atkinson, D.J., van der Ven, P.F., Fürst, D.O. a	nd
8	Hawke, T.J. Skeletal muscle regeneration is delayed by reduction in Xin expression:	
9	consequence of impaired satellite cell activation? Am. J. Physiol. Cell. Physiol.	
10	2012; 302 ,C220-227.	
11		
12	27/ Alonso, S., Garner, I., Vandekerckhove, J. and Buckingham, M. Genetic analysis of th	٦e
13	interaction between cardiac and skeletal actin gene expression in striated muscle of the	
14	mouse. J. Mol. Biol. 1990; 211 ,727-738.	
15		
16	28/ Nguyen, T.M. and Morris, G.E. A rapid method for generating large numbers of high	۱-
17	affinity monoclonal antibodies from a single mouse. In Walker, JM (ed), The Protein	
18	Protocols Handbook, 3rd Edition. Totowa NJ, Humana Press. 2010;1961-1974.	
19		
20	29/ Soni, S., Raaijmakers, A.J., Raaijmakers, L.M., Damen, J.M., van Stuijvenberg, L., Vos	,
21	M.A., Heck, A.J., van Veen, T.A. and Scholten, A. A Proteomics Approach to Identify New	
22	Putative Cardiac Intercalated Disk Proteins. PLoS One 2016;11,e0152231.	
23		

1	30/ Benevolensky, D., Belikova, Y., Mohammadzadeh, R., Trouvé, P., Marotte, F., Russo-
2	Marie, F., Samuel, J.L. and Charlemagne, D. Expression and Localization of the Annexins II,
3	V, and VI in Myocardium From Patients With End-Stage Heart Failure. Lab. Invest.
4	2000; 80 ,123-133.
5	
6	31/ Palatinus, J.A. and Gourdie, R.G. Xin and the art of intercalated disk maintenance.
7	Am. J. Physiol. Heart Circ. Physiol. 2007; 293 ,H2626-2628.
8	
9	32/ Wang, Q., Lin, J.L., Erives, A.J., Lin, C.I. and Lin, J.J. New insights into the roles of Xin
10	repeat-containing proteins in cardiac development, function, and disease. Int. Rev. Cell. Mol.
11	Biol. 2014; 310 :89-128.
12	
13	33/ Osler, M.E., Chang, M.S. and Bader, D.M. Bves modulates epithelial integrity through
14	an interaction at the tight junction. J. Cell Sci.2005; 118 ,4667-4678.
15	
16	34/ Jung-Ching Lin, J., Gustafson-Wagner, E.A., Sinn, H.W., Choi, S., Jaacks, S.M., Wang,
17	D.Z., Evans, S. and Li-Chun Lin, J. Structure, Expression, and Function of a Novel Intercalated
18	Disc Protein, Xin. J. Med. Sci. 2005; 25 ,215-222.
19	
20	35/ Xie, Y., Liu, S., Hu, S. and Wei, Y. Cardiomyopathy-Associated Gene 1-Sensitive PKC-
21	Dependent Connexin 43 Expression and Phosphorylation in Left Ventricular Noncompaction
22	Cardiomyopathy. Cell. Physiol. Biochem. 2017;44,828-842.
23	
	27

1
2
3
4
5
5
0
/
8
9
10
11
12
13
14
15
16
17
1/
T8
19
20
21
22
23
24
25
26
20
27
28
29
30
31
32
33
34
35
36
27
20
20
39
40
41
42
43
44
45
46
47
48
10
49
50
51
52
53
54
55
56
57
58
59
50
00 61
01
¢∠
63
64
65

1	36/ War	ng, Q., Lu, T.L., Adams, E., Lin, J.L. and Lin, J.J.Intercalated disc protein, mXin α ,		
2	suppresses p120-catenin-induced branching phenotype via its interactions with p120-			
3	catenin and cortactin. Arch. Biochem. Biophys. 2013; 535 ,91-100.			
4				
5	37/ Mat	teo, R.G. and Moravec, C.S. Immunolocalization of annexins IV, V and VI in the		
6	failing and r	non-failing human heart. Cardiovasc. Res. 2000; 45 ,961-970.		
7				
8	38/ Carr	neille, R., Bouvet, F., Tan, S., Croissant, C., Gounou, C., Mamchaoui, K., Mouly, V.,		
9	Brisson, A.R. and Bouter, A. Membrane repair of human skeletal muscle cells requires			
10	Annexin-A5. Biochim. Biophys. Acta. 2016; 1863 ,2267-2279.			
11				
12	39/ Cai,	C., Weisleder, N., Ko, J.K., Komazaki, S., Sunada, Y., Nishi, M., Takeshima, H. and		
13	Ma, J. Mem	brane repair defects in muscular dystrophy are linked to altered interaction		
14	between M	G53, caveolin-3, and dysferlin. J. Biol. Chem. 2009; 284 ,15894-15902.		
15				
16	40/ Haw	vke, T.J. , Atkinson, D.J., Kanatous, S.B., Van der Ven, P.F.M., Goetsch, S.C. and		
17	Garry, D.J. >	Kin, an Actin Binding Protein, Is Expressed Within Muscle Satellite Cells and		
18	Newly Rege	nerated Skeletal Muscle Fibers. Am. J. Physiol. Cell Physiol. 2007; 293, C1636-		
19	1644.			
20				
21	41/ Otte	en, C., van der Ven, P.F., Lewrenz, I., Paul, S., Steinhagen, A., Busch-Nentwich, E.,		
22	Eichhorst, J	., Wiesner, B., Stemple, D., Strähle, U., <i>et al</i> . Xirp proteins mark injured skeletal		
23	muscle in zebrafish. PLoS One 2012; 7 ,e31041.			
24				
		28		

1	42/ Cooper, S.T. and McNeil, P.L. Membrane Repair: Mechanisms and Pathophysiology.		
2	Physiol. Rev. 2015; 95 ,1205-1240.		
3			
4	43/ Wang, X., Tucker, N.R., Rizki, G., Mills, R., Krijger, P.H.L., de Wit, E., Subramanian, V.,		
5	Bartell, E., Nguyen, X.X., Ye, J., et al. Discovery and validation of sub-threshold genome-wide		
6	association study loci using epigenomic signatures. Elife. 2016; 5 ,10557.		
7			
8	44/ Hager, H.A., Roberts, R.J., Cross, E.E., Proux-Gillardeaux, V. and Bader, D.M.		
9	Identification of a novel Bves function: regulation of vesicular transport. EMBO J.		
10	2010; 29 ,532-545.		
11			
12	45/ Smith, T.K., Hager, H.A., Francis, R., Kilkenny, D.M., Lo, C.W. and Bader, D.M. Bves		
13	directly interacts with GEFT, and controls cell shape and movement through regulation of		
14	Rac1/Cdc42 activity. Proc. Natl. Acad. Sci. U.S.A. 2008; 105 ,8298-8303.		
15			
16	46/ Benesh, E.C., Miller, P.M., Pfaltzgraff, E.R., Grega-Larson, N.E., Hager, H.A., Sung,		
17	B.H., Qu, X., Baldwin, H.S., Weaver, A.M. and Bader, D.M. Bves and NDRG4 regulate		
18	directional epicardial cell migration through autocrine extracellular matrix deposition. Mol.		
19	Biol. Cell. 2013; 24 ,3496-3510.		
20			
21	47/ Parang, B., Kaz, A.M., Barrett, C.W., Short, S.P., Ning, W., Keating, C.E., Mittal, M.K.,		
22	Naik, R.D., Washington, M.K., Revetta, F.L. et al. BVES regulates c-Myc stability via PP2A and		
23	suppresses colitis-induced tumourigenesis. Gut. 2017; 66 ,852-862.		
24			

1	48/ Mamchaoui K, Trollet C, Bigot A, Negroni E, Chaouch S, Wolff A, Kandalla PK, Marie S,		
2	Di Santo J, St Guily JL, et al. Immortalized pathological human myoblasts: towards a		
3	universal tool for the study of neuromuscular disorders. Skelet. Muscle. 2011;1,34.		
4			
5	49/ Holt, I., Fuller, H.R., Lam, L.T., Sewry, C.A., Shirran, S.L., Zhang, Q., Shanahan, C.M.		
6	and Morris, G.E. Nesprin-1-alpha2 associates with kinesin at myotube outer nuclear		
7	membranes, but is restricted to neuromuscular junction nuclei in adult muscle. Sci. Rep.		
8	2019; 9 ,14202.		
9			
10	50/ Nguyen, T.M., Cartwright, A.J., Morris, G.E., Love, D., Bloomfield, J.F. and Davies, K.E.		
11	Monoclonal antibodies against defined regions of the muscular dystrophy protein,		
12	dystrophin. FEBS Lett. 1990; 262 ,237-240.		
13			
14			
15	Additional files		
16	Additional file 1: Table; Unfiltered mass spectrometry results; XLSX format.		
17	Additional file 2: Original uncropped blots used for Figure 1.		
18			
19			
20	Figure legends		
21			
22	Figure 1		
23	POPDC proteins pull down and co-immunoprecipitate XIRP1 and pull down annexin A5 from		
24	human myotube extract and pull-down XIRP1 from rat heart extract.		
	30		

Vertical black lines separate parts of blots that were originally non-adjacent or with different exposure times, as described. In all cases, the molecular weight markers in lane 1 were detected by colourimetric imaging and the antibody blots by chemiluminescence. (A) Glutathione beads were loaded with recombinant preparations of POPDC1, POPDC2 or not loaded (control for non-specific binding) and incubated with human myotube extracts. "Input" shows human myotube extracts diluted to 20%. Upper part of blot (A) (above 72kDa) developed with mAb against XIRP1 (sc-166658) shows XIRP1 in the input and pulled down by POPDC1 and POPDC2, but not in the control. The "Input" lane had a shorter exposure time than the others and the "Control" lane was originally non-adjacent. The Lower part of blot (A) (72kDa and below) was from a separate but identical gel to the upper blot and was developed with mAb against GST tag (17A10) as a loading control. The "Control" lane was originally non-adjacent to the others. The predicted molecular weights of the Popdc1 and Popdc2 fusion proteins are 58 and 57 kDa respectively, which are indicated approximately with a single arrow. (B) Pan mouse IgG Dynabeads were loaded with monoclonal antibody against POPDC1 (1C12) or not loaded (control) and incubated with human myotube extracts. The blot was developed with XIRP1 mAb and shows XIRP1 in the input and immunoprecipitated by the POPDC1 mAb but not by unloaded control beads. *=mouse Ig H and L chains. Predicted molecular weights of human XIRP1, isoforms 1,2 and 3 are 199, 122 and 56 kDa respectively. Arrows indicate bands corresponding to XIRP1 isoforms 1 and 3. The "POPDC1 1C12" lane had a longer exposure compared to the other lanes and the "molecular weight markers" were not originally in lane 1. (C) Glutathione beads loaded with recombinant POPDC1 or beads alone (control) were incubated with human myotube extract. The blot was developed with pAb against annexin A5. The arrow indicates a band corresponding to annexin A5 monomer (36 kDa) in the input and POPDC1

pull-down, but absent from the non-specific binding control. The "Control" lane was originally non-adjacent to the others. (D) Glutathione beads loaded with recombinant POPDC1 or beads alone (control) were incubated with rat heart ventricle extract. The blot was developed with mAb against XIRP1. Arrows indicate bands corresponding to XIRP1 in the input and POPDC1 pull-down, but absent from the non-specific binding control. The original colourimetric and chemiluminescence images are shown in Additional File 2. Figure 2 Immunofluorescent specificity of POPDC mAbs. COS-7 cells were transfected with either (A) human POPDC1 (GFP tag) or (B) human POPDC2 (FLAG tag) and processed as described in "methods". (A) mAb POPDC1 1C12 colocalised with transfected POPDC1 and (B) mAb POPDC2 12G12 colocalised with transfected POPDC2. The POPDC1 mAb did not recognise transfected POPDC2 and the POPDC2 mAb did not recognise transfected POPDC1 (not shown). (C) mAb POPDC1 1C12 localised to the sarcolemma of human skeletal muscle sections. This localisation was prevented by preincubation of the mAb with recombinant POPDC1 but not when the mAb was pre-incubated with recombinant POPDC2. Also, (D) localisation of mAb POPDC2 12G12 to the sarcolemma was not inhibited by pre-incubation with recombinant POPDC1 but this localisation was inhibited when the mAb was pre-incubated with recombinant POPDC2. Figure 3 Immunolocalisation of POPDC1/2, XIRP1 and annexin A5 at intercalated discs and myofibrils in rat heart.

1	In rat heart, intercalated discs were labelled with polyclonal antibody against connexin 43.
2	(A) XIRP1 colocalised with connexin 43 at the intercalated disc and also stained myofibrils.
3	(B) POPDC2 colocalises with XIRP1 at intercalated discs and myofibrils and POPDC2 is also
4	present at the cardiac sarcolemma. (C) POPDC1 is similar to POPDC2, locating to
5	intercalated discs and sarcolemma and to the centre of the actin I-band stained by
6	phalloidin. (D) Annexin A5 colocalises with XIRP1. (E) POPDC2 colocalises with amphiphysin
7	II, a marker for T-tubules.
8	
9	Figure 4
10	Immunolocalisation of POPDC1, POPDC2 and XIRP1 in human heart.
11	(A) POPDC1 and (B) POPDC2 were present at the sarcolemma and at intercalated discs.
12	XIRP1 was also present at intercalated discs. Cross striations were difficult to resolve in the
13	human heart sections.
14	
15	Figure 5
16	Immunolocalisation of POPDC1/2, XIRP1 and annexin A5 at myofibrils in longitudinal
17	sections of human skeletal muscle.
18	(A) POPDC1 and XIRP1, (B) POPDC2 and XIRP1, (C) XIRP1 and Actinin, (D) Annexin A5 and
19	XIRP1, all colocalised at myofibrils.
20	
21	Figure 6
22	Immunolocalisation of POPDC1/2 and XIRP1 in transverse section of human skeletal muscle.
23	(A) POPDC1 mAb 1C12 and (B) POPDC2 mAb 12G12 showed largely even and regular
24	sarcolemmal staining whereas XIRP1 was often discontinuous at the sarcolemma.
	33

	Accession number	Number of significant (unique) peptides	
		mPOPDC1	mPOPDC2
Actin, alpha cardiac muscle 1	ACTC_HUMAN	12	12
Xin actin-binding repeat-containing protein 1	XIRP1_HUMAN	11	19
Tubulin beta-4B chain	TBB4B_HUMAN	9	7
Tubulin beta chain	TBB5_HUMAN	9	6
Tubulin beta-6 chain	TBB6_HUMAN	9	4
ATP synthase subunit beta, mitochondrial	ATPB_HUMAN	7	5
Fructose-bisphosphate aldolase A	ALDOA_HUMAN	5	7
L-lactate dehydrogenase A chain	LDHA_HUMAN	5	5
Histone H2AX	H2AX_HUMAN	4	4
PDZ and LIM domain protein 3	PDLI3_HUMAN	3	5
Glucosidase 2 subunit beta	GLU2B_HUMAN	3	5
Lamina-associated polypeptide 2 (beta/gamma)	LAP2B_HUMAN	3	4
Histone H2A.V	H2AV_HUMAN	3	4
Phosphoglycerate mutase 1	PGAM1_HUMAN	3	4
Phosphoglycerate kinase 1	PGK1_HUMAN	3	4
Heat shock protein beta-2	HSPB2_HUMAN	3	4
Annexin A5	ANXA5_HUMAN	3	3
40S ribosomal protein S30	RS30_HUMAN	3	3

3 Table 1: Mass spectrometry abridged results of mouse POPDC1/2 pull-downs

 5 Only proteins that met the following criteria are shown: identified from more than two

6 significant unique peptide sequences, undetected in the control pull-down AND detected

7 across both mPOPDC1 and mPOPDC2 pull-downs. See Additional file 1 for full results.





















100 µm

Additional File 1

Click here to access/download Supplementary Material Additional file 1.xlsx Additional File 2

Click here to access/download Supplementary Material Additional File 2 - Uncropped Blots.pdf