

This work is protected by copyright and other intellectual property rights and duplication or sale of all or part is not permitted, except that material may be duplicated by you for research, private study, criticism/review or educational purposes. Electronic or print copies are for your own personal, noncommercial use and shall not be passed to any other individual. No quotation may be published without proper acknowledgement. For any other use, or to quote extensively from the work, permission must be obtained from the copyright holder/s. ASPECTS OF THE STRATIGRAPHY, AND AMMONITE FAUNAS OF THE AALENIAN - BAJOCIAN STAGES IN GREAT BRITAIN

BY

COLIN, F. PARSONS

A THESIS SUBMITTED FOR A DEGREE OF DOCTOR OF PHILOSOPHY

UNIVERSITY OF KEELE 1980





IMAGING SERVICES NORTH

Boston Spa, Wetherby West Yorkshire, LS23 7BQ www.bl.uk

BEST COPY AVAILABLE.

VARIABLE PRINT QUALITY



IMAGING SERVICES NORTH

Boston Spa, Wetherby West Yorkshire, LS23 7BQ www.bl.uk

The following has been redacted from this digital copy of the original thesis at the request of the awarding university:

7 publications/journal articles at the end of Volume 2 (after page 565) "Additions to fauna decrease the imperfection of the zoological, but increase that of any local geological record : the gaps caused by destruction stand revealed more plainly"

> S.S.BUCKMAN, 'Type Ammonites' Vol.III

"It is one of the greatest misfortunes for Jurassic geology that when increasing age and fraility prevented Buckman from continuing active field-work, he lost sight of the distinction between results obtained with hammer, collecting bag and field notebook, and those arrived at by speculation and deduction from matrices at home. The extent of the calamity is magnified by the fact that this loss of grasp of the importance of keeping surmise distinct from fact came when his reputation was at its highest, and the two kinds of results are almost inextricably interwoven in his later published works"

> W.J.ARKELL, 'The Jurassic System in Great Britain' p.36.



d'Orbigny's Bajocian : Les Hachettes, St.Honorine -des-Pertes, Normandy,France. ABSTRACT : A detailed stratigraphic revision of parts of the Aalenian-Bajocian Stages is undertaken. The status of the Sowerbyi zone is discussed and reasons given for its rejection and replacement by the Discites and Laeviuscula Zones. The Sauzei Zone is redefined. The detailed stratigraphy of the Humphriesianum/Subfurcatum Zone rocks of Southern England is described, its correlation discussed and a detailed analysis of their ammonite faunas given. The litho- and biostratigraphy of the Garantiana/Parkinsoni Zone rocks of the Sherborne area are revised. Formations and Members of the Cotswold Inferior Oolite Group are re-defined, and their Bajocian Ammonite faunas listed. The litho- and biostratigraphy of the Aalenian-Bajocian rocks of Dundry Hill, Bristol, are revised and the type horizons of Buckman's and Sowerbys' ammonite species located. The ammonite faunas of the Scarborough Formation of N.E.Yorkshire are revised and its lithostratigraphy described.

Two new Bajocian, microconch Otoitid species are described and their phylogenetic and stratigraphic significance discussed. A detailed systematic revision of the Sphaeroceratinae is given. Its constituent genera are re-defined and six species and subspecies of Sphaeroceras (S.), eight of <u>Chondroceras</u> (<u>C</u>.) ; including one new; and two subspecies of <u>Labyrinthoceras</u> are described. The evolution of the subfamily is discussed, its problems of sexual dimorphism summarised, and the stratigraphic distribution and significance of its members detailed.

Finally a summary of the stratigraphic part of this work is given in a discussion of the Bajocian standard zonal scheme. i ABSTRACT

ii INTRODUCTION

1. TOWARDS A REVISION OF THE STANDARD ZONAL SCHEME FOR	
THE BAJOCIAN AND PARTS OF THE AALENIAN STAGES	17
A.The Sauzei and so called Sowerbyi Zones of the	
Lower Bajocian (Middle Jurassic) of Europe.	18
B.A stratigraphic revision of the Humphriesianum/	
Subfurcatum Zone rocks of Southern England.	6 5
C.A stratigraphic revision of the Garantiana and	
Parkinsoni Zone (Upper Bajocian) rocks of the	125
Sherborne area, Dorset.	
2.A STRATIGRAPHIC REVISION OF THE BAJOCIAN ROCKS OF	
CERTAIN AREAS OF ENGLAND	167
A.A stratigraphic revision of the Aalenian-Bajocian	
rocks of Dundry Hill, Bristol.	168
B.A stratigraphic revision of the Bajocian (Jurassic)	
rocks of the Cotswold Hills.	215
C.A stratigraphic revision of the Scarborough Formation,	
Middle Jurassic , of north-east Yorkshire.	255
3.SYSTEMATIC DESCRIPTIONS OF MEMBERS OF THE BAJOCIAN	
AMMONITE SUPERFAMILY STEPHANOCERATACEA	302
A. Two new Bajocian microconch Otoitid ammonites and	
their significance.	303
B.A systematic revision of the Bajocian ammonite	
Subfamily Sphaeroceratinae.	335

B.i. Appendix. Dimensions of additional Sphaeroceratid material, not given in the main body of the text.

4. A STANDARD ZONAL SCHEME FOR THE BAJOCIAN STAGE

Publication Details

1A Published in Newsletters in Stratigraphy (1974), 3:153-179. 1B Shortened version published News.Strat.(1976),5:114-42. 2A Published in a modified form, Proc.Geol.Ass.(1979), 90: 133-51. 2B Submitted to Geol.Soc.Lond. 2C Shortened version published Proc.Yorks.geol.Soc.(1977), 43:203-22. 3A Published Palaeontology (1977), 20:101-18. The published versions of 2A &2C, together with copies of the following, are bound in at the rear of part 2 :i. Limonitic concretions from the European Jurassic .. Sedimentology, 1972. ii. The Bajocian stratigraphy of Eastern England. Proc. Yorks. geol.Soc. 1974. iii. Ammonites from the Doulting conglomerate Bed of Somerset. Palaeontology 1975. iv. The stratigraphy of the Stony Head Cutting. Proc.Dorset Nat.Hist.Arch.Soc. 1975. v. Ammonite evidence for dating some Inferior Oolite sections in the North Cotswolds. Proc.Geol. Ass. 1976.

2

498

<u>INTRODUCTION</u>

The rocks assigned to the Aalenian-Bajocian Stages in southern England, and in particular north Dorset, are some of the most highly fossiliferous beds in the world (Arkell, 1933, p.193). Over the years these beds have yielded a wealth of ammonites, which on one hand have provided the type material for numerous new genera and species (Smith, 1817; J.&J.de C.Sowerby, 1812-46; Buckman, 1909-30), and on the other they have provided the stratigraphic basis for much of the present standard zonal scheme. The original aim of this work was to have been a taxonomic revision of the ammonite Superfamily Stephanoceratacea, which forms an important part of these abundant Dorset faunas. At an early stage it became evident that before this primary aim could be accomplished, the stratigraphy of the Bajocian rocks of southern England would have to be revised. It was initially assumed that the standard zonal scheme for the Aalenian-Bajocian, as adopted by Spath (1936) and Arkell (1956), was a fair reflection of the stratigraphic distribution of the ammonite faunas. However, my research confirmed some already prevalent suspicions (Torrens, 1969, p. 302), that the Spath/Arkell scheme, based as it was on S.S.Buckman's scheme of hemerae (Buckman, 1893,1910,1909-30), was full of inconsistencies and in some cases outright errors. A high proportion of this thesis is thus taken up with stratigraphic problems, in first revising certain parts of the standard zonal scheme, and secondly in applying this scheme to some specific areas of English Bajocian rocks.



4

Figure 1

The main structural features controlling the deposition and subsequent preservation of the Middle Jurassic of the British Isles.

1. LITHOSTRATIGRAPHY

The original deposition, and to a certain extent the subsequent preservation of the British Middle Jurassic rocks has been controlled by important deep seated structural features. This has resulted in the thickest sequences being deposited in the centres of 'negative areas' or 'basins', whilst on the 'positive areas' or 'highs', only much attenuated or incomplete successions are preserved - see Figure 1 & Table 1. The most well known and frequently used lithostratigraphic term spanning the Aalenian-Bajocian, is the Inferior Oolite Group or 'series'. As originally defined (Townsend,1813), the use of this name should be restricted to those rocks on the western periphery of the Anglo-Paris Basin.

(a) The Anglo-Paris Basin

Whilst thick sequences of Aalenian-Bajocian rocks are preserved at the subsurface, towards the centre of this basin (e.g. Ashdown, No.2 borehole, see Gatrall, Jenkyns & Parsons, 1972, fig. 8), only the surface outcrops, with their relatively common ammonites, are of any real value for detailed stratigraphic work. These latter beds, which have long been known as the Inferior Oolite, are exposed in the area between Bridport on the English Channel and Hook Norton in north Oxfordshire. The Inferior Oolite, and in particular the middle and lower Inferior Oolite (Aalenian & Lower Bajocian), is preserved in two distinct facies, which are broadly separated by the present Mendip Hills, which appear to have acted as a 'positive area! throughout much of the Lower and Middle Jurassic. The upper Inferior Oolite, which is represented by colitic and bioclastic limestones, is relatively uniform in character across the whole outcrop. On the other hand, the lower and middle Inferior Oolite are largely represented by condensed

NORTH-WESTERN ANGLO-PARIS	PER BA:	IPHERY OF THE SIN		WESTERN PERIP South North S	HER 3 A	Y OF THE BASIN	INNER HEBRID
Dorset/Somers.		Cotswolds Upper		East Hidlands		Yorkshire	Great Estuarine
Inferior Oolite fossiliferous oolitic lsts. 31m.		Inferior Oolite pisolitic & bioclastic lsts. 14m	Estuari: 'series' (?mainly Bathon	Estuarine 'series' (?mainly Bathonian)	Ravenscar	Series'	
Middle Inferior Oolite highly fossiliferous 'iron-shot' lsts. 0.0-5.0m	MENDINS	Middle Inferior Oolite bioclastic Ists. 0.0-21m	RD SHALLOWS"	Lincolnshire Limestone Formation massive colit-	T WEIGHTON STRUCT	Group dominantly non-maring ssts. with thin marine intercalatio- ns, 200m	Bearreraig Sandstone Formation massive sst
Lower Inferior Oolite highly fossiliferous 'iron-shot' 1sts. 0.0-6.0m		Lower Inferior Oolite pisolitic & oolitic lsts. & marls 50m	"OXFU	Grantham Formation non-marine silts 0.0-5.0m Northampton Formation	MARKE	Dogger ferrug. sst.	

Table 1.

The main lithostratigraphic units of the British Aalenian-Bajocian, and their interelationship.

'iron-shot' and ferruginous limestones to the south of the Mendips, whilst to the north there is an expanded sequence dominated by bioclastic limestones in the middle, and by pisolitic limestones and marls in the lower Inferior Oolite, - see Table 1. These differences in lithology are mirrored by similar distinctions in the faunal characters of these beds, with marked differences in the brachiopod and bivalve populations, reflecting their different depositional environments. Perhaps the most marked faunal difference between these two areas, is to be seen in their ammonite distributions In the southern area the ammonites are abundant and well preserved, with intact fragile spines and lappets, whilst to the north they become increasingly rare and more badly preserved, with a generally worn and broken appearance, reflecting their allochthonous nature. It is thus hardly surprising that it is the southern area, which has in the past provided most of the figured and type ammonites.

(b) The south North Sea Basin

The advent of exploration for gas in the south North Sea, has produced a wealth of information relating to the subsurfac. Middle Jurassic. Unfortunately very few, if any of the boreholes drilled through this part of the sequence have been cored, hence our knowledge of these beds is mainly based on 'chippings' and various geophysical logging techniques.Based on this evidence, the Middle Jurassic ,West Sole Group (Rhys, 1974), appears to be very similar to the equivalent horizons at the western periphery of the basin, where they are exposed in the east Midlands and in Yorkshire. These beds consist of dominantly non-marine, clastic horizons, interbedded with marine limestones and ferruginous sandstones. In the east Midlands there is a rough balance between the non-marine silts and the marine elements : the Northampton Sand

Formation & the Lincolnshire Limestone Formation. However, towards the north there is an increase in the relative amount of clastic material, with a subsequent thickening of the non-marine elements. In north-east Yorkshire non-marine, deltaic sandstones dominate the Ravenscar Group (Hemingway & Knox, 1973), whilst the marine horizons (Eller Beck, Millepore and Scarborough beds), are relatively thin and of less significance.

(c) The Hebridean Basin

In the Inner Mebrides isolated patches of Jurassic sediments are preserved beneath the plateau basalts and cut by Tertiary dykes and sills. The Aalenian-Bajocian is represented by the Bearreraig Sandstone Formation (Morton, 1976), which is dominated by cross-bedded sandstones. Ammonites are not uncommon, and these beds appear to be wholly marine, although drifted wood and plant fragments are not uncommon at some horizons (Morton, 1965). There are rapid lateral facies changes in these beds, with massive, relatively unfossiliferous sandstones, such as the Druim an Fhurain sandstones of Strathaird (Skye), which pass laterally into thinner bedded highly fossiliferous units, such as the Rigg sandstones of Trotternish (N.Skye).

(d) Other, off-shore areas

The increasing momentum of the search for hydocarbons in the north-west european continental shelf is producing an immense quantity of information related to the distribution and nature of the off-shore Jurassic. Much of this data is still secret and commercial requirements will for some time militate against its general release. However, a broad overall picture is appearing, and it would seem that to the north of the Anglo-Paris Basin, the Aalenian Bajocian period is mainly

represented by clastic and sandstone facies. Thus in both the north North Sea (e.g. the Brent oil-field,Bowen,197⁴), and in Cardigan Bay (Penn & Evans,1976), there are thick,relatively unfossiliferous sandstone deposits. It is self-evident that sequences such as these are going to add little to our detailed biostratigraphic knowledge of this period. To the south, the English Channel has yielded sufficient information to suggest that the subsurface Inferior Oolite is very similar to the coastal exposures. Thus boreholes in Lyme Bay have revealed thicker, although otherwise very similar ammonite rich beds, to those of south Dorset. Detailed examination of this area could thus add considerably to the existing biostratigraphic picture.

2. BIOSTRATIGRAPHY

The Aalenian/Bajocian rocks, with their abundant and diverse ammonite faunas, have figured prominently in the development of both biostratigraphic methods and terminology (viz,d'Orbigny,1850,who was the author of the Bajocian Stage; Oppel,1856-8, who erected many of the Bajocian zones). The detailed stratigraphy of this period is closely linked with the study of the north Dorset successions. Many of the type specimens of the existing zonal and subzonal indices originate from this area, whilst the germinal work carried out by S.S.Buckman (1893), in the Sherborne district, forms the basis for much of the present subzonal scheme. However, whilst the existing biostratigraphic framework is apparently well founded and has had a long historical usage, it contains some inherent weaknesses. On one hand this has led to prolonged arguments and discussions concerning the interpretation and definition of the Bajocian Stage, whilst on the other hand

9

化学 化合成素化合

doubts have emerged concerning the validity of parts of the standard zonal scheme.

Many of these problems have arisen because of the peculiar nature of the mode of preservation of the Dorset Inferior Oolite, which has an extremely thin and attenuated sequence. The latter has strongly influenced the British view point on the status of the Aalenian Stage (see pp.501-4, below), and has helped to introduce errors into the interpretation of the faunal sequences, which consequently became incorporated in the standard zonal scheme. It is perhaps significant in this context to compare a section such as South Main-road quarry (Dundry Hill, Avon), where the discites, laeviuscula and sauzei Zone beds are less than 1.7m thick, with Lokut Hill (Bakony) Mountains, Hungary), where equivalent horizons in an 'ammonitico rosso' facies are over 4.0m thick (Galacz, 1976, p. 179). The Dundry beds are thus less than half the thickness of equivalent horizons, in what is commonly thought of as one of the most 'condensed' Jurassic facies. If the British Bajocian ammonite faunas, their stratigraphic distribution and rates of evolution are to be appreciated in comparison with those found in more expanded sequences, such as the Lias, then they must be compared in equivalent facies. The shales and mudstones of the Scarborough Formation (N.E.Yorkshire), are very similar to much of the Lias. Here at Hundale Point, Cloughton, the romani Subzone beds alone are 12m thick (compared to 0.30m, Oborne, Dorset), whilst the humphriesianum Zone as a whole is more than 20m thick (0.70-0.80m, Oborne).

The highly attenuated Dorset sequence also creates problems in the interpretation and recognition of biostratigraphic units. The ammonite faunas tend to occur in highly 'condensed', extremely fossiliferous 'fossil-beds' . Although very accurate collection can some times give the appearance of discrete

stratigraphic ranges within these beds (e.g. fig.3,p.39; figs. 5-6,pp.113-6), in most cases the faunas are homogeneous in any one particular layer or bed. Thus most of the faunas collected can only be assigned to'fauni-zones' in their broadest sense (Holland et al., 1978, p. 13), as an assemblage biozone. The interpretation of these zones or subzones in terms of accurate, relative stratigraphic ranges and first appearances (concurrent-range, partial-range biozones etc.), can only be done with reference to thicker, more expanded sequences. Although most of the British Lower Bajocian faunas in this light, are of limited international significance, their very abundance and diversity have given them a crucial importance in the interpret ation of the systematics of various ammonite groups. The real value thus comes, when the material obtained from these beds can be combined in a synthesis with the stratigraphic information from thicker sequences elsewhere.

(a) Development of the Standard Zonal Scheme

The backbone of the present zonal scheme for the Aalenian and Bajocian Stages is made up of the four Zones erected by Oppel (1856-8): that is the <u>parkinsoni</u>, <u>humphriesianum</u>, <u>sauzei</u> and <u>murchisonae</u> Zones. The majority of the other Zones have been added in a more piecemeal fashion (e.g. <u>subfurcatum</u> Zone, Terquem & Jourdy, 1869; <u>laeviuscula</u> Zone, Haug, 1894), and Buckman's work (1893, 1895, 1896, 1910), marks one of the first attempts, subsequent to Oppel, to produce a detailed and comprehensive Bajocian stratigraphy.

In his early work, Buckman (1881,1891)

used, with some modifications Oppel's Zones. However, in 1893, he introduced a new unit, the hemera, based on the acme of a particular ammonite species or genus. As originally defined (Buckman, 1893, p. 482), the hemera was a theoretically based, chronological unit, which was not intended as a replacement for the Zone. Most of Buckman's contemporaries did not understand the intrinsic difference between chronological and stratigraphic units, and tended to treat hemeræ as nothing more than stratigraphic subdivisions of a slightly lower order than the Zone. Indeed Buckman in his later works also seems to have lost sight of the distinction between the two, since Zones and hemeræ are interchanged in his later zonal tables (Buckman, 1913 in 1909-30;1915).

Buckman readily appreciated the highly condensed nature of the Inferior Oolite, which led him to greater refinements of his polyhemeral scheme. He argued (Buckman, 1893, p. 482), that if the acme of any one species was characteristic of a particular unit of time (a hemera), then in condensed sequences, it would be logical to find several different hemeral indices together in one bed, whilst in thicker sequence they would have distinct stratigraphic ranges. Taking this methodology to extreme lengths, in condensed horizons, virtua-11y every ammonite could be assigned to a different hemera . Whilst there is some logic to the above argument, in his later works, such as Type Ammonites (Buckman, 1909-30), Buckman went to absurd lengths in the creation of new hemerae, many of which were based on poorly localised material, or apparently in some cases, no ammonites at all . This led to the production of numerous totally artificial hemerae, which either were define as containing several different ammonite species, which can now be shown to occupy different horizons, or which were

duplicated, with several hemeraebeing applied to the same bed. In the case of some Lower Bajocian hemerae, it is evident that there is an inextricable mixture of these two faults (see Parsons, 1974, Table 1).

With the recognition of the dubious nature of Buckman's later polyhemeral schemes, there came a reaction against all his work. This is unfortunate, since much of his detailed research carried out prior to 1910, is of outstanding value. Later workers (Spath, 1936) were forced to go back to earlier zonal concepts, and "Oppel's Zones were re-instated, often with Buckman's earlier hemeræ as subzones (e.g.the 'Sowerbyi' Zone). Arkell (1956) merely used a modification of Spath's scheme, it hence contained the inherent weaknesses carried over with Buckman's hemerae.

There has recently been a resurgence of interest in the problems of Bajocian stratigraphy, with some important studies in thick ,uncondensed sequences (e.g.Pavia & Sturani, 1968). This would suggest that we are gradually approaching the situation, where a consensus can be reached on a Bajocian stratigraphic scheme, which will be a fairer representation of the ammonite faunal successions.

3. THE ORGANISATION AND SCOPE OF THE THESIS

The arrangement of material within this thesis does not follow the generally accepted practice of subdivision into closely related chapters. Instead an attempt has been made to present the material in the form of self-contained papers, which are either in a readily publishable form, or indeed have already been accepted for publication. In consequence there is a certain amount of duplication between sections, mainly in the reference lists, as well as some disparity in style, as individual papers have been tailored to the 'house



Figure 2

A sketch-map showing the location of the main areas of the Inferior Oolite, and other, Bajocian rocks, which are described in the following sections: 1, south Dorset (Eridport /Beaminster); 2, north Dorset (Sherborne-Yeovil); 3, Doulting district; 4, Dundry Hill, near Bristol; 5, the Cotswold and Mendip Hills; 6, the north Cotswolds; 7, the north Yorkshire Moors; 8, south Lincolnshire (the Grantham district). style' of their intended destinations. Another result of this method of arrangement has been to produce a degree of discontinuity between individual sections. The aim here is thus to draw together the individual strands, and to clarify the underlying objectives of this work.

(a) The stratigraphic range of the study

In the course of this work, the zonal scheme for the Lower Bajocian (Middle Bajocian, Arkell, 1956), was found to be the most unsatisfactory. Hence most attention has been paid here to a revision of the horizons between the concavum and garantiana Zones. The first two sections (1A,1B) are thus devoted to a discussion and revision of the 'sowerbyi' (now discites & laeviuscula Zones) - sauzei Zones, and the humphriesianum -subfurcatum Zones, whilst the third section (1C) gives an account of the garantiana and parkinsoni Zones. These, and subsequent units set out to apply this revised stratigraphic scheme to the British Bajocian. This work has not been totally restricted to this horizon, since the Aalenian beds of Dundry Hill (section 2A), have been the subject of some attention. All of the above work, together with a few additional points, are summarised in the final section (4). (b) The geographical range of the study

The geographical limits of the study have been wide, and included much of Europe, but in the revision of the zonal scheme, most attention has been paid in sections 1A-C, to the Sherborne area of Dorset (Figure 2, area 2). In the stratigraphi revision of parts of the British Bajocian rocks, there has been a concentration on specific areas; on Dundry Hill (section 2A- area 4, Fig. 2) and the Cotswold and Mendip Hills (section 2B - area 5, Fig. 2). Accounts of the stratigraphy of the Doultin district, Somerset (area 3), the north Cotswolds (area 6) and south Dorset (area 1) have been given elsewhere (Parsons, 1975a, 1976 & 1975b respectively). There is a sparcity of Bajocian ammonites in Britain north of the Anglo-Faris basin, but there are several horizons, which have yielded isolated, but important faunas. Thus in section 2C the stratigraphy of the Yorkshire Scarborough Formation is revised (area 7), whilst in a paper from the Proceedings of the Yorkshire geological Society (Parsons, 1974b), there is a brief discussion of ammonites from the Lincolnshire Limestone Formation (area 8); a subject which in collaboration with Dr.M.Ashton, is being treated in greater detail elsewhere. I have refrained from any detailed discussion of the Hebridean basin, apart from a brief reference in section 4, since my field-work has added little to the recently published studies of this area by Dr.N.Morton (1965, 1971, 1976).

4. ACKNOWLEDGEMENTS

Individual acknowledgements follow each of the separate sections. Here I will restrict myself to thanking Hugh Torrens for his help during his supervision of this research project, which was initially supported by a University of Keele Research Studentship : this award is gratefully acknowledged. Finally I must express a great debt of thanks to my wife Marion,who for longer than she would care to remember, has had to put up with living twenty-four hours of the day with ammonites !

5.REFERENCES

Most of the references cited in the Introduction will be found in the reference lists following the individual sections, particularly 1A-B,2B & 4.

1. TOWARDS A REVISION OF THE STANDARD ZONAL SCHEME FOR THE BAJOCIAN AND PARTS OF THE AALENIAN STAGES

There are inherent problems in the interpretation of the Bajocian stratigraphy of southern England, which are the result of vagaries in preservation and exposure of the rocks concerned. These difficulties fall into two groups. First the so called 'Sowerbyi' and Sauzei Zones are preserved in a series of disjunct and discontinuous, but highly fossiliferous 'fossil-beds' . The faunal sequence for this part of the column has thus to be pieced together from separate exposures. The problems with this part of the stratigraphic scheme, have thus been the result of previous mis-correlation of these various exposures. The second difficulty is the result of limited preservation, linked with lack of exposure. Thus the Humphriesianum/Subfurcatum Zones are only preserved to any great thickness in a small area to the north-east of Sherborne Considering these difficulties, Buckman (1893), did a superb job in unravelling the stratigraphy of this period.Unfortunately most of the ammonites, which he subsequently figured (Buckman, 1909-30), were based on poorly localised material. A serious past omission has been the lack of work on the Garantiana Zone rocks. These beds, which are thick and relatively fossiliferous have been largely ignored, subsequent to Buckman's introduction of the garantiana hemera. The aims of this part of the work are thus to redescribe the faunal successions to locate as many topotypes as is possible of Buckman's species, and lastly to use this more reliable stratigraphic information to re-interpret the zonal scheme.

IA.THE SAUZEI AND 'SO CALLED' SOWERBYI ZONES OF THE LOWER BAJOCIAN (MIDDLE JURASSIC) OF EUROPE

SUMMARY

The Sauzei and Sowerbyi Zones of the Lower Bajocian (Middle Jurassic) are discussed in the light of new evidence from Normandy (France), Schwabia (South Germany) and Southern England. The Sauzei Zone is re-interpreted as a 'fauni-zone'. The Sowerbyi Zone is rejected and evidence is given for its replacement by the Laeviuscula Zone (HAUG, 1894). It is shown that the subzonal divisions of the Sowerbyi Zone were based on incorrect stratigraphic evidence. The Laeviuscula Zone is subdivided into a redefined Laeviuscula subzone (senior synonym of the Trigonalis subzone) and an Ovalis subzone; the Discites subzone being elevated to full Zonal rank. A detailed analysis is given of the ammonite faunas of Southern England from the Sauzei to the Concavum Zones, as well as a brief summary of the correlation of equivalent horizons in Normandy and Schwabia.

Modified from:

"The <u>Sauzei</u> and 'so called' <u>Sowerbyi</u> Zones of the Lower Bajocian". Newsl. Stratigr. <u>3</u> (3.) pp 153-179.

CONTENTS

1.	INTRODUCTION AND	ACKNOWLEDGEMENTS
2.	THE SAUZEI ZONE:	a/ Schwabia 24
		b/ Normandy 25
		c/ Southern England 26
		d/ Typical fauna of the Sauzei Zone 27
3.	THE 'SO CALLED' S	OWERBYI ZONE 29
4.	THE LAEVIUSCULA Z	ONE
5.	THE SUBZONES OF T	HE SOWERBYI ZONE 34
6.	THE SUBZONES OF T	HE LAEVIUSCULA AND CONCAVUM ZONES
7.	THE SUCCESSIVE AM	MONITE FAUNAS OF THE SAUZEI, LAEVIUSCULA,
	DISCITES AND CONC	AVUM ZONES IN SOUTHERN ENGLAND: 38
		a/ The Sauzei Zone 39
		b/ The Laeviuscula Zone and Subzone 41
		c/ The Ovalis Subzone of the Laeviuscula
		Zone
		d/ The Discites and Concavum Zones 45
8.	THE CORRELATION C	F SCHWABIA AND NORMANDY:
		a/ Schwabia 49
		b/ The Wutach 51
		c/ Normandy 52
9.	CONCLUSION:	a/ The Zone of Graphoceras concavum 53
		b/ The Zone of Hyperlioceras discites 53
		c/ The Zone of Witchellia laeviuscula 54
		d/ The Zone of Otoites sauzei
10.	REFERENCES	56

1. INTRODUCTION

Although one of the first stages in the Jurassic to succumb to modern, detailed, stratigraphic methods, (S. BUCKMAN, 1893), the Bajocian has been largely ignored by later workers. The early work done by Buckman in the Sherborne area (Dorset, England) was readily accepted by Spath (1936) and Arkell (1954, 1956) in an only slightly modified form: 'Thanks to the work of the late S.S. Buckman and Mr. L. Richardson, the Inferior Colite Series from Dorset to Northamptonshire is so thoroughly known that almost perfect generalisation is possible: a detailed section of the beds can be drawn right across the country'. (ARKELL, 1933 p.vi). This has created the erroneous impression that the fundamental stratigraphic work, particularly in the Lower Bajocian (= middle Bajocian olim) has long since been completed. However, current research has suggested that the existing Zonal and Subzonal scheme (see Table 1), far from being acceptable, is often based on erroneous or inadequate information. In this the Sauzei and 'so-called' Sowerbyi Zones have proved to be the most serious offenders. It has been found necessary to re-examine in some detail the stratigraphic basis for the very existence of these two Zones and attendant subzones.

The Sauzei and Sowerbyi Zones, introduced by Oppel, (OPPEL, 1856 and 1862), were replaced by Buckman with a series of smaller units, hemerae, based on the acme of a particular species or genus, (BUCKMAN, 1893, 1895, 1910). Subsequently Spath reinstated Oppel's Zones on grounds of priority, relegating Buckman's earlier hemerae to subzonal status, (SPATH 1936). It was this latter scheme which was accepted by Arkell as the standard for the World Bajocian.

oppel 1856-62	S. DUCKNAN 1891	S.BUCKMAN 1893-6 'Hemerae'	BUCKMAN, 1930, Inferred position of Hemerae, from range of ammonites assigned to them.	SPATH , 1936 Arkell, 1956	SCHEME SUGGESTED HERE
SAUZEI (1856)	SAUZEI	SAUZEI	alsaltica propinguans pars sauzei " Labyrinthoceras " mollis	SAUZEI	SAUZEI
SOWERBYI	SOWERBYI	WITCHELLIAE SONNINIAE (1896) <u>non</u>	ruber brocchij Shirbuirnia Witchellia pars sauzei " Labyrinthoceras " mollis fissilobatum hebos ovalis pars mollis ? Bradfordia	Laeviuscula Trigonalis	Laeviuscula Vinoscula Ovalis
(1002)		DISCITAE	Docidoceras Trilobiticeras Rudidiscites Depaceras Reynosella Platygraphoceras ? eudmetum	Discites	DISCITES
NURCHI SONAE	CONCAVUM	CONCAVI	stigmosum crassispinata casta cornu cornu concava	CONCAVUM	Concavum

Table 1.

Development of the standard zonal scheme for parts of end the British Aalenian/Bajocian.

(ARKELL, 1954, 1956, etc), and which up to recently (TORRENS, 1969) no one has seriously questioned. None the less serious confusion has, and still does exist over the exact boundaries and faunas of these two Zones. This has at least in part arisen from the publication of Buckman's 'Type Ammonites' (BUCKMAN, 1909-1930). In this work Buckman created large numbers of new genera and species of ammonites, which were figured, but only accompanied with the most abbreviated of descriptions. Many of these figured ammonites were unlocalised specimens, which Buckman had acquired from various sources, and then assigned to their respective hemerae purely by matrix. Whilst in his early years Buckman had an extensive first hand knowledge of the Inferior Oolite and an intuitive judgement which probably still could not be bettered, towards the end, his work came to be based more on doubtful 'Biological Laws' and slight differences in matrix, than on in situ collection. This regrettable fact led to many key species of ammonites, common and highly characteristic of certain horizons, being recorded by Buckman, as coming from a whole series of, largely fictional, hemerae. As an example one may cite the fauna of the 'green marl bed' of Frogden quarry, Sherborne, Dorset (BUCKMAN, 1893, Sect. xv, bed 9), where the Stephanoceratid ammonites, (such as Frogdenites spiniger) were recorded by Buckman as coming from the Sauzei hemera, whilst the Witchellia spp. were recorded from the Witchellia hemera, in spite of their occurrence together in the same bed. This confusion, once created, became firmly entrenched in the subsequent literature, until Arkell was able to quote faunal lists from the Lower Bajocian of Dorset, which bore no resemblance to those produced by in situ collection (ARKELL, 1956, p.33).



Figure 1.

A map showing the main areas of Europe under discussion : 1. Southern England ; 2. Normandy, north-west France ; 3. Schwabian Albe, south-west Germany. The time has thus come for a critical re-appraisal of Lower Bajocian stratigraphy, based on accurately localised collections. The following account is an attempt at a long overdue revision of the stratigraphy of beds of this age in certain key areas of North-West Europe. The main areas under discussion are those which figured largely in the work of the earlier stratigraphers, Oppel, Haug, Buckman etc., that is the Schwabian Albe, Normandy and Southern England, (see fig. 1).

ACKNOWLEDGEMENTS

I should like to thank the curators of the following museums for the loan and study of material in their care: The British Museum (N.H.) and the Geological Survey Museum in London, Oxford University Museum, the Sedgwick Museum, Cambridge, Manchester Museum, Bristol City Museum, Bristol University Museum, Reading University Museum, Geologisch-Paläontologisches Institut und Museum Tübingen, and the Staatliches Museum für Naturkunde Stuttgart; as well as many other institutions, and numerous private collectors, in particular, J. Whicher (Bristol) and U. Bayer (Stuttgart).

For much help and discussion in the field, I would like to thank J. Callomon, U. Bayer, M. Gatrall, J. Senior and in particular H. Torrens.

In the following work, numbers prefixed by these abbreviations refer to specimens in the following Institutions and private collections.

BMNH.	-	British	Mus	seum (N.H.), London.		
Ce	-	Geologisch-Paläontologisches Museum, Tubingen.				
ST		Staatliches Museum, Stuttgart.				
CP	-	The aut	hor	's collection.		
Bayer Co	ll e	ction	-	Private collection of U. Bayer, Deizisau,		
				near Stuttgart, W. Germany.		
Morton C	011	ection	-	Private collection, N. Morton, Birkbeck		
				College, London.		
Whicher	Col	lection	-	Private collection, J. Whicher, Bristol,		
				England.		

2. THE SAUZEI ZONE

Oppel first introduced the term 'beds with Ammonites sauzei', for the lower part of his Humphriesianum Zone (OPPEL, 1856-58, p.305) but later separated them first as a Subzone (op. cit., p.334) and later as a full Zone (op. cit., p. 369). For the defining fauna of the Sauzei Zone Oppel gave Ammonites sauzei, A. brocchii, A. brongniarti (op. cit., p.305) and subsequently added A. bayleanus, A. jugosus, A. sowerbyi and A. tessonianus (op. cit., p.334). It is almost impossible to interpret these specific names in terms of the present more restricted concepts of these species. Hence in order to define the fauna of the Sauzei Zone in modern terms it is necessary to examine the actual strata cited by Oppel as belonging to his Zone and to re-identify any ammonites produced by these beds. Amongst other regions noted as having representatives of the Sauzei Zone (Horizon 21 in Table 64, OPPEL, 1856-58) Oppel specifically included three areas; Schwabia, South-western Germany, Normandy, North-west France and Southern England, (see Text Fig. 1).

a/ Schwabia

The bed cited by Oppel in this area was described as "Blaulich, graue, harte, sandige kalke, mit <u>Ammonites sauzei</u> of Neuffen and Hohenzollern" (OPPEL, 1856-58, Table 64). There can be no doubt that Oppel was here referring to Quenstedt's Blaukalke, since in this connection he refers to the Blaukalke (QUENSTEDT, 1843) as "Die blauen kalke sind das lager des <u>A. sauzei</u>" (op. <u>cit.</u>, p.336). It is to be stressed that the Blaukalke must be considered the type horizon of the Sauzei Zone and Schwabia the type area (ARKELL in DONOVAN & HEMINGWAY, 1963, p. 307).

The Blaukalke is only sparsely fossiliferous and it is now poorly exposed. The few fossils present tend to occur in clusters on the upper surface of the bed which in the Beuren region has recently yielded several ammonites (Emileia (E.) cf. polyschides (Waagen) & E. (Otoites) sp. - U. Bayer collection). The matrix of this bed is sufficiently characteristic to enable use to be made of geographically localised material. The following have been recognised in museum collections (see also BUCK, HAHN & SCHADEL, 1966):-

Emileia (E.) polyschides, Bayer Collection & ST10028.

E. (E.) pseudocontrahens Maubeuge, ST22551.

E. (<u>Otoites</u>) <u>contractus</u> (S. Buckman <u>non</u> Sow.) - <u>sauzei</u> (d'Orb.) group, Bayer col.

Kumatostephanus aff. kumaterus S. Buck., ST22549.

K. triplicatum (Renz), Ce5-66-13.

K. turgidulum (Qu. emend. Welsch), Ce5-66-3.

b/ Normandy

The bed which Oppel correlated with the Sauzei Zone in Normandy was the basal member of the 'Oolithe Ferrugineuse de Bayeux', which rests on top of the Malière. This horizon agrees well with the 'Couche verte' of subsequent French authors (BIGOT, 1900; ERASIL, 1895). Although most of the famous exposures of this latter bed have disappeared, the extensive coastal exposures at St. Honorinedes-Pertes, Normandy, are still available for study (RIOULT, 1962; RIOULT & GABILLY, 1967). The matrix of this bed; rich in glauconite, which often replaces the shells of the fossils (DANGEARD, 1940); is unmistakable even in unlocalised museum material. The following have been collected in situ from the coastal exposures :-

Emileia (E.) polyschides, CP 1591.

E. (E.) pseudocontrahens, CP 1590.

E. (E.) cf. polymera (Waagen), CP 1593.

Kumatostephanus turgidulum (= K. bigoti, Haug ex. Munier-Chalmas

m.s. per. QUENSTEDT, 1886, pl. 65, fig. 9), CP 1585.

K. cf. perjecundus S. Buckman, CP 1586.

Mollistephanus sp. nov., CP 1597.

Sonninia felix (S. Buckman), CP 1607.

S. cf. propinquans (Bayle), CP 1605.

Amongst numerous other specimens which I have seen in various museum collections, the following, cited by d'Orbigny (1842-51) from the Tesson collection are now in the British Museum:-

<u>Skirroceras bayleanum</u> (Oppel), BMNH, 37069. <u>Emileia (E.) polymera</u>, BMNH, 37320.

Labyrinthoceras meniscum (Waagen), BMNH, 37268.

All three specimens come from the old quarry at St. Vigor and are of considerable value for the correct interpretation of these three species.

c/ Southern England

The localities cited by Oppel in Southern England as having Sauzei Zone rocks were Burton Bradstock (Dorset), Dundry Hill near Bristol and the Cheltenham area of the Cotswolds (OPPEL, 1856-8). The beds of Sauzei Zone age near Cheltenham are the Bourguetia/ Phillipsiana beds (BUCKMAN, 1897), at Burton Bradstock the Red Bed (BUCKMAN, 1910) and at Dundry the 'Brown Iron-shot bed' (BUCKMAN & WILSON, 1896). Ammonites are relatively rare at the first two localities and it is only at Dundry that abundant Sauzei Zone faunas may be collected. The following have been collected <u>in situ</u> from South Main-road Quarry, Dundry (BUCKMAN & WILSON, 1896, p.691, section ix, bed 4):-

Emileia (E.) polyschides, CP 2414.

<u>E.</u> (<u>E.</u>) polymera, CP 2423.

E. (Otoites) cf. sauzei (d'Orb.), CP 2448.

E. (O.) cf. fortis Westermann, CP 2425.

Kumatostephanus kumaterus, CP 2413.

Labyrinthoceras sp. CP 2402.

Skirroceras bayleanum, CP 2400.

Sonninia aff. felix, CP 2422.

<u>Witchellia</u> (W.) <u>hebridica</u> (Morton), CP 2442.

W. (Pelekodites) cf. sulcata (S. Buckman), CP 2451.

A further discussion of other Sauzei Zone ammonite faunas from Southern England is given in Section 7.

d/ The typical Faunal and zonal Index of the Sauzei Zone.

It is impossible to be certain of the absolute range of \underline{E} . (<u>O</u>.) <u>sauzei</u> as no collections have yet been made through continuous and 'expanded' beds of this age, hence the Sauzei Zone must be considered as an assemblage zone, based on the occurrence of the following characteristic fauna:-

Emileia (E.) <u>bulligera</u> S. Buckman - <u>greppini</u> Maubeuge group. <u>E. (E.) polymera</u>

 \underline{E} . (\underline{E} .) <u>pseudocontrahens</u>

Kumatostephanus kumaterus
K. perjucundus S. Buckman - triplicatum (Renz) group.

K. turgidulum

Labyrinthoceras meniscum

Skirroceras bayleanum

Sonninia propinquans - felix group.

The index species of the Sauzei Zone, Emileia (Otoites) sauzei (d'Orb.) is at this moment under rather a cloud. Whilst specimens do exist which are very close to d'Orbigny's original figure (such as BMNH, 37323, St. Vigor, near Caen, Normandy, ex. Tesson collection), the subsequently designated neotype (WESTERMANN, 1954) has completely changed the interpretation of this species. However, this neotype designation must be considered totally invalid under article 75c, 3 & 5, of the Code of Zoological Nomenclature, since syntype material still exists. The specimen figured by d'Orbigny was only one of at least eight syntypes mentioned by him, of which the specimens figured by Westermann from the d'Orbigny collection form two (WESTERMANN, 1954, pl.1 fig. 1 - neotype; pl. 1, fig. 2). No type specimen was designated by d'Orbigny and no subsequent author has designated any of the syntypes lectotype. It was thus impossible for Westermann to erect a neotype as a replacement for a type specimen which in fact has never, as yet, existed. The d'Orbigny collection has not yet been re-organised, it would thus be impossible to be certain of the absence of the original figured specimen of \underline{E} . (0.) sauzei. Thus no action should be taken until it is certain that the latter is missing, then one of the original syntypes (which need not be from the d'Orbigny collection) should be chosen as lectotype. There is no need for the suppression of the existing neotype via the International Commission on Zoological Nomenclature, since it was both invalid from

its inception and published prior to 1961 (Article 75c of the 'code').

Until the d'Orbigny collection has undergone its present reorganisation, <u>E</u>. (<u>O</u>.) <u>sauzei</u> must remain in a state of flux. This need not however effect its status as Zonal index, since its position will finally be stabilised by the selection of a lectotype; its existing use is merely as a 'label' for an assemblage zone.

3. THE'SO CALLED' SOWERBYI ZONE

Unlike the Sauzei Zone and most of the other Zones erected by Oppel (1856-8) in 'Die Juraformation' the Sowerbyi Zone was a poorly documented after thought by Oppel (1862-3) in 'Uber Jurassiche Cephalopoden'. Whilst the Sowerbyi Zone's relative position was clearly indicated 'rather lower than Ammonites sauzei and higher than A. murchisonae base above the ferruginous sandstone of Gingen, Fils and neighbouring exposures' ... (Oppel, 1862-3, p.128), (the sandstone referred to is the Concavum or Donzdorf sandstone of subsequent German authors), no reference was made to the characteristic ammonite fauna of the new Zone. It was left to his pupil Waagen who monographed the Sowerbyi Zone over much of extra-alpine Europe, to produce the still generally accepted interpretation of this Zone (Waagen, 1867). The beds which have been correlated with the Sowerbyi Zone in its type area (Gingen, Wurtemburg, South Germany) are the clays and 'Wedel' sandstones below the Blaukalke down to and including the 'Sowerbyi Bank' or 'Oolite', all of which have been included in the 'Sowerbyi Schicten', (see fig. 3). Thus, whilst the litho-stratigraphic limits of the Zone in the type area are clearly delimited and, thanks to the work of Quenstedt (1886), Stahlecker (1926), Dorn (1935) and Oechsle (1958),

its fauna is adequately described, the use of <u>Sonninia sowerbyi</u>, as its index is totally unsuitable. The impediment to its continued use is the incorrect identification of <u>S. sowerbyi</u> by Oppel, Waagen and most subsequent workers.

The type specimen of <u>S</u>. <u>sowerbyi</u> is an exceedingly poorly preserved ammonite, of which Sowerby's figure leaves much to be desired. It has however been carefully re-figured (BUCKMAN, 1904, pl.52) and after examining the holotype in the Bristol City Museum, I have come to very similar conclusions to those of Buckman (<u>op</u>. <u>cit</u>.), which are:-

i. The type specimen came from the top most, densely 'ironshot' horizon at Dundry Hill, which is entirely Sauzei Zone in age (BUCKMAN & WILSON, 1896).

ii. It is the totally septate inner whorls of what would be, if complete, a large species. It is impossible to compare it with any but similarly incomplete specimens. Unfortunately a spinose nucleus is a feature common to the Sonninids.

iii. Taking into account its type horizon, whorl cross-section and spinose nature, the holotype of <u>S. sowerbyi</u> is most easily reconciled with a nucleus of a specimen of the <u>Papilliceras mesacantha</u> (Waagen) group.

<u>Sonninia</u> <u>sowerbyi</u>, whose type specimen is quite indeterminate and which came from the Sauzei Zone, is thus totally unsuitable as an index for the bio-stratigraphic unit previously called the 'Sowerbyi Zone'. This is a conclusion obviously reached by Buckman (1887-1907, p.63-4), who rejected its use and replaced it with a series of hemerae (BUCKMAN, 1893, et seq.). Arkell (1954 & 1956) was of the opinion that Buckman had never found the true horizon of <u>S. sowerbyi</u> and taking into account reports of the occurrence of the latter in the Sauzei Zone, suggested the possible inclusion of the senior Sauzei Zone in the junior Sowerbyi, as a subzone (ARKELL, 1956), an indefensible recommendation subsequently followed by the British Geological Survey (WILSON, WELCH, ROBBIE & GREEN, 1959).

The type horizon of <u>S</u>. <u>sowerbyi</u> was $\int as$ the Sauzei Zone by Buckman, and has been confirmed as such here. The reports of this species in the so called Sowerbyi Zone <u>sensu stricto</u> are due to the presence of other spinose Sonninids, such as <u>Euhoploceras</u>, to which the type of <u>S</u>. <u>sowerbyi</u> has a great superficial similarity. This inconsistency has two possible solutions:-

i. keep to the boundaries of the present Sowerbyi Zone and merely choose a new index species.

ii. revert to one or more of the zones proposed as replacements for the Sowerbyi Zone .

On grounds of priority the second alternative must be accepted. The first, partial replacement was the Concavum Zone, which as originally described (HUDLESTON, 1887), included the present Discites Subzone. The remainder of the Sowerbyi Zone can be replaced by the Laeviuscula Zone, which had considerable popularity in France at the turn of the Century. It was rejected by Arkell (1954, p.73) as a synonym of the Sowerbyi Zone, on the grounds that Haug (1910) had cited <u>S. sowerbyi</u> as part of its characteristic fauna. However it is unlikely that Haug's identification of <u>S. sowerbyi</u> would be any more reliable than Waagen's or Oppel's, there thus seems no barrier to the use of <u>Witchellia</u> (<u>Witchellia</u>) <u>laeviuscula</u> as an index for a replacement Zone for the Sowerbyi Zone, which is here recommended.

A problem does exist with the positioning of the Discites Subzone, the basal unit of the Sowerbyi Zone . Whilst it would be logical to combine the Discites faunas within an extended Concavum Zone, as originally described (HUDLESTON, 1887; EUCKMAN, 1887-1907), this creates considerable problems concerning the finally accepted Aalenian/Bajocian boundary. The best, indeed the only solution to this problem is to elevate the Discites Subzone to full Zonal rank, a procedure which is followed here. The lectotype of the index species of the Discites Zone appears to be lost (J. Callomon pers. com.). However since Waagen (1867, p. 600) mentions twenty-six syntypes of <u>Hyperlioceras</u> ('<u>Ammonites'</u>) <u>discites</u>, of which at least one survives (the holotype of the invalid species <u>H. desori</u> (Moesch <u>non</u> Pictet), refigured by BAYER, 1969); there should be no problem in designating a new type specimen.

4. THE LAEVIUSCULA ZONE

Based on the earlier use of a '<u>Witchellia</u> horizon or beds', between the Concavum and Sauzei Zones (HAUG, 1893), Haug introduced the Zone of <u>Witchellia laeviuscula</u> (type specimen, recently refigured, WESTERMANN, 1969, Text fig. 35), in 'La Grande Encyclopédie' (HAUG, 1894). Earlier Haug had discussed the possibility of the existence of such an horizon (HAUG, 1893) and had referred the so called Sowerbyi Zone of Toulon (DOUVILLÉ, 1885), the '<u>Witchellia</u> beds' of Normandy (MUNIER-CHALMAS, 1892) and Dundry

(S. BUCKMAN, 1887-1907) as being its possible equivalents. The exact biostratigraphic limits of the Laeviuscula Zone can be defined precisely. Haug used the Concavum Zone in its original extended sense, including the fauna, now separated within the Discites Zone (HAUG, 1891). Although originally including the Sowerbyi Zone of Toulon in his interpretation of the Sauzei Zone, Haug was well aware of the true fauna of the latter Zone, as found on the Normandy coast in the 'Couche Verte' and in the Blaukalk of Wurtemburg (HAUG, 1891, p.67). The limits of the Laeviuscula Zone can thus be defined at the base, as coming above the Discites fauna, and at the top, as coming below the Sauzei Zone, as redefined here. Haug's next use of the Laeviuscula Zone was in connection with a description of the Bajocian of the Basses Alpes(HAUG, 1900), and this area would probably be the most suitable for the designation of Type area and Horizon for this Zone.

Recent work in the Basses Alpes has produced a slightly clearer picture of the stratigraphic succession (PAVIA & STURANI, 1963). It is noticeable that these authors had some difficulty in distinguishing any more than one fauna between their Sauzei Zone and Discites subzone, which they included in the Laeviuscula subzone of the Sowerbyi Zone . Thus whilst the Laeviuscula Zone can be recognised in the Basses Alpes, its precise basal limits must be determined by those of its basal Ovalis subzone. However, it has not yet been possible to recognise the latter subzone in the Basses Alpes, an omission which will no doubt be rectified by future collecting.

5. THE SUBZONES OF THE SOWERBYI ZONE

The subzonal divisions of the Sowerbyi Zone, in present use, are based on the Buckman scheme of hemerae, as modified by Spath, (SPATH, 1936).

After having given reasons for rejecting 'Ammonites', (Sonninia) sowerbyi, as Zonal index. (BUCKMAN, 1887-1907, p.63-4), Buckman introduced a series of hemerae as replacements, (BUCKMAN, 1893). At first he was content with two, the Witchellia hemera, based mainly on the lower half of the 'Sandford Lane Fossil Bed; (BUCKMAN, 1893, Sect. IX), and the Discites. Buckman later confined the Witchellia hemera to the middle and introduced the Sonniniae hemera for the basal third of the 'Fossil bed', (BUCKMAN 1895). The table which Buckman produced for the correlation of the Cotswold Inferior Oolite, (BUCKMAN, 1895, Tab. V.) perhaps gives the most accurate picture of Buckman's mature views on the correlation of these beds, and it is reproduced here in a slightly modified form, (Tab. 2). After the restriction of the genus Sonninia to those ammonites of the S. propinguans Bayle group, the Sonniniae hemera took the new generic name Shirbuirnia, (BUCKMAN, 1910) - see Tab. 1 for fuller details of the evolution of these Subzones.

These Buckman hemerae were merely modified by Spath, who gave them valid specific, rather than generic indices, when he incorporated the Witchellia and Shirbuirnia hemerae in the reinstated Sowerbyi Zone as the Laeviuscula and Trigonalis subzones. Although these two latter subzones were recognised by Buckman, (as hemerae), at several localities in Southern England, (BUCKMAN, 1887-1907, p.ccvii), the faunas he figured from them were almost entirely from the Sandford Lane 'Fossil Bed'. This single bed, thin and highly condensed, (less than 0.6m. in thickness) was virtually the sole source of ammonites of, in particular, the Trigonalis subzone. It is surprising that no one has seriously questioned Buckman's division of this thin bed initially into 2-3 horizons, and later into more than 9 hemerae (BUCKMAN 1909-30). Of the three hemerae described by Buckman up to 1910, the Shirbuirnia hemera or Trigonalis subzone was the most restricted in its distribution; Buckman himself was the first to admit that this fauna from Sandford Lane was virtually unique. He could not point out any other horizon in England with an exactly comparable fauna to the latter, but none the less, thought it the direct equivalent to that described by Waagen from Gingen (WAAGEN, 1867). For the last sixty years, the stratigraphic divisions, on which the subzones of the present Sowerbyi Zone rest, have been based on the faunas collected from one bed, in one isolated quarry. This quarry was only commercially worked for a short period of time in the 1870's, and was reopened under the direction of Buckman and Hudleston in 1892, in order to collect its well preserved and abundant fauna. However, most of the ammonite species later figured by Buckman from this locality were unlocalised specimens from the Darrel Stephens collection. No one in the last eighty years has seen the need for a re-examination of this key section, which is still available for study. It was thus with considerable interest, that, with the consent and assistance of Sherborne Castle Estates and their tenant farmer, I re-excavated the 'Fossil Bed' (in 1970-1), in order to test the validity of Buckman's subdivisions. The results of this work are to be found in the sections on the Subzones of the Laeviuscula Zone and on the successive ammonite faunas of S. England.

In brief, it was found that it was impossible to separate a <u>Witchellia</u> and Shirbuirnia fauna from the basal half of the 'fossil bed's the two were totally mixed together. Underlying the 'fossil bed' were found sandy limestones, containing Sonninia (Fissilobisceras) ovalis, following the faunal sequence originally described by Buckman from the Cotswolds (BUCKMAN, 1895) and Dundry, (BUCKMAN and WILSON, 1896). There has been much confusion over the stratigraphic relationship between the genus Shirbuirnia and the Sonninia ovalis group. Thus for example the alleged occurrence of Shirbuirnia directly above the Discites fauna in Skye (Inner Hebrides, Scotland, MORTON, 1965), has been accepted as proving the present subzonal scheme, (WESTERMANN and RICCARDI, 1972). The fact that this particular occurrence of Shirbuirnia is in fact based on mis_identified Sonninia of the S. ovalis group, (J. Callomon, pers. comm.), and that the true Shirbuirnia, as in Dorset, occur at a higher horizon, makes a nonsense of any stratigraphic or systematic conclusions, based on such erroneous records, (WESTERMANN and RICCARDI, 1972).

6. THE SUBZONES OF THE LAEVIUSCULA AND CONCAVUM ZONES

The re-examination of the Sandford Lane 'fossil bed' has proved that it is impossible to separate the faunas of Buckman's Witchellia and Shirbuirnia hemerae, as based on this bed. Since <u>Witchellia</u> <u>laeviuscula</u> has precedence over <u>Shirbuirnia trigonalis</u> as index for this subzonal fauna, it is used as such here. Whilst it would be possible to use <u>S. trigonalis</u> in order to prevent the double use of <u>W. laeviuscula</u> as both Zonal and subzonal index, too much confusion

		,				
HEMERAE	COTSWOLDS	SANDFORD LANE				
SAUZEI	BOURGUETIA and	Limestone with				
	PHILLIPSIANA DEDS	Otories sauzer and				
		Sonninia propinquans				
·		(6a)				
WITCHELLIAE	WITCHELLIA GRIT	Limestone with				
•	NOTGROVE FREESTONE	Witchellia (6b)				
		and				
		Sonninia cf.				
		<u>fissilobatum</u> -				
SONNINIAE	GRYPHITE GRIT	ovalis, mainly at				
		the bottom. (6c)				
	<u>ovalis</u>	Sandy limestone				
÷		with <u>S</u> . <u>ovalis</u>				
		(7-8)				
	BUCKMANI GRIT	Sandy parting with				
POST	with	Terebratula				
DISCITAE	Terebratula	contonensis (0)				
	DUCKMANT					
	LOWER TRIGONIA	Sandy limestone				
DISCITAE	CDTO	with				
	GRTL	Hyperlioceras (10)				
CONCAVI	SNOWS HILL CLAY	Sandy stone				
		(11-13)				
	TILE STUNE etc.					
L	1					

Table 2.

A correlation of parts of the Middle Inferior Oolite by S.S. Buckman ; modified from Buckman (1895),table 2, with bed numbers inserted after.Buckman (1893).

would be created by its use in a position very different to that originally intended by Buckman and Spath. However if there were any strong objections to the double use of W. laeviuscula, Shirbuirnia stephani, inadvertently validated as a Zonal index by Arkell (ARKELL, 1933, plate 33), prior to Spath's Trigonalis Subzone (SPATH, 1936), would make a good alternative choice. The horizon below the 'fossil bed' at Sandford Lane yields a fauna which may be correlated with that obtained from the 'lower white iron-shot' of Dundry and the Gryphite Grit of the Cotswolds. This was recognised at an early stage by Buckman (see Tab. 2), who later confused the issue by wrongly correlating the fauna from this lower horizon, with that artificially split from the basal third of the Sandford Lane 'fossil bed', and combining both within the Shirbuirnia hemera; Spath's Trigonalis Subzone. This lower horizon, originally designated by Buckman as the 'Sonninia fissilobatum - ovalis horizon', (BUCKMAN, 1895) can take a pre-existing subzonal index, Sonninia (Fissilobisceras) ovalis (Qu. emend. S.B. 1893) as used by Oechsle in South Germany, (OECHSLE, 1958, p.182, with Lectotype designation). This follows current French practice, (GABILLY, 1964; CONTINI, 1970; GABILLY et al. 1971) and is in line with Buckman's use of S. ovalis as a hemeral index, (BUCKMAN, 1930 in 1909-30).

The use of a Discites Zone produces few problems. The fauna of this Zone is easily recognisable, in spite of the fact that further, more detailed stratigraphic division of it may be possible (BAYER, 1969, p.35). Problems arise in the recognition of an additional subzone above the Concavum <u>B. str.</u>, variously called the 'Amplectens faunule' or subzone (WESTERMANN, 1969; BAYER, 1970) and the Formosum subzone, (CONTINI, 1970; CABILLY <u>et al.</u> 1971). There are indications



Figure 2.

A correlation of certain beds of the Inferior Oolite in the Sherborne area, north Dorset. that the middle part of the Bradford Abbas 'fossil bed' yields a fauna equivalent to those from this 'so called subzone'. However until a detailed revision of the Graphoceratid ammonites has been completed, no final decision should be made on this point.

7. THE SUCCESSIVE AMMONITE FAUNAS OF THE SAUZEI, LAEVIUSCULA, DISCITES AND CONCAVUM ZONES IN SOUTHERN ENGLAND

The following can only be a summary of work, which will be published in greater detail elsewhere. Beds of this age in Southern England tend to be both highly condensed and laterally impersistent. This attempt to establish a series of consecutive ammonite faunas can only be considered provisional, as it is often difficult to establish any correlation between isolated exposures. In this respect, the beds of this age in the Sherborne district of Dorset were particularly difficult to place in sequence until an exposure at Clatcombe Farm, (BUCKMAN, 1893, section xiv) was examined. This enabled a correlation to be made between the Blue Bed of Oborne (op. cit. section xv, bed 10) and the Sandford Lane 'Fossil Bed', (on. cit. section ix, bed 6), which was at variance with the results obtained by Buckman (see fig. 2 for further details). The faunal sequence established by this revised correlation was fully upheld however by the continuous series of faunas which it has recently been possible to collect from Dundry Hill, near Bristol (fig. 3). Certain limits were needed to restrict the extent of this study and the prominent unconformities at the base of both the Humphriesianum and Concevum Zone rocks in the Sherborne district formed natural lines of division.

		EMILEIA POLYSCHIDES	E. POLYMERA	E.(OTOI TES)c(SAUZEI	KUMATOSTEPHANUS SP.	LABYRINTHOCERAS SA	ZWITCHELLIA HEBRIDICA	7 <u>W(PELEKODITES) SULCATA</u>	PROTOECOTRAUSTES sp.	W. FALCATA	W. LAE VIUSCULA	W. SPINIFERA	W. cf SUTNERI	W(PELEKODITES) MACER	FROGDENTES Sp.	MOLLISTEPHANUS Sp.	SHIRBUIRNIA of STEPHANI	W. PAVIMENTARIUS	W. ROMANOI DE S	EUHOPLOCERAS	ACAN THERUM	SONNINIA OVALIS	W. CONNATA	EMILEITES (TRILOBITICERAS)		
SAUZEI	ZONE	ļ	ł	ł	Ţ		ł	Ţ	8																U.	
	LA	-		-	.Y .	-		<u> </u>		Ö		 		Š	Ī	-	 	-	~~	-	-	-	-		-	20
ZONE	VIUSCU									1						ł		ļ							-	45
SCULA	LAE	-			• -	-	-	•			. 0.	• ••		ٳ	-	•	 -	-	¢	- <	ļ.	Ą	•	- 4	-	·65
LAEVIU	OVALIS															•							ł			
	-		-	-	-	-				•		-	- •		• •	• •••	-		• , •			-	-	- 6	- •	•95

Figure 3

The stratigraphic distribution of certain ammonites from the South Main-road quarry, Dundry Hill, Avon. The O.Ocm. line marks the planed surface at the top of the Lower Bajocian rocks. The ammonite records which are given here are all based on material from my own collection, collected in situ, unless otherwise stated. In the following lists:-

T = Topotype
C = Chorotype
M = Macro-conch
m = Micro-conch

a/ The Sauzei Zone

The fauna of part of the Irony Bed (GATRALL, JENKYNS & PARSONS, 1972, p.81) to the west of Sherborne, together with that from the top half of the Sandford Lane 'Fossil Bed' and its equivalent at Clatcombe (BUCKMAN, 1893, pp.493-5) are all identical to those from the Couche Verte and Blaukalke of Normandy and Schwabia; as is that from the upper part of the Brown Iron-shot Bed of Dundry (BUCKMAN & WILSON, 1896). The Pecten Bed of the Bruton district of Somerset (RICHARDSON, 1916), the Phillipsiana/Bourguetia Beds of Cleeve Hill near Cheltenham (BUCKMAN, 1897; BUCKMAN in RICHARDSON, 1904) and the upper part of the Red Beds of South Dorset (SENIOR, PARSONS & TORRENS, 1970) have also yielded typical Sauzei Zone faunas.

The following ammonites come from the most extensive Sauzei Zone fauna collected during the course of this work, that is from the top of the Sandford Lane 'Fossil Bed'. This bed was found to be divisible into only two horizons; at the base a light yellow-green coloured limestone with yellow ooliths and green glauconite grains and at the top a deep blue coloured 'iron-shot' limestone, which Weathers to a brown colour; hence the third matrix and hemera of Buckman. These two highly distinctive matrices are separated by an irregular parting, which in fact marks the position of an extensive 'hard-ground', on to the top of which the blue 'iron-shot' limestone has been cemented. The 'hard-ground' shows half eroded ammonites, limonite- and serpulid-encrustation and is surmounted by a conglomerate of soft, light grey coloured limestone lithoclasts.

Sandford Lane ST628,179*1

(BUCKMAN, 1893, section ix, bed 6a)

Emileia (Emileia) bulligera S. Buckman, T.M.

E. (E.) pseudocontrahens Maubeuge, M., J. Whicher Collection.

E. (Otoites) fortis (Westermann), m.

E. (O.) sauzei (d'Orbigny, pl.139 non neotype), m.

E. (O.) cf. delicata (Buckman), m.

Kumatostenhanus cf. kumaterus S. Buckman (J. Whicher

collection), T.M.

?Normannites aff. kialagvikensis Imlay, m.

?N. aff. sauzeiformis Westermann, m.

Labyrinthoceras meniscum (Waagen), M.

Sphaeroceras manselii (J. Buckman), C, M & m.

Skirroceras kalus (S. Buckman), T.M.

Sonninia felix (S. Buckman), T.M.

S. cf. propinquans (Bayle), M.

S. cf. corrugata (J.de C. Sowerby),? M.

Witchellia (?Witchellia) hebridica (Morton), M.

W. (Pelekodites) sulcata (S.Buckman), C,m.

Lissoceras semicostulatum S. Buchman, T.M.

<u>Strigoceras</u> (<u>Strigoceras</u>) <u>strigifer</u> (S. Buchman), T, M. <u>Toxamblyites</u> aff. arcifer S. Buchman,? m.

The conglomerate at the base of this bed has a thicker lateral equivalent at Clatcombe Farm (see fig 2, = bed 4, EUCKMAN, 1893, p.498). This stratigraphically lower horizon yields ammonites which are too poorly preserved for anything other than generic identification. The next ammonite fauna is difficult to place, since it is transitional in character between the Sauzei and Laeviuscula Zones. Taking into account the wealth of <u>Witchellia</u> present and the absence or rarity of many species characteristic of the Sauzei Zone, this has been placed in the Laeviuscula Zone.

b/ The Laeviuscula Zone and Subzonei. The 'Green grained marl' of Oborne

The highly glauconitic marl bed of the Oborne area, Sherborne, Dorset (BUCKMAN, 1893, Frogden quarry, p.500, bed 9) has yielded extensive and superbly preserved faunas, which, apart from specimens from the lower part of the main 'Iron-shot' bed at Dundry (BUCKMAN & WILSON, 1896, p.681, bed 6), have no other English equivalent. The following collection is from the Marl Bed at a recent temporary exposure at Oborne, near Sherborne, Dorset (WHICHER & PALMER, 1971) although a similar fauna was obtained from the South Main-road quarry, Dundry, near Bristol, (BUCKMAN & WILSON, 1896, p.691, bed 5)

Oborne Wood, ST648,188

(Equivalent to bed 9, Frogden quarry) (BUCKMAN, 1893)

Emileia (E.) brocchii (J. Sowerby), ?T.M.

E. (E.) bulligera, J. Whicher collection, M.

E. (E.) polyschides, M.

E. (Otoites) contracta (S. Buckman non Sow.), m.

Frogdenites spiniger S. Buckman, T,M & m.

F. cf. profectus S. Buckman, m.

Mollistephanus aff. mollis S. Buckman, M.

Skirroceras aff. kalus S. Buckman, M.

<u>S. leptogyrale</u> S. Buckman (including <u>S. macrum</u> Buckman <u>non</u> Quenst.), M.

Bradfordia cf. inclusa S. Buckman.

'Ambloxyites' amblys S. Buckman, M.

Protoecotraustes spiniger (S. Buckman), m.

Papilliceras arenatus (Quenst. emend. Buckman), N.

Shirbuirnia superba (S. Buckman), T, M.

S. cf. trigonata (Quenst. emend. Dorn), M.

Witchellia (Witchellia) falcata S. Euckman, T, N.

W. (W.) glauca S. Buckman, T, M.

W. (W.) actinophora S. Buckman, T, M.

The above three 'species' form part of a highly variable plexus, varying from smooth involute to coarsely ribbed evolute morphotypes.

W. (W.) cf. patefactor S. Buckman, M.

W. (W.) plena (S. Buckman), T.N.

W. (Pelekodites) aurifer (S. Buckman), m.

W. (P.) maora (S. Buckman), T, m.

ii. The basal half of the Sandford Lane 'Fossil-bed'

The 'green grained marl' of the Oborne district grades down into a hard glauconitic limestone (= bed 10, Frogden quarry, BUCKMAN, 1893), which contains a fauna very similar to that obtained from the basal part of the Sandford Lane 'Fossil-bed'. Comparable faunas have also been collected from the basal part of the Pecten Bed of the Bruton district, Somerset (RICHARDSON, 1916, p.495, bed 4), the Witchellia Grit of the Cotswolds (BUCKMAN in RICHARDSON, 1904) and from the middle of the 'Iron-shot' beds on Dundry Hill, near Bristol (BUCKMAN & WILSON, 1896, p.681, bed 8). Due to the intractable nature of the Sandford bed it has not proved possible to collect topotypes of all the ammonite species figured by Buckman from this horizon. However the matrix of specimens from this horizon is so characteristic as to preclude any error in the identification of their horizon.

It cannot be over stressed that it has proved totally impossible to separate two faunas from the basal half of the 'Fossil-bed'; <u>Witchellia and Shirbuirnia are inextricably mixed together</u>. Indeed one of the highest ammonites found in this portion of the bed was a specimen of <u>Shirbuirnia stephani</u> (S. Buckman), CP 1930, at 0.31m. below the top of the 'Fossil-bed'.

Sandford Lane, ST628,179

(BUCHMAN, 1893, bed 6c)

Emileia (E.) <u>catamorpha</u> S. Buckman, J. Whicher collection, T, M.

E. (E.) cf. polyschides (Waagen), J. Whicher collection, N.
E. (Otoites) contracta (S. Buckman non Sow.), m.
E. (O.) cf. delicata (S. Buckman), T, m.
<u>Mollistephanus (M.) cf. mollis S. Buckman, T, M.</u>
<u>Trilobiticeras (Emileites) liebi (Maubeuge), M.</u>
<u>Bradfordia of. liomphala S. Buckman</u>

<u>Euhoploceras acantherum</u> (S. Buckman), T, N. <u>Shirbuirnia stephani</u> (S. Buckman), T, M. <u>Sonninia gingensis</u> (Waagen), M. <u>Witchellia (W.) gelsina</u> (S. Buckman), T, M. <u>W. (W.) plena</u> (S. Buckman), M. <u>W. (W.) sutneri</u> (Branco), M. <u>W. (W.) glauca</u> S. Buckman, M. <u>W. (W.) patefactor</u> S. Buckman, M.

<u>W.</u> (<u>W.</u>) cf. <u>pavimentaria</u> (S. Buckman), N.

The last three 'species' form part of a plexus of variation, similar to that of the 'Marl Bed' fauna, but they are consistently smoother and more evolute.

W. (Pelekodites) macra (S. Buckman), m.

c/ The Ovalis Subzone of the Laeviuscula Zone

The beds immediately below the Sandford Lane 'Fossil-bed' yield large members of the <u>Sonninia ovalis</u> (Quenst. emend. Buckman) group and evolute Witchellids, which enable a correlation to be made with a bed on Dundry Hill yielding a similar fauna, that is the 'Lower white iron-shot' bed (BUCKMAN & WILSON, 1896, p.676, beds 4-8). Other areas which have produced similar ammonite faunas include the Bruton district of Somerset (RICHARDSON, 1916, p.495, bed 4b), the 'Gryphite Grit' of the Cotswolds (BUCKMAN, 1895, p.393) and Seavington St. Mary, Somerset (PARSONS & TORRENS in TORRENS, (Editor) 1969, p.A27, bed 4a). The following ammonite fauna comes from the 'Lower white iron-shot' bed of Barns Batch Spinney, Dundry Hill.

> Barns Batch Spinney, ST557, 659, (BUCKMAN & WILSON, 1896, p.689) bed 2.

Sonninia (Fissilobisceras) ovalis (Quenst. emend.

Buckman), M.

<u>Witchellia</u> (<u>W.</u>) <u>albida</u> (S. Buckman), T, M.
<u>W.</u> (<u>W.</u>) <u>connata</u> (S. Buckman), M.
<u>W.</u> (<u>W.</u>) <u>romanoides</u> (Douvillé), M.
<u>W.</u> (<u>W.</u>) cf. <u>sutneri</u> (Branco), M.
<u>W.</u> (<u>Pelekodites</u>) <u>pelekus</u> (S. Buckman), T, m.
<u>W.</u> (<u>P.</u>) cf. <u>macra</u> (S. Buckman), m.
<u>Bradfordia</u> cf. <u>inclusum</u> S. Buckman
<u>Strigoceras</u> (<u>S.</u>) <u>compressum</u> (S. Buckman), T, M.
<u>Docidoceras</u> (<u>D.</u>) <u>cylindroides</u> S. Buckman, M.
<u>Trilobiticeras</u> (<u>T.</u>) aff. <u>punctum</u> (Vacek), m.
<u>T.</u> (<u>Emileites</u>) <u>liebi</u> (Maubeuge), m.
<u>Emileia</u> (<u>Otoites</u>) <u>sauzei</u> (Douvillé <u>non</u> d'Orb.), m.

d/ The Discites and Concavum Zones

The most important single locality for the study of the Discites and Concavum Zones in England is undoubtedly the Bradford Abbas district, near Sherborne, Dorset. Here a relatively thin (0.50 -70m.) highly condensed limestone bed yields abundant ammonite faunas, which were minutely studied by Buckman (1893; 1887-1907). Above this Bradford Abbas 'Fossil-bed' there is a large stratigraphic break. The junction between the Discites and Laeviuscula Zones can however be studied at Dundry (Castle Farm and Barns Batch Spinney, BUCKMAN & WILSON, 1896), Lusty quarry, Bruton, Somerset (RICHARDSON, 1916, p.495) and Frogden quarry, Oborne, Dorset (BUCKMAN, 1893). Whilst all of the quarries made famous by J. and S.S. Buckman in the Bradford Abbas district are now filled in, the extensive section of the 'Fossil-bed' in the Bradford Abbas Railway-cutting is still available for study. The following faunas are all <u>in situ</u> collections from this locality and bed. It is to be noted that the Graphoceratid ammonites, like the Witchellids, are extremely variable and until a full systematic revision is made of them, any specific names given merely represent single morphotypes.

Buckman mainly on lithological criteria, divided the Bradford Abbas 'Fossil-bed' into two horizons; an upper black stained 'ironshot' limestone and a lower soft brown 'iron-shot', which he included in the Discites and Concavum hemerae respectively. In the Bradford Abbas Railway-cutting the 'Fossil-bed' was found to be divided by two marl partings into three approximately equal components. The upper, main parting, which separates the black stained limestone from the rest is consistent in thickness over the length of the cutting, whilst the second, lower parting is less persistent, but it is still traceable over most of the cutting. These two partings enabled three successive ammonite faunas to be collected.

i. The top third of the Bradford Abbas 'Fossil-bed'

Whilst the following fauna comes from the Bradford Abbas 'Fossil-bed', exactly similar ammonites have been found in the lower exposure at Frogden quarry (BUCKMAN, 1893), in the 'Snuff-box' bed of South Dorset (SENIOR, PARSONS & TORRENS, 1970), in the lower Trigonia Grit of the Cotswolds (BUCKMAN, 1895) and on Dundry (BUCKMAN & WILSON, 1896)

Bradford Abbas Railway-Cutting, ST594, 145 (Equivalent to bed 7, BUCKMAN, 1893, p.485)

Braunsina aspera 5. Buckman

<u>Graphoceras</u> (<u>Graphoceras</u>) cf. <u>apertum</u> (S. Buckman) <u>G. (?G.) hamatum</u> (S. Buckman) <u>G. (?L.) mecticostata</u> (S. Buckman) <u>Darellia</u> cf. <u>coela</u> (S. Buckman) <u>Darellia</u> cf. <u>coela</u> (S. Buckman) <u>D. cf. laxa</u> (S. Buckman) <u>Hyperlicceras</u> (<u>Hyperlicceras</u>) <u>walkeri</u> (S. Buckman) <u>H. (H.) licdiscites</u> S. Buckman <u>Reynesella piedes</u> S. Buckman <u>Nannoceras nannomorphum</u> S. Buckman <u>Nannolytoceras cf. liccyclum</u> (Brasil) <u>Euhoploceras acanthodes</u> (S. Buckman) <u>E. submarginata</u> (S. Buckman) <u>Trilobiticeras (Trilobiticeras) trilobitoides</u> S. Buckman <u>T.(Emileites</u>) aff. <u>liebi</u> (Maubeuge)

ii. The middle of the Bradford Abbas 'Fossil-bed'

The middle of the 'Fossil-bed' yields a fauna intermediate in character between the previous and that from the basal part of the 'Fossil-bed' (Concavum Subzone <u>sensu stricto</u>). Similar faunas have been collected from the top of the Iron-shot' Bed at Horn Park quarry, south Dorset (SENIOR <u>et al.</u>, 1970) and from bed 3, Seavington St. Mary (PARSONS & TORRENS in TORRENS (Editor), 1969).

> Bradford Abbas Railway-Cutting <u>Graphoceras (G.) apertum</u> (S. Buckman) <u>G. (G.) scriptitatum</u> S. Buckman <u>G. (Ludwigella) compactum</u> (S. Buckman) <u>G. (L.) limitatum</u> S. Buckman

<u>G. (L.) subrudis</u> S. Buckman <u>Bradfordia liomphala</u> S. Buckman <u>Eudmetoceras amplectens</u> S. Buckman <u>Euhoploceras acanthodes</u> (S. Buckman) <u>E. simplex</u> (S. Buckman) <u>Haplopleuroceras subspinatum</u> (S. Buckman) <u>Trilobiticeras (T.) cf. punctum</u> (Vacek)

iii. The lower part of the Bradford Abbas 'Fossil-bed'

The lower part of the 'Fossil-bed', together with the main 'iron-shot' bed at Horn Park quarry, Dorset (SENIOR <u>et al.</u>, 1970), bed 2 Seavington St. Mary (PARSONS & TORRENS, in TORRENS (Editor), 1969), and the lower limestones at Dundry (BUCKMAN & WILSON, 1896) have all yielded a similar fauna, that of the Concavum Subzone <u>sensu</u> <u>stricto</u>.

> Bradford Abbas Railway-Cutting <u>Graphoceras</u> (G.) <u>concavum</u> (Sow.) <u>G. (G.) magna</u> (S. Buckman) <u>G. (G.) sublineata</u> (S. Buckman) <u>G. (Ludwigella) attenuata</u> S. Buckman <u>G. (L.) cornu (S. Buckman)</u> <u>G. (L.) flexilis S. Buckman</u> <u>G. (L.) micra S. Buckman</u> <u>G. (L.) rotabilis (S. Buckman)</u> <u>G. (L.) stigmosum S. Buckman</u> <u>Bradfordia liomphala S. Buckman</u> <u>Euhoploceras cf. crassiformis</u> (S. Buckman)

8. THE CORRELATION OF SCHWABIA AND NORMANDY

The correlation of the 'Couche verte' and 'Blaukalk' with the Sauzei Zone, has already been discussed in some detail, and presents no problems; unfortunately the same cannot be said of the subjacent strata.

a/ <u>Schwabia</u>

Correlation of the beds between the 'Blaukalk' and the 'Oberer Donzdorf' sandstone (= Concavum Zone), with the British succession present considerable problems for the following reasons. Firstly these Schwabian horizons are poor in ammonites, secondly many typical British ammonites, such as <u>Witchellia</u> are rare or absent and thirdly there exists a prominent unconformity, showing a marked discordance, which at some localities cuts out several beds, (The 'Unter Wedel Sandstein', see Fig. 4). Exposures in the Gingen area, notable in the past for its 'Sowerbyi Zone' ammonites (WAAGEN, 1867, DORN, 1935, OECHSLE, 1958), are now very poor. The following is an undescribed stream section at Grunbach near Goppingen, and less than 10 km. from Gingen, which had a very similar succession, (OECHSLE, 1958, with extensive list of ammonites etc.).

i. Stream section in a wood just to the east of Grunbach

(4) "Mittel -X- tone"

A blue tenacious clay limonite streaked at the base, and with intercalations of thin, sandy limestone ('Wedel Sandstein') in the upper part.

seen to 1.0 m. ++

(3) A conglomerate consisting of dark blue 'iron-shot'

limestone lithoclasts, heavily limonite-encrusted and set in soft blue limestone matrix, which becomes harder and more indurated towards the base.

There is a fairly prolific remanié ammonite fauna, mainly distorted, broken and limonite-encrusted internal moulds, resting on the irregular and limonite coated basal erosion surface.

Shirbuirnia cf. stephani, CP. 1810.

Shirbuirnia spp.

0.15m.

prominent 'hard-ground' ----

(2) "The Sowerbyi Oolith"

A hard, blue, heavily 'iron-shot' limestone, with a tendency to be conglomeratic; very hard at the top and softer below, with a lighter, more grey coloured matrix.

Euhoploceras cf. adicra (Waag.), CP. 1818.

(1b) A thin sandy parting

(1a) "The Oberer Donzdorf, or Concavum Sandstein"

A massive, brown sandstone, heavily ironstained and with some "box structure" weathering.

seen to 1.0 m. +

0.45m.

0.04m.

The "Sowerbyi Oolith", (Discites Zone) is undoubtedly the source of Waagen's type specimens of <u>Euhoploceras adicra</u> and <u>E. polycantha</u>. The others, apart from <u>Hyperlioceras discites</u>, came either from the suprajacent remanie bed or the 'Unter Wedel sandstein", which in the



Figure 4

A schematic correlation of parts of the Lower ond Bajocian/Aalenian, in the Schwabian Albe, southern Germany. past have all been grouped, together with the 'Sowerbyi Oolith', in the 'Sowerbyi bank' or 'Schichten'.

b/ The Wutach

There is a slightly more expanded sequence in the Wutach, with several courses of very sandy limestone (the 'Unter Wedel Sandstein') coming in below the erosion plane, and above the 'Sowerby' Oolith' equivalent, (BAYER, 1970; BUCK, HAHN & SCHADEL 1966; and HAHN. 1971). The 'Sowerbyi Oolith' equivalent (Eichberg, Fig. 4), yields Hyperlioceras cf. rudidiscites (CP.2397) (see RIEBER, 1963, p.6) and brachiopods, Sphaeroidothyris eudesiana and 'Rhynchonella' forbesi, identical to those from southern England. The 'Unter Wedel Sandstein' has produced several Sonninia (Fissilobisceras) cf. ovalis (Bayer Col.), whilst resting on the overlying erosion plane are poorly preserved, remanic ammonites which can be referred to the genus Shirbuirnia, (cf. S. trigonata (Qu. emend. Dorn), Scheffhue, CP 1820). The correlation of the 'Wedel Sandstein' with the Ovalis subzone is further strengthened by the isolated occurrence of Trilobiticeras (Emileites) cf. liebi (Maub.), above the highest occurrence of Hyperlioceras, (BAYER, 1968 and pers. comm.).

To the west, at the old Iron-stone mines at Ringsheim, similar sections are visible to those of the Wutach, (BAYER, 1970) where a reduced 'Unter Wedel Sandstein', yielding <u>Fissilobisceras</u> <u>ovalis</u> (CP.1821) overlies beds containing <u>Hyperlioceras</u> spp. and <u>Euhoploceras</u> cf. <u>polycantha</u>

The correlation of the Schwabian lower Bajocian is summarised in a general way in Fig. 4.

c/ Normandy

Below the 'Couche Verte' on the Normandy coast, there is a series of light grey, slightly sandy limestones, rich in black chert bands - the Malière. At St. Honorine des Pertes, the upper part of the Malière has yielded <u>Kumatostephanus</u> cf. <u>triplicatum</u> (CP.1600 & 1601), and <u>Emileia (E.) greppini</u> Maubeuge, (N. Morton col.), and is thus still Sauzei Zone in age. At lower horizons the Malière is less fossiliferous, but past records suggest that it ranges down, at least as far as the Murchisonae Zone, (RIOULT and GABILLY, 1967).

The large inland quarries at May-sur-Orne show numerous exposures of the Iron-shot Aalenian/Bajocian resting on steeply dipping palaeozoics. The Murchisonae - Concavum Zones are well represented as are the Upper Bajocian Zones, but the intermediate horizons tend to be very lenticular (BIGOT, 1900). From the work of Munier-Chalmas (1892) Brasil (1895), Bigot (1900) and more recently Ricult and Gabilly (1967) it seems likely that at least the Ovalis subzone has been represented by a small lens of 'iron-shot' limestone. The preservation and mode of deposition of the Lower, Bajocian in Normandy thus follows a very similar pattern to that of South Dorset.

9. CONCLUSION

The following is a resumé of the Standard Zonal scheme for parts of the Aalenian-Bajocian Stages, including any changes made necessary by the present work.

aphoceras (Graphoceras) concavum (J. Sow.)
Hudleston (1887, in 1887-96) restricted by
Buckman (1889, in 1887-1907).
North Dorset.
Basal part of the Bradford Abbas 'Fossil-
bed!.
The wealth of <u>Graphoceras</u> s. str.,
particularly <u>G</u> . (<u>G</u> .) <u>concavum</u> , <u>G</u> . (<u>G</u> .)
magna, G. (Ludwigella) cornu and G. (L.)
stigmosum, which is typical of the Concavum
Zone sensu stricto. There may be a possibi-
lity of subdivision of this unit, with a
higher horizon characterised by Eudmetoceras
and <u>Haplopleuroceras</u> .

Ъ/	The	Zone	of Hyperlio	ceras (Hyp	erliocera	s) <u>discites</u>	(Waa	gen)
				A second s	a tha an an an an an		•.	

Authors-	Buchman (1893).						
Type area:-	North Dorset.						
Type Horizons-	Top of the Bradford Abbas 'Fossil-bed'.						
Characteristic Fauna:-	This Zone is characterised by the abundance						
	of the genus <u>Hyperlioceras</u> , particularly <u>H</u> .						
	(<u>H.) walkeri - rudidiscites</u> S. Buckman group,						
	as well as by the genus <u>Reynesella</u> and by						
	Trilobiticeras trilobitoides.						

c/ The Zone of <u>Witchellia</u> (Witchellia) <u>laeviuscula</u> (J. de C.

Authors-Haug (1894). Type area:-To be designated. Type Horizon:-To be designated. Characteristic Fauna:-The abundance of the dimorphic pair <u>Witchellia</u> (W.) and <u>W.</u> (Pelekodites) and the species Trilobiticeras (Emileites) liebi (Maubeuge) characterise this horizon. i. The Subzone of Sonninia ovalis (Quenst. emend. Buckman) Authors-Oechsle (1958). Type area:-The Schwabian Albe. The 'Unterer Wedelsandstein'. Type Horizon:-Characteristic Fauna:-This horizon is typified by the presence of large smooth Sonninids, such as S. ovalis, and by the appearance of T. (E.) liebi and the genus Witchellia.

ii. The Subzone of Witchellia (Witchellia) laeviuscula (J. de C. Sow.)

Author:-Type area:-

Still to be designated.

Haug (1894), restricted Spath (1936).

Type Horizon:- Still to be designated.

Characteristic Fauna:- This subzone is characterised by the more involute, costate Witchellids and by the genus <u>Shirbuirnia</u>. There is a possibility that this unit may be subdivided on the basis of the distribution of certain species of <u>Witchellia</u> and <u>Shirbuirnia</u>, the higher

Sow.)

horizon being defined by the appearance of the genus <u>Frogdenites</u>.

d/ The Zone of Emileia (Otoites) sauzei (d'Orb.)

Authors-

Oppel (1856 in 1856-8).

Type area:-

Type Horizon:-

Schwabian Albe. The Blaukalke.

Characteristic Fauna:-

This has already been given in some detail (section 2), but the base of this Zone is defined by the appearance of the genera <u>Kumatostephanus</u> and <u>Labyrinthoceras</u> and the species <u>Sphaeroceras manselii</u> and <u>Witchellia</u> hebridica.

10. REFERENCES

Arkell, W.J., 1933 (Reprinted 1970)

The Jurassic System in Great Britain. Oxford Univ. 681 p.41 pl.

in Arkell, W. & Playford, P.E., 1954

The Bajocian Ammonites of Western Australia. Phil. Trans. Roy. Soc. London 237 (B): 547-604. pls. 27-40.

1956

The Jurassic Geology of the World. Oliver & Boyd, Edinburgh, 806 p. 46 pls.

Bayer, U. 1968

Docidoceras cf. <u>liebi</u> Maubeuge aus dem Unteren Bajocium des Wutachgebietes. Stuttgt. Beitr. z. Naturk. <u>183</u>: 1-3.

1969

Die Gattung <u>Hyperlioceras</u> Buckman. Jber. u. Mitt. oberrh. geol. Ver. <u>51</u>: 31-70.

1970

Das Profil des Erz-Tagebaus Ringsheim (Ober- Aalenium/ Unter-Bajocium). N. Jb. Geol. Paläont. Mh. Jg. 1970, H.5: 261-9.

Bigot, A. 1900

Normandie. Livret Guide des excursions du VIII Congres Géologique International. Paris. 33 p.13 fig.

Brasil, L. 1895

Observations sur le Bajocien de Normandie.

Bul. Lab. Geol. Fac. Sci. Caen, 2 : 222-48.

Buck, E., Hahn, W., & Schadel, K. 1966

Zur stratigraphie des Bajociums und Bathoniums der

Schwäbischen Alb. Jh. geol. Landesant. Baden-Nürttemberg, 8 : 23-46.

Buckman, S.S. 1887-1907

A Monograph of the Ammonites of the Inferior Colite Series. Palaeontogr. Soc. cclxii + 456 p. 127 pls.

_____ 1893

The Bajocian of the Sherborne district; its relation to subjacent and superjacent strata. Q. Jl. geol. Soc. Lond. <u>49</u>: 479-522.

____ 1895

The Bajocian of the mid-Cotteswolds. Q. Jl. geol. Soc. London. <u>51</u> : 388-462.

1897

Deposits of the Bajocian Age in the Northern Cotteswolds; the Cleeve Hill Plateau. Q. Jl. geol. Soc. Lond. 53 : 607-629.

_____1904

Ammonites soverbyi (Killer m.s.) J. Sowerby 1821.

Palacontologia Universalis Pl. 52.

_ 1910

Certain Jurassic (Lias-Oolite) strata of South Dorset. Q. Jl. gool. Soc. Lond. <u>66</u> : 52-89. Buckman, S.S. 1915 in

Fox-Strangways, C. & Barrow, G.

Whitby and Scarborough. 2nd edit. Mem. Geol. Surv. Great Britain. iv + 144 p.

_____ 1909–1930,

'Type Ammonites' London & Thame, vols. 1-7: 790 pls.

_ & Wilson, E. 1896

Dundry Hill: its upper portion, or the beds marked as Inferior Oolite in the maps of The Geological Survey. Q. Jl. geol. Soc. Lond. <u>52</u>: 669-720.

Contini, D. 1970

L'Aalénien et le Bajocien du Jura Franc - comtois, etude stratigraphique. Ann. Sci. Univ. Besançon, 3e ser. - Géologie. fasc. <u>11</u>: 1-204.

Dangeard, L. 1940

Sur les diverses manières d'être de la glauconie dans la couche verte du Bajocien Normand. C.R. Acad. Sci. <u>211</u>: 264-266.

Donovan, D. & Hemingway, J. 1963

Lexique Stratigraphique International Europe Fasc. 3a -England, Wales & Scotland. Part 3a X - Jurassic. Paris, 394 p.

Dorn, P. 1935

Die Hammatoceraten, Sonninien, Ludwigien, Dorsetensien und Witchellien des süddeutschen insbesondere fränkischen Doggers. Palaeontographica <u>82</u> (A): 1-124, pls. I-XXIX Douville, H. 1885

Sur quelques fossiles de la zone à <u>Ammonites sowerbyi</u> des environs de Toulon. Bul. Soc. geol. Fr. (3) <u>13</u>: 12-44, 3 pl.

Gabilly, J. 1964

Le Jurassique inférieur et moyen sur le littoral Vendéen. Trav. Inst. Géol. Anthropol. Préhist. Fac. Sci. Poitiers. 5: 65-107.

Les Zones du Bajocien. C.R.S. Soc. geol. Fr. : 85-88.

Gatrall, M., Jenkyns, H. & Parsons, C., 1972

Limonitic concretions from the European Jurassic with particular reference to the 'snuff-boxes' of Southern England. Sedimentology <u>18</u>: 79-103.

Hahn, W. 1971

Der Jura, p.117-133, in 'Die Wutach' - Erd- und Landschaftsgeschicte des Wutachgebeites. Freiburg-i-Breslau. 161 p., 40 figs., 10 pl.

Haug, E. 1891

Les Châines Subalpines entre Gap et Digne. Bul. Serv. Carte géol. Fr. <u>111</u>, No. 21: 199 p.

1893

Etude sur les ammonites des etages moyen du System Jurassique. Bul. Soc. Geol. Fr. 3e ser. <u>20</u>: 277-333, 3 pl. 1894

Le System Jurassique. in La Grande Encyclopédie. Vol. 21, Paris. p.322-331.
Haug, E. 1900

Des Basses Alpes. Livret guide des excursions de VIII congrès Géologique International, Paris. : p.1-28.

1910

Traité de Géologie. Vol. 2, pt. 2 (1st ed.), (Colin, Paris): 929-1396.

Hudleston, W.H., 1887-96

A Monograph of the British Jurassic Gasteropoda. pt. I, the Gasteropoda of the Inferior Oolite. Palaeontograph. Soc. 514 p., 44 pl.

Morton, N., 1965

The Bearreraig Sandstone Series (Middle Jurassic), of Skye and Raasay. Scot. J. Geol. 1: 189-216.

Munier-Chalmas, 1892,

Etude préliminaire des terrains Jurassiques de Normandie. C.R.S. Soc. Géol. Fr. (3) 20: 161-170.

Oechsle, E., 1958

Stratigraphie und Ammonitenfauna der Sonninien-Schichten des Filsgebeits unter besonderer Berucksichtigung der Sowerbyi-Zone. Palaeontographica (A) <u>111</u>: 47-129, pls. 10-20.

Oppel, A., 1856-8

Die Juraformation Englands, Frankreichs und des südwestlichen Deutschlands. Jb. Ver. vaterländ. Naturkde. Württ. (Stuttgart). <u>12-14</u>: 1-837. Oppel, A., 1862-3

Ueber jurassische Cephalopoden. Pal. Mitt. Mus. k. Bay. staates, 3, : 127-266, pls. 40-74.

Orbigny, A. d'., 1842-51

Paléontologie Française; Terrains jurassiques, I Céphalopodes. Paris.

Pavia, G., & Sturani, C., 1968

Etude biostratigraphique du Bajocien des Châines Sub-Alpines aux environs de Digne. Boll. Soc. Geol. Italy, <u>87</u>: 305-316.

Quenstedt, F.A., 1843

Das Flözgebirge Württembergs mit besonderer Rücksicht auf den Jura. 1st edit. Tübingen, (Laupp): 558 p.

1886-7

Die ammoniten des Schwäbischen Jura, II. Der Braune Jura. Schweizerbart, Stuttgart: p.441-815, pls. 55-90.

Renz, C., 1904

Der Jura von Daghestan. Neu. Jb. Min. Geol. 2: 71-85.

Richardson, L., 1904

The Geology of Cheltenham. Sayer, Cheltenham: 303 p.

1916,

The Inferior Oolite and Contiguous deposits of the Doulting -Milborne Port district (Somerset). Q. Jl. geol. Soc. Lond., <u>71</u>: 473-520.

Rieber, H., 1963

Ammoniten und stratigraphie des Braunjura der Schwäbischen Alb. Palaeontographica (A) <u>122</u>: 1-89

Rioult, M., 1964

Le Stratotype du Bajocien. Coll. Jurassique Luxembourg, 1962. Publ. Inst. Grand-Ducal Luxembourg, Sect. Sci., Nat., Phys., Math. p.239-258.

Rioult, M., & Gabilly, J., (1967). 1974

Le Bajocien inférieur et le Toarcian supérieur sur les bordures du Massif amoricain. Prétirage, Coll. Luxembourg, 1967.

Senior, J.R., Parsons, C.F., & Torrens, H.S., 1970

New sections in the Inferior Colite of South Dorset.

Proc. Dorset Nat. Hist. Archaeol. Soc. 91: 114-9

Spath, L.F., 1936

On Bajocian Ammonites and Belemnites from eastern Persia

(Iran). Pal. Indica (N.S.) 22, mem. 3.: 1-21, 1 pl.

Stahlecker, R., 1926

Brauner Jura und Tektonik im Kirchheim - Uracher Vulkangebeit. N. Jb. Min. Geol. B-B. liv. B. : 157

Torrens, H.S., 1969

Field meeting in the Sherborne - Yeovil District. Proc. Geol. Assoc. 80 : 301-330.

(Editor) 1969

International Field symposium on the British Jurassic. Excursion Guide I - Dorset and South Somerset. University of Keele, 1969 : 71 p. Waagen, W., 1867

Über die Zone des Ammonites sowerbyi. Geogn. - Paläont. <u>1</u>: 511-607, pls. xxiv-xxix.

Wilson, V., Welch, F., Robbie, J. and Green, G., 1959

The Geology of the country around Bridport and Yeovil. Mem. Geol. Surv. G.B. 239p.

Welsch, J., 1905

Etude sur les Terrains du Poitou. Bul. Soc. geol. France, (4), <u>3</u> : 798-881.

Westermann, G.E.G., 1954

Monographie der Otoitidae. Beih. Geol. Jb. <u>15</u> : 364 p. 33 pls.

1964

Sexual Dimorpismus Bei ammonoideen und seine bedeutung Für die Taxionomie der Otoitidae. Palaeontographica (A) <u>124</u>: 33-73, pls. 6-9.

1969

The ammonite fauna of the Kialagvik formation at Wide Bay, Alaska Peninsula. pt. II - <u>Sonninia sowerbyi</u> Zone. Bul. Amer. Palaeont. <u>57</u>, pt.255 : 5-226, pls. 1-47.

& Riccardi, A., 1972

Middle Jurassic ammonite fauna and Biochronology of the Argentine - Chilean Andes. pt I. - Hildocerataceae. Palaeontographica (A) 140 : 1-116, pls. 1-31. Whicher, J. & Palmer, C., 1971

Preliminary description of a temporary exposure in the Middle Inferior Oolite, near Sherborne, Dorset. Proc. Dorset, Nat. Hist. Archaeol. Soc. <u>92</u>: 110-9.

IB. A STRATIGRAPHIC REVISION OF THE HUMPHRIESIAMUM/SUBFURGATUM ZONE ROCKS (BAJOCIAN STAGE, MIDDLE JURASSIC) OF SOUTHERM

ENGLAND

ABSTRACT

The development of the Standard Zonal scheme for this part of the Bajocian Stage is discussed, and a detailed, bed by bed, analysis of rocks of this age in Southern England (mainly Dorset) is given. The type horizons of Smith's, ^{fic} Sowerbys' and Buckman's ammonite species are discussed, and as many as possible of these are placed in their correct stratigraphic position. A detailed description of the stratigraphic distribution of the major ammonite species and genera is given. Finally the subzonal divisions of these Zones, which have been recognised in Southern England are redefined; that is the Romani, Humphriesianum and Blagdeni Subzones of the Humphriesianum Zone and Eanksi, Polygyralis and Eaculata Subzones of the Subfurcatum Zone.

	CONTENTS	
1.	INTRODUCTION AND ACKNOWLEDGEMENTS	67
2.	DEVELOPMENT OF THE STANDARD ZONAL SCHEEP	68
	2.1 The Zone of Stephanoceras (Stephanoceras) numphriesianum $(J. de C. Sow.)$	
	2.2 The Zone of <u>Strenoceras</u> (<u>Strenoceras</u>) <u>subjurcatum</u> (Zieten)	
3•	DISCUSSION OF THE SUB-ZONAL SCHEME	73
4•	THE HUMPHRIESIANUM AND SUBFURCATUM ZONE ROCKS OF SOUTHERN ENGLAND	75
	4.1 The Sherborne Region	
	4.2 The North-west Dorset - South Somerset Area	
	4.3 The South Dorset Area	
	4.4 The 'Cole Syncline'	
	4.5 The Doulting District	
5.	THE HORIZONS OF SMITH'S, SOMERBYS' AND BUCKMAN'S TYPES	107
	5.1 Smith's and Sowerbys' specimens	
	5.2 Buckman's specimens	
6.	CONCLUSIONS	111
	6.1 The geographical distribution of the Humphriesianum and	
	Subfurcatum Zones.	
	6.2 The Stratigraphic distribution of ammonites in the	
	Humphriesianum/Subfurcatum Zones.	
	6.3 The Standard Zonal Scheme	
7.	REFERENCES	117
	가 있었다. 이렇게 가지 않는 것 같아요. 이렇게 가지 않는 것은	

1. INTRODUCTION AND ACKNOWLEDGEMENTS

The Humphriesianum and Subfurcatum Zones are some of the most poorly represented of all the British Bajocian Zones. Humphriesianum Zone rocks are only found sporadically in the British Isles, South of the Mendips in Dorset and Somerset, in Yorkshire (Gross's record of <u>Stephanoceras humphriesianum</u> (Sow.) from the Lincolnshire Limestone cannot be confirmed (Gross, 1875, p.121)) and in the Inner Hebrides, Skye and Raasay in particular. The Subfurcatum Zone is even more restricted in its distribution, rocks of this age having only been found in a few isolated areas of Southern England. The supposed Subfurcatum Zone ammonites from Skye and Raasay are too fragmentary and poorly preserved to be certain of their true identity or horizon, (Norton, 1971). The Scottish occurrences of these two Zones have been the subject of recent detailed work, and little needs to be added to this, (Norton, 1965, 1971), whilst I aim to revise the ammonite faunas from the Yorkshire,Scarborough Formation elsewhere.

The main areas of preservation of these Zones to be discussed here are those of Dorset and Somerset - see Text fig. 1. These have not been the subject of a detailed study for over half a century, (S. Buckman, 1893, and Richardson 1931 - most of whose field work was completed before the first World War). Considering that this region of England is the type area for numerous Zonal and subzonal indices, (S. humphriesianum, Teloceras blagdeni (Sow.), T. banksi (Sow.) etc.) as well as numerous other characteristic species of ammonites, (S. Buckman, 1909-30), a revision of the stratigraphy of these beds is long overdue. Hence an attempt is made here, firstly to reinterpret the present standard Zonal scheme, secondly to give a



Text figure 1

A sketch map of part of south-west England, showing the location of the main localities mentioned in the text. The area 1. is that shown in Text fig. 2, and the area 2. is that shown in Text fig. 4. detailed bed by bed analysis of beds of this age in Southern England and lastly to determine the type horizons of the numerous ammonite species which have been described from this area.

At this point I should like to thank the curators of the various museums, who have allowed me to study type material in their care, particularly those in the British Museum (N.H.) and Institute of Geological Sciences, London; the Sedgwick Museum Cambridge and the Manchester City Museum. I should also like to thank various individuals who have allowed me access to their private collections, notably Brigadier G. Bomford, Dr. J.R. Senior and Dr. J. Whicher. Lastly I should like to give my special thanks to Dr. J. Callomon and Dr. H.S. Torrens for both help in the field and in the discussion of the numerous problems which arose during the course of this work. The majority of the work for this paper was undertaken during the tenure of a University of Keele research studentship, at that University; this award is gratefully acknowledged.

 DEVELOPMENT OF THE STANDARD ZONAL SCHEME
 The Zone of <u>Stephanoceras</u> (<u>Stephanoceras</u>) <u>humphriesianum</u> (J. de C. Sow.)

Oppel (1856) restricted his Humphriesianum Zone, in the Schwabian Albe (S. Germany), to the beds below the <u>subfurcatum</u> Oolithe, (included by him in the Parkinsoni Zone) and above the Blaukalke (= Sauzei Zone by original definition - Oppel, 1856, pp. 308, 344, etc.); that is the <u>gicanteus</u> Tone, Ostreen-Kalke and <u>blagdeni</u> Schichten of subsequent authors. Many later workers were of the opinion that S. humphriesianum was not restricted to its Zone,

	3) 3) 3)		SUBFURCATUM		HIBATIAL USI ANDM				
	SCHENE USED HE		BACULATA	SI IYI ADATOJ	ISAAKSI	INAGARNI		HUMPHALESI ANUM	INTRO
	STURANI 1971		BACULATUM	POLYGYRALIS	BANKSI	BLAGDENI	×	* STEPHANOC- CRAS*	CYCLUIDES
•	ARKELL, 1956	lanti ANA		SUBPURCATUM		INAGPAL	-	AUNT IS A LAHAMAN	
	MULLER, 1941	GAI		SUBFURCATUM		INAGSV18		HURDHRI ESI ANUM	ROMANI
	SPATH 1936			NIORTENSIS		INGODVTR			
	ARKELL, 1933			SISNALHOIN				BLAGDENT	u et di sea se di set u se ust its di se etap
	BUCKMAN, 1909- 1930		NIORTENSIS	Leptosphinctes (1923)	BAKKSI	UMBILICATA (1926)	PYGMAEUH (1927)	EPALAXITES (1922)	PARCICANINATUS (1925) MASCKEITES (1920)
	BUCKMAN, 1910 8 1913			NIORTENSIS	D ANK SI	IN: GUY TO			STENNIATOCZAAS (1913)
	BUCANAW 1891 1893 \$1898	GANAN FI ANA (1895)	CADOMENTISIS (1891) (1893) (1893)					BLAGDANI (1693)	
	OFPEL, 1856	F. A.K. INSONI		Sel The					

Table 1. The development of the standard zonal scheme for

the British Humphriesianum/Subfurcatum Zones.

and hence introduced, what they considered, were more suitable indices, such as <u>Dorsetensia romani</u> (Oppel), Haug(1891); <u>D. eduardiana</u> (D'Orb.), Brasil(1895), etc. However, reports of <u>S. humphriesianum</u> from the Sauzei Zone are due to mis-identification of such forms as <u>Skirroceras</u> and in any case an extended stratigraphic range would not preclude its use as a Zonal index.

Subdivision of the Humphriesianum Zone commenced at an early stage, with Waagen probably being the first to recognise the distinct stratigraphic distribution of 'Ammonites' (Teloceras) blagdeni Sow. (Maagen, 1867, p.539). However, the first use of Teloceras blagdeni as a stratigraphic index was as a replacement for the Humphriesianum Zone, (Six, 1879; Gosselet, 1881; Buckman, 1898). The first real attempt to subdivide the Humphriesianum Zone into separate horizons was made by Mascke, whose work on the North German Bajocian was unfortunately only published in an abbreviated 'Innaugural Dissertation', (Mascke, 1907). The four zones erected by Mascke were thus poorly defined and largely un-interpretable. The publication of Mascke's work, however, prompted Buckman to subdivide his blacdeni hemera; introduced previously as a replacement for the humphriesianum hemera (Buckman, 1898); into a restricted blacdeni hemera and an 'inter-Zone' (Buckman, 1910) or following Masche a Stemmatoceras zone, (Buckman, 1913, in 1909-30). Subsequently Buckman introduced a whole series of hemerae (stratigraphic units approximately equivalent to the subzone in present usage), of which even the relative order is doubtful, (Buckman, 1909 - 1930) - see Table 1 for further details. Spath (1936) returned to the earlier zonal concepts, when he suggested the use of a Humphriesianum Zone with Blagdeni and Romani subzones, a scheme followed by Arkell (1956), who, however, preferred

the use of a Humphriesianum subzone to that of a Romani (Aricell 1954, p.596).

In North Germany attempts have been made to give some validity to the zones introduced by Masche; thus Kumm (1952) and Westermann (1967) introduced a series of valid modifications to these zones.

Mascke 1907	Kumm 1952	Westermann 1967
<u>Teloceras</u> z one	= <u>blagdeni</u> Zone	= <u>blagdeni</u> subzone
Stepheoceras Zone	= <u>humphriesianum</u> Z one	= humphriesianum subzone
Stephanoceras zone	= umbilicum zone	= <u>umbilioum</u> subzone
Stemmatoceras Zone	= coronatum zone	= frechi subzone

Some attempts were also made in Germany to use species of <u>Dorsetensia</u> for stratigraphic indices for the basal part of the Humphriesianum Zone, with Westermann's use of a '<u>pinguis</u> Schicten' and a '<u>romani</u>/ <u>complanata</u> Schicten', (Westermann, 1954). Huf's use of the <u>pinguis</u> Zone (Dorn, 1935), however, was as a synonym of Oppel's Sauzei Zone, (Huf, 1968).

The most important recent work on the Bajocian is probably that of Pavia and Sturani (1963) in the Basse Alpes of S.E. France. Here a thick limestone/marl sequence provides one of <u>the</u> standards of reference for the world Bajocian, (Arkell, 1956, p.157). Unfortunately the base of the Humphriesianum Zone is difficult to define, since certain ammonites, particularly the genus <u>Dorsetensia</u>, are either absent, or poorly preserved. None the less a three-fold subdivision of the Humphriesianum Zone was suggested by the study of this sequence, into Blagdeni, '<u>Stephanoceras</u>' and '<u>Poecilomorphus</u>' subzones, (Pavia & Sturani, 1968, p.312). The latter subzono was subsequently given a more valid, specific rather than generic, index by Sturani (1971, p.50), who suggested the use of <u>Poecilomorphus</u> <u>cycloides</u> (d'Orb.). This three-fold subdivision of the Humphriesianum Zone appears the most satisfactory, and further subdivision into six 'horizons', (Gabilly <u>et al.</u>, 1971) would seem unnecessary.

2.2 The Zone of <u>Strenoceras</u> (<u>Strenoceras</u>) <u>subfurcatum</u> (Zieten)

Oppel recognized the distinct nature of the Subfurcatum Zone fauna (Oppel, 1856, p.308), but Terquem and Jourdy, (1869, p.2) were the first to give it Zonal status. Their Zone was later restricted by Buckman's introduction of a Garantiana hemera (Buckman, 1893) or Zone, (Buckman, 1913 in 1909-30), the basal subzone of which; the Dichotoma; must define the top of the Subfurcatum Zone. There have been numerous, subsequently designated, alternative indices for the Subfurcatum Zone, mainly related species of <u>Strenoceras</u>, such as <u>S</u>. <u>bajociensis</u>, <u>S</u>. <u>niortensis</u>, etc. but <u>S</u>. <u>subfurcatum</u> must have priority.

The first person to grasp the possibility of subdivision of the Subfurcatum Zone was undoubtedly Buckman, whose <u>banksi</u> hemera is only partly the equivalent to Mascke's <u>Teloceras</u> zone, (Buckman, 1910). Buckman later further restricted his <u>miortensis</u> hemera, by the introduction of a <u>leptosphinetes</u> hemera, (Buckman, 1923 in 1909-30). Whilst Buckman never clearly indicated the actual beds to which these hemerae referred, they can be deduced by a study of the ammonite species assigned to them in 'Type Ammonites', (Buckman, 1909-30) see Norley-Davies in Richardson (1931, p.48).

<u>niortensis</u> = <u>niortensis</u> = <u>leptosphinctes</u> (1923) - Bed 4, Frogden Quarry, Dorset. hemera, (1893), (<u>banksi</u> (1910) - Bed 5, Frogden Quarry, Dorset. Zone (1913)

Subsequent to Buchman, most work on the Subfurcatum zone has been done on the thick clay, sequences of north-west Germany. Here, mainly following the work of Althoff, (1914 and 1928) and Bentz (1924 and 1928), the following horizons have been recognised -

	Althoff 1928 Bentz 1928	Kumm 1952	Westermann 1954 and 1967
	Upper Subfurcatum Zone =	<u>schroederi</u> zone	= <u>schroederi</u> subzone
Subfurcatum Zone	Lower Subfurcatum Zone =	<u>baculata</u> zone	= <u>subfurcatum</u> subzone
	'Leptosphinctes Schicten'		= <u>phaula</u> subzone

The faunas recorded from these horizons, (Althoff, 1928; Kumm, 1952) show that they closely follow the stratigraphic succession worked out by Buckman in North Dorset. Pavia and Sturani (1968), whose work has already been discussed in connection with the Humphriesianum Zone, have largely confirmed the German results, and they recognized the following horizons in the Basse Alpes -

Subfurcatum Zone <u>baculatum</u> subzone <u>phaulus/polycyralis</u> subzone plous subzone

Subsequent modifications of this scheme have included the selection of <u>Caumontisphinctes</u> polygyralis Buckman, as the sole index for the polygyralis/phaulus horizon (Sturani, 1971) and the recognition of the priority of Buckman's banksi Zone (Buckman 1910) over Pavia and Sturani's aplous subzone, (Parsons in Sturani, 1971, p.49). French workers have only accepted a three-fold subdivision of the Subfurcatum Zone, with the omission of the schroederi subzone, (Gabilly et al.

1971). Pavia's recent work on the upper Bajocian (Pavia,1973) has largely confirmed this three-fold division, since it proved impossible to separate the <u>schroederi</u> and <u>baculatum</u> subzones in the Basse Alpes. It must be noted at this point that Pavia and Sturani's <u>baculatum</u> subzone is not the same as Kumm's <u>baculata</u> zone, (Kumm, 1952, Gabilly <u>et. al.</u>, 1971) since the index is <u>Apsorroceras baculatum</u> (Qu.) rather than <u>Garantiana baculata</u> (Qu.).

3. DISCUSSION OF THE SUB-ZONAL SCHEME

During a recent study of the Bajocian rocks of Britain, the subzonal scheme shown in Table 1 was found to give the best interpretation of the ammonite distributions. The subzones of the Humphriesianum Zone follow Pavia and Sturani's work on the Basse Alpes, but using pre-existing indices. The middle of the Humphriesianum Zone, the 'Stephanoceras subzone', (Pavia & Sturani, 1963), is characterised by the wealth of Stephanoceras s. str., and since it is the type horizon of S. humphriesianum (Sow.) in Dorset, it would seem logical to adopt this species as both the Zonal and subzonal index. This subzone would thus approximate to Kumm's restricted humphriesianum zone of N.W. Germany, which is distinguished by the occurrence of a large variety of different Stephanoceratids, including S. crassicostatum (Qu. emend. Renz), S. mutabile (Qu. emend. Mascke), S. triplex (Mascke in Weisert) and S. zieteni (Qu. emend. Renz), (Kumm, 1952, p. 388). As index for the basal subzone of the Humphriesianum Zone, I here suggest the use of Dorsetensia romani (Oppel), which was originally introduced, like Teloceras blagdeni, as a replacement index for the Humphriesianum Zone, (Haug, 1891). However, this zone has since been used in a more restricted sense, as a subzone of the Humphriesianum Zone, (Spath,

1936) and as a separate zone below a vestigal Humphriesianum Zone, (Huf, 1968). Arkell's rejection of D. romani as the index for the lower subzone of the Humphriesianum Zone (Arkell, 1954, p.596), was based on the assumption that this species occurred above S. humbriesianum, in the Blagdeni subzone. However, there is now ample evidence to show that D. romani occurs abundantly at a relatively low level in the Humphriesianum Zone; in any case an extended range up into the Blagdeni subzone would not preclude its use as an index for the lower horizon. In much of N.W. Europe the basal part of the Humphriesianum Zone is characterised, not by the fauna ascribed to the cycloides subzone, (Sturani, 1971, p.50), but by a wealth of the genus Dorsetensia. Thus in Skye, N.N. Scotland, there is a profusion of Dorsetensia and only relatively rare members of other genera, whilst the Scarborough Formation of Yorkshire, England, like North-Nest Cermany, (Westermann, 1954) and South Germany, (Stahlecher, 1935), has an abundance of Dorsetensia, often to the virtual exclusion of other genera. Taking this distribution of Dorsetensia into account, along with the priority of D. romani over Poecilomorphus cycloides (d'Orb.) as a subzonal index, there seems no alternative but to accept the former as the valid, basal index for the Humphriesianum Zone. The choice of a type horizon for the Romani subzone is clear. In his original description of the romani zone, Haug said of the 'Les couches a ammonites ferrugineuses de Beaumont', near Digne, Basse Alpes "they form one of the best types of the zone of Sonninia Romani" (Haug, 1891, p.70). The section in these beds at Truyas, Beaumont, is thus formally designated as the type locality of the Romani subzone. Since these very beds were cited in the description of the 'Poecilomorphus' subzone (Pavia & Sturani, 1968, p.312), which was

later renamed the <u>cycloides</u> subzone (Sturani, 1971, p.50), then the latter must by original definition be totally synonymous with the Romani subzone. The subzonal scheme for the Humphriesianum Zone suggested here, thus coincides with that already put forward by Muller (1941, Table 2).

The subzones of the Subfurcatum Zone adopted here, are essentially those utilised by Pavia and Sturani (1963) in the Basse Alpes, but with certain modifications. Kumm's <u>baculata</u> zone (Kumm, 1952) has priority over Pavia and Sturani's <u>baculatum</u> subzone, (Gabilly <u>et. al. 1971</u>), and is used as the highest subzone of the Subfurcatum Zone, since there is very little published evidence for a separate <u>schroederi</u> subzone, (Gabilly <u>et. al. 1971</u>; Sturani, 1971, p.49; Pavia, 1973, p.86).

4. THE HUMPHRIESIANUM AND SUBFURCATUM ZONE ROCKS OF SOUTHERN ENGLAND

The Humphriesianum/Subfurcatum zone rocks of Southern England are now preserved in five distinct areas and in two different litho-facies. The thickest deposit of rocks of this age is just to the east of Sherborne, Dorset, where they attain a maximum thickness of approximately 2.0m. The 'iron-shot' limestones from this area are relatively 'condensed' and are rich in well preserved ammonites, but the faunas have not been re-worked or mixed, and accurate collecting enables detailed stratigraphic subdivisions to be made - see Fig.3. Elsewhere the Subfurcatum and Humphriesianum Zones are only sporadically preserved in highly condensed, remanié horizons, which are rich in limonite laminae and limonitic, algal concretions, (= 'snuff-boxes', q.v. Gatrall, <u>et. al.</u> 1972). These latter horizons are useless for



Text figure 2.

A sketch map of the Sherborne-Yeovil district, showing the position of the localities cited in the text :

Bradford Abbas Railway-cutting, 2 - Halfway-House road-cutting,
 3 - Louse Hill quarry, 4- Sandford Lane quarry, 5- Clatesabe Road
 bection, 6 - Frogden quarry, 7 - Oborne Wood section, 8 - Oborne
 Lane section and 9 - Millorno Wick Lane section.

most stratigraphic purposes, but they have yielded some well preserved faunas, and they are interesting, since they give an indication of the original area of deposition, during this period.

No new formal lithostratigraphic terminology will be introduced here, as this must await a complete revision of the Dorset Inferior Oolite. Thus the existing informal bed names are used where they are applicable. The thickest deposit of Humphriesianum/Subfurcatum Zone rocks, to the east of Sherborne (beds 6-4, Oborne Wood section), have been collectively called the 'Oborne Road-stone' (Buckman, 1893. p.501; 1887-1907, p.ccvi). The upper, Subfurcatum Zone, part of these rocks (beds 6b-d, Oborne Wood), has been separated as the 'cadomensis beds' (Hudleston, 1886, p. 193; Buckman, 1891, p. 655), whilst the lowest layer of the latter beds has been called the 'Sphaeroidothyris bed' (Hudleston, 1886, Buckman, 1909-30, Pl. 543). Elsewhere the Humphriesianum/Subfurcatum Zone rocks are more condensed and thinner; thus in north Dorset and Somerset they are represented by one, thin, hard limestone bed called the 'Irony bed' (Buckman, 1893, p.487), whilst in south Dorset these Zones are represented in a thin remanie horizon here called the 'Red conglomerate' (Richardson, 1915, p.58; Gatrall, Jenkyns & Parsons, 1972, p.81).

4.1 The Sherborne Region

The area directly to the east of Sherborne, North Dorset, is a classic locale for the study of the Humphriesianum and Subfurcatum Zones; since apart from having produced the type specimens of so many Zonal and subzonal index species figured by the Sowerbys, (1812-45) and Buckman, (1909-30), it was the scene of the first real attempts



and an experiment there is that which and

to elucidate the more detailed stratigraphy of this period, (Buckman, 1893, 1910, 1909-30). The main localities described by Buckman (1893) are now poorly exposed, and many have disappeared. The following sections (see Text Fig. 2), represent all those now visible in the district, and two; Oborne Wood and Clatcombe; although very similar to nearby sections described by Buckman, were not actually seen by him. All the faunas listed in the following sections, unless otherwise stated, are based on material in the author's collections. All the bed numbering of the sections has been revised, but where applicable Buckman's numbers are given in brackets, following the new ones. The sections are described starting with those nearest Sherborne and thence eastwards to Milborne Wick. The correlation of the four main exposures in the Sherborne district is given in Text figure 3.

4.1.1 Clatcombe farm section

This is a shallow, poorly exposed, road-cutting on the west side of the road from Sherborne to the Golf course (ST636184); an old, virtually ploughed out quarry to the east of the road also shows some exposures in the condensed Humphriesianum/Subfurcatum Zones. This section must be very close to the location of that described by Buckman (1893, p.498, section XIV).

Subfurcatum Zone - Humphriesianum Zone; Blagdeni subzone

6. (7, at Lower Clatcombe, Section XIII, Euclman, 1893)

A very coarsely colitic limestone, from which the large limonite coliths tend to drop out, giving the rock a porcus, sponge like texture. Poorly exposed as a loose rubble at the top of the hedge, this bed is more extensively represented in the small quarry in the

field opposite.

Teloceras cf. blagdeni (Sow.) CP.1276

Orthogarantiana sp.

Caumontisphinctes (Infraparkinsonia) phaulus (S.B.) CP.1277

<u>C. (C.)</u> sp.

seen to 0.2m. +

Humphriesianum Zone & subzone

5c (pars 1)

A grey coloured limestone, with matt brown limonite coliths. The base of the bed is marked by a poor parting, with abundant ammonite fragments, at 0.63 m. above the top of bed 2.

> <u>Stephanoceras</u> cf. <u>humphriesianum</u> (Sow.) CP.1305 <u>S</u>. sp.

S. (Normannites) sp.

Sphaeroidothyris sphaeroidalis (Sow.)

0.13m.

0.16m.

Romani subzone

5b (pars 1)

A grey coloured limestone, with dark brown, shiny, coliths.

Chondroceras aff. evolvescens (Waag.), CP.1273

Stephanoceras of. plicatissimum (Qu. emend. Hyatt) (=

S. humphriecianum, J. Euchman, 1881, non. Sow.) CP.1310

S. (Normannites) sp.

Dorsetensia tecta S.B. CP.1276

Lissoceras colithicum (d'Orb.) CP.1293

Oppelia cf. nubradiata (Sow.) CP.1297

Poecilomorphus cycloides (d'Orb.) CP.1292

5a (2)

A grey, densely colitic limestone, with numerous dark brown coliths.

Emileia sp. (derived)

Pseudomelania sp.

0.15m.

Sauzei Zone

Planed surface, bored and limonite-encrusted

4 (3)

A light grey limestone, with yellow-brown limonite coliths, which are sparser than in bed 5a. The top of the bed contains serpulidencrusted, rolled and reworked ammonites. There is a very irregular junction with bed 3.

> Emileia (Emileia) multifida S. Buckman CP.1269 E. (E.) cf. polyschides (Waag.) CP.1270 Skirroceras cf. leptocyrale S.B. CP.1291 Pelekodites sulcata (S.B.) CP.1281 Sonninia cf. propinguans (Bayle) CP.1284 Lissoceras semicostulatum S.B. CP.1233 Acanthothiris paucispina S.B. & Walker

0.19-0.28m.

3 (4)

A light grey, soft limestone, with yellow coliths. This bed contains many badly preserved, rolled and broken ammonites, which have a characteristic soft, light grey fill to their casts. The bottom of the bed is very irregular and conglomeratic. <u>Emileia</u> (<u>E</u>.) sp. <u>E</u>. (<u>Otoites</u>) sp. <u>Bradfordia</u> sp. <u>Sonninia</u> sp. <u>Acanthothiris paucispina</u> (S.B. & Walk.) '<u>Pleurotomaria</u>' <u>granulata</u> Sow.

> 0.05-0.12n.

Laeviuscula Zone and subzone

2 (5)

A soft, light grey, marly limestone, with sparse limonite ooliths, speckled with glauconite grains, and typical of the unweathered 'green grained marl' of Oborne, (see Buckman, 1893, Frogden Qy. bed 9).

> ?<u>Witchellia</u> cf. <u>nodatipincuis</u> (S.B.) CP.1278, (= <u>Sonninia</u> sp. nov. top bed 6, Nilborne Wick, L. Richardson, 1916) <u>Mollistephanus</u> (<u>M.</u>) aff. <u>mollis</u> S.B. CP.1290

Cenoceras sp.

0.0-0.1m.

1 (6)

A 'blue hearted', crystalline, slightly sandy and glauconitic limestone, (= 'Blue bed', bed 10 Frogden) which forms a 'kerb' on the west side of the road.

> ?<u>Witchellia</u> of. <u>nodatipinguis</u> (S.B.) CP.1279 seen to 0.15m.

4.1.2 Frogden Quarry Oborne See And Meridian (559)
The section described by Eucliman from this quarry (ST642185) is
ctill visible from his bed 6 up (Buckman, 1893, p.500, section XV).

This quarry was the source of numerous specimens figured by Buckman (1883, 1909-30) and Hudleston (1887-96) and has been described by Hudleston (1886), Richardson (1932) and more recently by Macfadyen (1970, p.158). The bed numbers used here are the same as those for the Oborne Wood section.

7 (pars 2)

A sandy, marly limestone, heavily stained with limonite and with a conglomerate of limonite-stained, re-worked ammonites at its base. 0.30m.

- - - planed surface -

'Oborne Road-stone' (6-5), - 'cadomensis Beds' (6)

Subfurcatum Zone, Baculata subzone

6d (pars 3)

A highly fossiliferous, hard colitic limestone. There are many fragmentary fossils, which towards the top of the bed, are often limonite coated.

> Caumontisphinetes (Infraparkinsonia) cf. bonarellii (Parona), CP.3099 Leptosphinetes (Leptosphinetes) cf. leptus S. Buckman, CP.2802 L. (L.) cf. davidsoni (S. Buckman), CP.2803 L. (Cleistosphinetes) cf. asinus (Zatvornitzki), CP.2796 Strenoceras (S.) subfurcatum (Zieten), CP.2795 Strenoceras (Garantiana) baculata (Qu.), CP.2793 S. (G.) spp. Apsorroceras baculatum (Qu.), CP.2792 Lissoceras (Lissoceras) colithicum (d'Orb.), CP.2797 L. (L.) psilodiscus (Schloenbach) s. sp. inflatum Wetzel, CP.2793

L. (<u>Microlissoceras</u>) cf. <u>pusillun</u> Sturani, CP.2799 <u>Strigoceras</u> (<u>Strigoceras</u>) sp.

S. (Cadomoceras) sullyence Brasil, CP.2801

S. (C.) nevos (Parona), CP.2800

Chondroceras canovense (de Gregorio), CP.2331

Sphaeroceras auritum cf. subsp. auritum Parona, CP.2332

0.15-0.20m.

Polygyralis subzone

6c. (pars.3)

A hard, blue 'hearted', oolitic limestone, highly bioturbated, with a mixture of more sandy limestone. This bed is poorly fossiliferous, with the few fossils being badly preserved and difficult to extract. There is a very large species of <u>Pholadomya</u> present, which was probably the source of the bioturbation.

Caumontisphinetes (Caumontisphinetes) cf. polycyralis S. Buckman, CP.2806

Leptosphinctes (Cleistosphinctes) sp.

<u>Oppelia (Oecotraustes</u>) cf. <u>longarae</u> Sturani, CP.2805 0.35-0.40m

'Sphaeroidothyris Bed'

6b. (4)

Lenticular masses of hard, blue 'hearted', crystalline, 'ironshot'limestone, which are very irregular in thickness, and are sometimes only with some difficulty separated from the bed above. This bed is very sandy in patches, and 'nests' of <u>Sphaeroidothyris</u> are common. <u>Caumontisphinetes</u> (<u>Infraparkinsonia</u>) <u>phaulus</u> (S.Buckman), <u>CP.2808</u> <u>Strenoceras</u> (<u>Strenoceras</u>) sp. <u>Orthogarantiana</u> (<u>Orthogarantiana</u>) cf. <u>densicostata</u> (Qu. emend. Douv.), CP.2809 <u>O. (O.) haugi</u> Pavia, CP.2807 <u>Torrensia gibba</u> (Parona), CP.2804 <u>Oppelia (Oppelia</u>) sp. <u>Strigoceras (Cadomoceras</u>) sp. <u>Cadomites (Cadomites</u>) sp. <u>Chondroceras canovense</u> (de Gregorio), CP.2345

> 0.10-0.20m.

Banksi subzone

6a. (pars ? 5)

A soft brown, colitic limestone only sporadically preserved, where the large specimens of <u>Teloceras</u>, resting on the basal erosion plane, 'stick up' into the overlying bed.

Teloceras banksi (Sow.), CP.2647

0.0-0.18m.

<u>Total</u> for bed 6 = 0.66 - 0.71 m.

planed surface -

56. (5)

A thin, soft, marly limestone; densely colitic, with shiny, brown limonite coliths set in a yellow-brown, often purple stained, matrix. Belemnites are particularly common. Divided by two poor partings, into three courses. Caumontisphinctes (Caumontisphinctes) cf. aplous S. Buokman, CP.2811

C. (Infraparkinsonia) sp.

Leptosphinctes (Leptosphinctes) sp.

L. (Cleistosphinctes) sp.

?Cadomites cf. humphriesiformis Roche, CP.3071

Chondroceras cf. tenue (West), CP.2278

Teloceras banksi (Sow.), CP.2810

T. aff. blagdeni (Sow.), CP.2988

T. sp.

0.31 -0.37m.

Humphriesianum Zone, Blagdeni subzone

5a. (6)

A hard, limestone with shiny orange coliths set in a grey-brown matrix. There are patches of more sandy limestone, due to bioturbation, and numerous belemnites present.

Teloceras blagdeni (Sow.)

seen to 0.30m.

seen to 1.Om.

Laeviuscula Zone, Ovalis subzone

1b. (13)

A series of thin bedded, grey, glauconitic limestones, with soft, glauconitic marl partings.

Trilobiticeras (Emileites) aff. liebi (Maub.), CP.1503

Explanation of Plate 1.

Figure 1.

A general view of the temporary section at Oborne Wood, near Sherborne, showing the 'Oborne Road-stone', <u>cadomensis</u> beds and the basal part of the Sherborne Building-stone. The bed marked 'C', with the flat top is the massive block of the cadomensis beds.

e se la construir e la serie de la construir d En la construir de la construir

Figure 2.

A close-up of the Humphriesianum Zone beds at Oborne Wood.



Figure 1.



Discites Zone

la. (14)

A soft, grey, slightly conglomeratic, glauconitic limestone, with many fossils, particularly ammonites, which are preserved as distorted internal moulds.

Docidoceras (Docidoceras) cylindroides S. Buckman, CP.1637 Trilobiticeras (Trilobiticeras) platygaster S. Buckman, CP.1723

<u>T.</u> (<u>T.</u>) cf. <u>punctum</u> (Vacek), CP.1724

Euhoploceras of. polycanthum (Naagen), CP.1710

Hyperlioceras (Hyperlioceras) walkeri (S.Buckman), CP.1630 Reynesella sp.

0.30m.

4.1.3 Oborne Wood

This is a large temporary section (ST64S18S), very near to, and showing an exactly similar succession to that of Frogden Quarry (see previous section). A series of exposures have been available for study at various times (see Plate 1, figs. 1 and 2), and at different points the beds exposed showed a considerable variation in thickness; the section recorded here is thus to a certain extent a composite one. These exposures have already been the subject of a general description (Whicher & Palmer 1971).

Garantiana Zone? + ?Subfurcatum Zone (see Section 4)

'The Sherborne Building Stone Series'

9 (1)

A massive, grey, crystalline, sandy and slightly colitic limestone which weathers to a yellow-brown colour.

seen to 1.2m.

8 (pars 2)

A series of five (a-e) hard, yellow-grey limestone bands very similar to the bed above, alternating with, grey, sandy marls. These beds are relatively unfossiliferous, bed 8b having yielded -

<u>Cadomites</u> (<u>Cadomites</u>) cf. <u>deslongchampsi</u> (d'Orb.) CP.2812 <u>Acanthothiris</u> sp. - common

Total l.lm.

7 (pars 2)

Two thin (0.08m.) yellow-brown colitic limestone bands. The upper bed (b), has an iron-stained upper surface, whilst the lower bed is slightly softer. These beds are separated by brown sandy marks of similar thickness.

0.3m.

A planed surface, with a conglomerate of derived, iron-coated fossils, cemented to the bed below by a limonite crust.

'The Oborne Road-stone' (6-4)

Subfurcatum Zone

'The cadomensis beds' (6)

This is a solid block of 'iron-shot' limestone of variable lithology, being more colitic and fossiliferous in the top 0.15m., whilst elsewhere, due to bio-turbation, there are patches of sandy material mixed with more crystalline limestone. Bounded at top and bottom by two prominent erosion planes, this bed is divisible into four horizons (6a-d), by three irregular partings.

Baculata subzone

6d. (pars 3)

A highly fossiliferous, brown colitic limestone. The fossils

are often fragmentary, particularly at the top of the bed, where they are also abraded and limonite stained.

<u>Strenoceras</u> (S.) <u>subfurcatum</u> (Zieten), CP.3103 <u>Strenoceras</u> (<u>Garantiana</u>) <u>baculata</u> (Qu.), CP.3100 <u>S.</u> (<u>G.</u>) spp. <u>Orthogarantiana</u> cf. <u>inflata</u> (Bentz), CP.3105 <u>Leptosphinotes</u> (<u>Leptosphinotes</u>) cf. <u>leptus</u> S.B., CP.3082 <u>L.</u> (<u>L.</u>) <u>rotula</u> (Parona), CP.3101

L. (Cleistosphinctes) asinus (Zat.), CP.3102

Chondroceras canovense (de Greg.), CP.2321

Sphaeroceras auritum (Parona) cf. subsp. auritum, CP.2724

Apsorroceras baculatum (Qu.), CP.2695

Spiroceras spp.

Lissoceras (L.) colithicum (d'Orb.)

Oppelia (Oppelia) flexa (S. Buckman), CP.3083

Strigoceras (Strigoceras) sp.

S. (Cadomoceras) sullyense Brasil, CP.3104

0.15-0.18m.

Polygyralis subzone

60. (pars 3)

A bio-turbated mixture of blue colitic and sandy limestone, Much of this bed is unfossiliferous, the fossils being concentrated at the base, just above the parting which divides the whole of bed 6 in half.

Leptosphinetes (L.) cf. davidsoni, CP.2701

L. (L.) leptus, CP.2702

L. (Cleistosphinctes) asinus, CP.2700

<u>Strenoceras</u> (<u>Strenoceras</u>) cf. <u>subfurcatum</u> <u>S. (S.) apleurum</u> S. Buchman, CP.3106 <u>Caumontisphinotes</u> (C.) aff. <u>polygyralis</u> S. Buchman <u>C. (Infraparkinsonia) phaula</u> (S. Buchman), CP.3107 <u>Cadomites</u> (C.) cf. <u>deslongchampsi</u>, CP.3031 <u>C. (Polyplectites</u>) sp. nov. <u>Chondroceras canovense</u> <u>Sphaeroceras aff.auritum</u> CP.2336 <u>Lissoceras colithicum</u>, CP.3109 <u>Oppelia</u> (O.) <u>flexa</u> (S. Buchman), CP.3108 <u>O. (Oecotraustes) pulchra</u> (S. Buchman), CP.3110

0.18m.

'The Sphaeroidothyris Bed'

6b. (4)

A lenticular mass of blue 'hearted', crystalline, 'iron-shot' limestone, with a large amount of detrital quarts. This bed is very irregular in thickness, it fills the umbilical areas of the <u>Teloceras</u> in the bed below, and where these are absent, it extends down to the basal erosion plane. <u>Sphaeroidothyris</u> <u>sphaeroidalis</u> (Sow.) is very common, both as individuals, and in 'nests'.

Caumontisphinctes (C.) polycyralis, CP.2693

C. (Infraparkingonia) phaula, CP.2699

Leptosphinetes (L.) davidsoni, CF.2690

L. (L.) lentus, CP.2683

L. (<u>Cleistosphinotes</u>) asinus

Strenoceras (S.) cf. subfurcatus

Strenoceras (Garantiana) sp.

Orthogarantiana cf. densicostata (Qu.) O. haugi Pavia, CP. 3079 Cadomites (C.) deslongchampsi, CP.2969 C. (Polyplectites) sp. nov. cf. P. linguiferus (d'Orb.) Chondroceras canovense, CP.2335 0. (Oecotraustes) cf. pulchra

> 0.08-0.25%.

Banksi subzone

6a. (pars ?5)

A soft, brown, colitic limestone, very similar to the bed below. but which is often absent, especially when the common Teloceras banksi (Sow.), which lay flat on the basal erosion surface, are also missing.

> 0.0-0.18m. Total for bed 6 = 0.5-0.56m.

A planed surface 0.90 - 1.0m. above bed 3, surmounted by 0.05n. of marl

5b. (5)

A soft, brown, 'earthy', colitic linestone, divided by two irregular partings into three courses. There are numerous small corale, Discacyathue, and Belennites present. The larger

C. (C.) aff. homalogaster S. Buckman, CP.2693

Torrensia gibba (Parona), CP.2804

Oppelia (0.) aff. flexa

Lissoceras oolithicum

Strigoceras (Cadomoceras) sullyense
ammonites lay parallel to the bedding, with their upper surfaces planed off.

<u>Caumontisphinctes</u> (<u>C</u>.) <u>aplous</u> S. Buckman, CP.2981 <u>C</u>. (<u>C</u>.) <u>diniensis</u> (Pavia), CP.2982 <u>C</u>. (<u>Infraparkinsonia</u>) cf. <u>phaula</u>, CP.2983 <u>Leptosphinctes</u> (<u>L</u>.) sp. nov. <u>L</u>. (<u>Cleistosphinctes</u>) sp. nov. <u>Stephanoceras</u> (<u>Normannites</u>) sp. <u>Teloceras</u> (<u>T</u>.) <u>banksi</u> (Sow.) <u>T</u>. (<u>T</u>.) cf. <u>blagdeni</u> (Sow.) <u>T</u>. (<u>T</u>.) <u>lotharingicun</u> Maubeuge <u>T</u>. (<u>Epalaxites</u>) sp.

Humphricsianum Zone, Blagdeni subzone 5a. (6)

> A hard, grey-brown, colitic linestone, with a brittle fracture. This bed is highly bio-turbated; with the introduction of softer, sandier material. The common, large ammonites tend to be 'rotten', - they have a soft limonite rich fill to their inner whorls, - and, unlike the bed above, they occur at all angles to the bedding.

- - Marl parting - - -

<u>Teloceras</u> (<u>T</u>.) <u>blardeni</u>, CP.2646 <u>T. (Epalaxites</u>) sp. <u>Stephanoceras</u> (Normannites) sp.

> 0.2-0.25m.

0.08-0.3m. - - - Marl parting 0.01 - 0.1m. thick -

A hard, grey, colitic limestone, divisible by fauna, and to a lesser extent by lithology into three horizons, (4a-c).

Humphriesianum subzone

4c. (pars 7)

A hard, grey limestone with matt brown coliths and soft sandy patches, particularly at the top. The bottom of the bed is marked by a poor parting, associated with a layer of large, serpulid-encrusted, flat lying ammonites, indicating a pronounced pause in sedimentation.

Teloceras (T.) acuticostatum Weisert, CP.2714 T. (T.) cf. lotharingicum, CP.2686 T. (Epalaxites) sp. Stephanoceras (S.) gibbosum (S. Buckman), CP.2705 S. (S.) crassicostatum (Qu. emend. Renz) S. (S.) humphriesianum (Sow.), CP.2719 S. (S.) pyritosum (Qu. emend. Renz), CP.2972 S. (S.), cf. scalare (Masche emend. Weisert), CP.2712 S. (S.) mutabile (Qu. emend. Renz), CP.2716 S. (S.) zieteni (Qu. emend. Renz), CP.2718 S. (Skirroceras) sp. S. (Normannites) formosum (S. Buckman), CP.2977 S. (N.) latansatus (S. Buckman), CP.2979 Chondroceras evolvescens (Waag.) C. gracile (Westermann), CP.2273 Poecilomorphus (P.) cycloides (d'Orb.), CP.2457 Dorsetensia regrediens Haug, CP.2720

92

0.15

Romani subzone

4b. (<u>mars</u> 7)

A hard, grey collic limestone. This horizon is extremely fossiliferous and it is characterised by the very shiny collichs, by the soft, light grey coloured limestone fill to the fossils casts, and by the abundance of ammonite fragments.

Chondroceras delphinus (S. Buckman), CP.2266

C. evolvescens, CP.2225

C. gervillii (Sow.), CP.2267

Sphaeroceras brongniarti, CP.2325

Phaulostephanus paululum S. Buckman, CP.2978

Stenhanoceras (S.) nodosum Hyatt, CP.1532

<u>S. (Normannites) bicostatum</u> (Westermann, 1954, Pl.30, fig.3 <u>non</u>. 1)

S. (N.) latansatum, CP.2980

S. (N.) of. orbitryi (S. Buchman)

Teloceras (T.) blagdeniformis (Roche), CP.2965

T. (Epalarites) cf. laticostatus (Westermann), CP. 3070

T. (E.) portitor (Maubeuge), CP.2966

Dorsetencia alsa tica (Buchman, 1909-30, pl.528, non Haug.), CP.2116

D. complanata S. Buckman, CP.2496

D. pulchra S. Buckman, CP.2497

D. regrediens Haug, CP.2498

D. subtecta S. Buckman, CP.2502

Onnelia (0.) subradiata (Sow.), CP.2477

0. (Cecotraustes) renicularis, CP.2488

Poecilomorphus (P.) cyclaides, CP.2455

P. (Micropoecilomorphus) vicetinus (Parona), CP.2473 <u>Stegoxyites parcicarinatus</u> S. Buchman, CP.2493 <u>Strigoceras</u> (S.) <u>bessinum</u> Brasil, CP.2453 <u>S. (Cadomoceras</u>) sp. <u>Nannolytoceras pygnaeum</u> (d'Orb.), CP.2399

> 0.15-0.20m.

4a. (8)

A dark brown, densely colitic limestone with a basal conglomerate consisting of derived lumps of, 'green grained marl' and soft grey limestone with yellow-brown coliths. As well as re-worked pebbles there are also limonite coated fossils and rare 'Snuffboxes', analagous with those of south Dorset, (Gatrall, Jenkyns & Parsons, 1972, p.81).

in situ Dorsetensia subtecta, CP.1357

<u>Chondroceras delphinus</u>, CP.1351 <u>Sphaeroceras brongniarti</u>, CP.1352

derived

Emileia (E.) sp. Frogdenites spiniger S. Buckman, CP.1189 'Skirroceras' cf. skolex (S. Buckman), CP.1192 Sonninia sp. <u>Witchellia (W.) sp.</u> <u>N. (Pelekodites) sp.</u>

> 0.1-0.15m.

Laeviuscula Zone and subzone

3 (9)

A soft, light grey limestone, speckled with green glauconite

grains and with patches of yellow-brown ooliths. On weathering this bed has the appearance of a soft white marl. There is no sharp boundary with the bed below, only a gradual transition. This bed is full of fossils, many of which are oyster-and serpulid-encrusted and, at the top of the bed, eroded and iron stained.

Emileia (E.) brocchii (Sow.), CP.1353

E. (E.) polyschides (Waag.), CP.1359

E. (Otoites) contracta (S.B. non. Sow.), CP.1334

Frogdenites spiniger S. Buckman, CP.1210

F. cf. profectus S. Buckman

Chondroceras sp. nov.

Mollistephanus aff. mollis S. Buckman

Skirroceras aff. kalus (S. Buckman), CP.1332

S. leptogyrale S. Buckman, CP.1333

Bradfordia inclusa S. Buckman, CP.1371

Oppelia (0.) amblys (S. Buckman), CP.2996

Papilliceras arenatus (Qu. emend. S.B.), CP.1373

Shirbuirnia superba (S. Buckman), CP.1374

S. cf. trigonata (Qu. emend. Dorn), CP.1375

Mitchellia (W.) actinophora S. Buckman

W. (W.) falcata S. Buchman

W. (W.) glauca S. Buckman, CP.1394

<u>N. (N.) plena</u> (S. Buckman), CP.1390

N. (Pelekodites) aurifer (S. Buckman)

M. (P.) macra (S. Buckman), CP.1387



A hard, blue 'hearted', glauconitic limestone. The upper surface of the bed is the most fossiliferous and it is also softer and more glauconitic. Elsewhere fossils are sparse and poorly preserved. This bed has been extensively bored, the narrow vertical borings allowing the marl bed to be 'piped down'.

Emileia (E.) crater S. Buckman, CP.1638

E. (E.) aff. catamorpha S. Buckman, CP.1641

E. (E.) polyschides, CP.1640

E. (Otoites) sp.

Papilliceras arenatus, CP.2406

Sonninia sp.

Witchellia (W.) laeviuscula (Sow.)

W. (W.) rubra (S. Buckman), CP.2407

W. (Pelekodites) sp.

Lissoceras semicostulatum S. Buckman, CP.2408

0.23m.

Ovalis subzone

lc.

A more massively bedded top to the subjacent thinly bedded glauconitic limestones. This bed is only occasionally present.

Sonninia ovalis (Qu. emend. S.B.)

0.0-0.45m.

16. (13)

A series of thin, yellow-grey, sandy, glauconitic limestones, interbedded with soft, brown, sandy marls. These beds are not very fossiliferous and have not been well exposed. <u>Witchellia (W.)</u> sp.

Sonninia sp.

Emileia (E.) sp. nov. aff. E. catamorpha S. Buckman, CP.1624 circa 4.0m.

Discites Zone

la. (14)

A grey, glauconitic limestone, slightly conglomeratic and with many fossils, particularly distorted internal moulds of ammonites.

<u>Sonninia</u> (<u>Euhoploceras</u>) cf. <u>polycantha</u> (Waag.), CP.1616 <u>Hyperlioceras</u> cf. <u>liodiscites</u>, CP.1619 Reynesella sp.

Graphoceras apertum S. Buckman, CP.1621

seen to 0.30m.

4.1.4 Cborne Lane Section

A very poor section is still visible in the floor of this lane (ST.656186). The old cutting is too slipped and overgrown to make out anything of Buchman's section (Buchman, 1893, section XVI).

4 (4)

A soft, brown, iron-shot linestone.

scen

Laeviuscula Zone and subzone

3 (5)

A coft, white clauconitic marl.

<u>Witchellia</u> (<u>Witchellia</u>) <u>laeviuscula</u> (Sow.), CP.3098 <u>Papilliceras</u> aff. <u>arenatus</u> (Qu. emend. S.E.), CP.3097 0.10m. A massive, hard, glauconitic limestone.

0.25%.

Hard, thin bedded, glauconitic limestones, with marl partings. seen

(Further up the lane a single specimen of <u>Teloceras</u> sp. was found <u>in situ</u>, in the basal part of the Sherborne Building Stones. This shows that, as at Milborne Port, (Kellaway & Wilson, 1941, p.154), the Subfurcatum Zone beds, become more 'expanded' and less iron-shot, towards the east, and thus become indistinguishable from the overlying Building Stones.) See p. 146 below.

4.1.5 Milborne Wick Lane Section

Although now poorly exposed and overgrown, this road-cutting (ST663205), still shows a similar section to that of Buckman's, (Buckman, 1893, section XVII). This section has always been famous for the abundant and superbly preserved Romani subzone fauna, which may be collected from bed 5. Material from this bed has found its way into virtually every museum in the United Kingdom, as well as many abroad.

6 (1)

A hard, crystalline, light yellow coloured, sandy limestone. The basal few centimetres contain pebbles derived from bed 5, as well as eroded, limonite coated fossils.

- Planed curface

seen to 0.90m.

2 (6)

1

Humphriesianum Zone, Romani subzone

5 (2)

A soft, white, marly limestone with numerous yellow limonite ooliths and black dendritic manganese staining. This bed is highly fossiliferous, with many of the shells having been replaced by a pink coloured calcite.

Chondroceras evolvescens (Waagen), CP.2178

C. gervillii (Sow.), CP.2214

Phaulostephanus cf. paululum S. Buckman, CP.2616

Stephanoceras (Stephanoceras) cf. scalare (Mascke, emend. Weisert), CP.2601

 \underline{S} . (\underline{S} .) spp. (many fragments)

S. (Normannites) crassicostatum (Westermann), CP.2606

S. (N.) mitis (West.), CP.2610

S. (N.) cf. portitor (Maubeuge), CP.2604

S. (?Germanites) bicostatus Nest., CP.2605

Teloceras blagdeniformis (Roche), CP.2555

Dorsetensia deltafalcata (Qu.), CP.2550

D. equardiana (d'Orb.), CP.2556

D. liostraca S. Buckman, CP.2554

D. regrediens Haug, CP.2551

Lissoceras (Lissoceras) colithicum (d'Orb.), CP.2575

Oppelia (Oppelia) subradiata (Sow.), CP2562

O. (O.) aff. skrodzkii Brasil, CP.2574

0. (Occotraustes) genicularis Waagen, CF.2576

Poecilomorphus (Poecilomorphus) cycloides (d'Orb.), CP.2558

P. (Micronoecilomorphus) vicetinus (Parona), CP.2599

Stegozyites (Stegozyites) aff. percicarinatus S. Buckman, CP.2563 0.07-

0.10m.

4 (3)

A hard limestone, with yellow-brown ooliths set in a light bluegrey matrix. In contrast to the bed above, there is very little manganese staining, and fossils are sparser and more difficult to extract. There is a layer of bivalves towards the centre of the bed , and a conglomerate of derived lumps of bed 3 at the base.

<u>Stephanoceras (Normannites) formosum</u> (S. Buchnan), CP.2679 <u>S. (N.) portitor</u> (Naub.), CP.2644 <u>Dorsetensia liostraca</u> S. Buchman, CP.2678 <u>Oppelia (Oppelia) subradiata</u> (Sow.), CP.2638 <u>Stegozyites (Stegozyites) parcicarinatus</u>, CP.2634 0.20-0.25m.

?Sauzei Zone

3 (4)

A soft, white marl, speckled with green glauconite grains. There are no limonite coliths present.

- Unconformity

<u>Kumatostephanus</u> (<u>Kumatostephanus</u>) cf. perjucundus 5. Buckman, CP.1563

K. (?Gerzenites) rugosus Westermann, CP.1530

?Labyrinthoceras sp. nov.

Skirroceras bayleamun (Oppel), CP.1531

S. skolex (S. Buchman), CP.1564

Emileia (Emileia) polyschides (Waag.), CP.1565

E. (Otoiter) contractus (S.B. non. Sow.), CP.1574

Papilliceras arenatus (Qu. emend. S.B.), CF.1551

0.10-0.15m. Laeviuscula Zone and subzone

2b. (pars 5)

A hard, blue-green limestone, highly glauconitic and forming a softer, irregularly preserved, top to the sub-adjacent bed.

K. (?Gerzenites) cf. rugosus, CP.1576

Mollistephanus (Mollistephanus) sp.

Emileia (Emileia) sp.

E. (Otoites) sp.

Witchellia (Witchellia) laeviuscula (Sow.)

N. (Pelekodites) aff. macra (S. Buckman), CP.1559

0.0-0.10m.

2a. (pars 5)

A hard, blue limestone, in two-three layers, and more sparsely glauconitic than the bed above.

Emileia (Emileia) cf. brocchii (Sow.), CP.1543

Shirbuirnia (Stiphromorphites) cf. <u>nodatipinguis</u> S. Buckman. 0.45-0.60m.

1 (6)

A series of hard, sandy, sparsely glauconitic limestones,

intercalated with soft, sondy marls.

<u>Witchellia</u> (<u>Witchellia</u>) aff. <u>zugophorus</u> (S. Buckman) CP.1545 seen to 3.0m.

Note

There has been some confusion in the past literature concerning this section. Richardson (1916, p.516) considered that Euchman (1893) had been overzealous in splitting up of some of his beds (5-3, here), since Richardson considered that all three were of the same age and lithology. However in this case there can be no doubt that Buckman was correct, since bed 3 is the only one of this group of marly limestones to contain glauconite grains and it is also older, yielding as it does Sauzei rather than Humphriesianum Zone ammonites.

4.2 The North-west Dorset - South Somerset Area

The Humphriesianum/Subfurcatum beds thin rapidly to the west of Sherborne and at Sandford Lane (ST628179), they have disappeared completely. West from this point, these Zones are represented in the highly condensed deposit known as the 'Irony bed', (S. Buckman, 1893, p.485). This bed, which consists of a thin (-0.18m.), hard, crystalline limestone, containing numerous large, limonite ooliths, comminuted crinoid debris and many small 'snuff-boxes' (limonite rich, algal concretions), is consistently heavily stained with limonite; hence its name. The Irony bed is always non-sequentially related to adjacent beds and it is found as far west as Seavington-St.-Mary, the most westerly outcrop of the Inferior Dolite, and as far south as Bradford Abbas, beyond which the Inferior Colite is mainly faulted out (see Fig.2). Although this horizon has yielded amonites representative of all the Lover and basal Upper Bajocian Zones (Gatrall, Jenhyns & Parsons, 1972, p.83), several localities have produced exclusively Humphriesianum and Subfurcatum Zone faunas.

4.2.1 Halfway House - Louse Hill.

There were once several old quarries in this area, showing exposures of the 'Irony-Bed' normally resting on Discites Zone rocks. There are now only two exposures; Buckman's old locality - Louse Hill, (ST608152), (Buckman, 1893, p.488), where this bed has yielded

a Romani subzone fauna - <u>Chondroceras cervillii</u> (Sow.), <u>Sphaeroceraa</u> <u>brongniarti</u> (Sow.), <u>Stephanoceras</u> (<u>Normannites</u>) <u>variecostatus</u> (Westermann, 1954, Pl.25, fig. 5 <u>non</u> 3), <u>Oppelia</u> (<u>Oppelia</u>) sp., <u>O</u>. (<u>Oecotraustes</u>) <u>genicularis</u> (Waag.) and <u>Poecilomorphus cycloides</u> (d'Orb.); and a new road cutting at Halfway House, (ST602163), where the 'Irony Bed' has produced a Banksi subzone fauna of, <u>Caumontisphinctes</u> (<u>Infraparkinsonia</u>) cf. <u>phaulus</u> (S.B.), <u>Leptosphinctes</u> (<u>Cleistosphinctes</u>) sp. and <u>L.</u> (<u>Cleistosphinctes</u>)/<u>C</u>. (<u>Infraparkinsonia</u>), (intermediate sp.).

4.2.2 Bradford Abbas

The 'Irony Bed' now exposed in the Bradford Abbas railway cutting, as well as that once exposed in the old quarries, is mainly Sauzei Zone in age. However a recent Gas Pipe trench in this area has yielded numerous Subfurcatum Zone ammonites from the top of the 'Irony Bed', mainly <u>Strenoceras</u> (<u>Strenoceras</u>) <u>subfurcatum</u> (Zieten), (<u>teste</u> H.S. Torrens).

4.2.3 Seavington-St.-Mary

The outlier at Seavington-St.-Mary is the most westerly outcrop of the Inferior Oolite - see Fig.1. This section (ST396144), has recently been described in detail (Parsons in Torrens 1969, p.A26), but further collecting has revealed the presence of a Romani subzone fauna in bed 7, directly below the stromatolite layer, including, <u>Stephanoceras (Stephanoceras) sp., Oppelia (Oppelia) sp., Poecilo-</u> <u>morphus cycloides (d'Orb.) and Strigoceras (Strigoceras) bessinum</u> (Sturani non Brasil).

4.3 The South Dorset Area

In South Dorset the Humphriesianum Zone is represented in the



Text Figure 4.

A sketch map of the Burton Bradstock - Beaminster district, showing the position of localities cited in the text : 1 - Burton Cliff section, 2- Bonscombe Hill section, 3 - Stony Head Cutting, 4 - Upton Manor Farm quarry, 5 - Loders Pipe trench, 6 - Welcome Hill quarry, 7 - North Poerton, 8 - North Marren Hill and 9 - Hern Park quarry. 'Red Conglomerate'. This consists of re-worked and remanié fossils set in a limonite rich red clay or marly limestone replete with crinoid ossicles and pebbles, and which is cemented to the top of the 'Red Bed', (Richardson, 1928-30, p.48). There is some evidence at Burton Bradstock that this bed was subsequently further re-worked during the Subfurcatum Zone. Whilst the 'Red Conglomerate' is not present at every South Dorset Inferior Oolite locality, its position is always marked by a limonite stained and encrusted surface, - see Text fig. 4.

4.3.1 Burton Bradstock

The 'Red Conglomerate' is well exposed at the Cliff Hill section, Burton Bradstock, (SY487891), but it is easier to collect from the fallen blocks at the foot of Burton Cliff (SY482892). Both of these localities were described by Richardson (1928-30), the second having produced the following ammonites; <u>Stephanoceras (Stephanoceras) kreter</u> (S.B.), <u>S</u>. cf. <u>humphriesianum</u> (Sow.), <u>S</u>. aff. <u>nodosum</u> Hyatt, <u>S</u>. <u>trichalus</u> (Westermann), <u>S</u>. (Normannites) of. <u>orbignyi</u> (S.B.), <u>Teleceras</u> sp., <u>Oppelia</u> (<u>Oppelia</u>) <u>flexa</u> (S.B.), <u>O</u>. (<u>O</u>.) <u>lectotypa</u> S.B. and <u>Poecilomorphus cycloides</u>. At the latter locality a more white coloured matrix has penetrated down into the 'Red Conglomerate' and this has yielded Subfurcatum Zone ammonites; <u>Strenoceras</u> (<u>Strenoceras</u> sp., <u>Lentosphinotes</u> (<u>L</u>.) aff. <u>davidsoni</u> (S.B.) and <u>Spiroceras</u> sp., (Gatrall <u>et al</u>. 1972, p.82).

4.3.2 Bonscombe Hill - Hyde Quarry

A recent temporary section at Bonscombe Eill (SY488919), produced Humphriesianum Zone ammonites from bed 29 (Senior, Parsons & Torrens, 1970, p.116); <u>Stephanoceras humphriesianum</u> (Sow.) and Sphaeroceras sp. The nearby Hyde Quarry (SY434922) has also produced a specimen of S. humphriesianum (Woodward, 1394).

4.3.3 Loders Cross

There are several localities in this area which have produced Humphriesianum Zone ammonites. A new road cutting at Stony Head (SY496927) yielded <u>Sphaeroceras brongniarti</u> (Sow.) and <u>Stephanoceras</u> of. <u>plicatissimum</u> (Qu. emend. Hyatt) from the 'Red Conglomerate', whilst the old Stony Head quarry to the North (SY497929) has produced a specimen of <u>S. humphriesianum</u> (Sow.) - I.G.S. 56552, from the same horizon. The small exposure to the north-east of Loders Gross (SY506929), described by Bomford (1948) has also produced Humphriesianum Zone ammonites, mainly uninterpretable stephanoceratid nuclei, and one specimen of <u>Chondroceras</u> of. <u>gracile</u> (Westermann).

4.3.4 Upton Manor Farm

The section at Upton Manor Farm (SY512936) was described by Richardson (1928-30, p.164), since which time a Humphriesianum Zone fauna has been found at the top of the 'Red Bed', (Senior <u>et al.</u> 1970, p.117, bed 6). This fauna is the most extensive from this Zone found in South Dorset, and includes - <u>Sphaeroceras brongniarti</u>, <u>Stephanoceras</u> (<u>S.</u>) <u>kreter</u> (S.B.), <u>S.</u> cf. <u>humphriesianum</u>, <u>S. plicatissianum</u>, <u>S.</u> (<u>Phaulostephanus</u>) <u>paululum</u> (S.B.), <u>S.</u> (<u>Normanuites</u>) cf. <u>orbignyi</u> and <u>Oppelia</u> (<u>0.</u>) <u>flexa</u>.

4.3.5 Other 'Red Conglomerate' localities

Most exposures at the top of the 'Red Bed' show signs of the presence of remanié style deposits, (Richardson, 1928-30, p.45). Limonite rich, red clay yielding small, un identifiable ammonite nuclei has been found at : Loders Pipe trench, (SY503945 - Parsons, 1972); North Poorton, (SY520986); North Warren, (SY486998); Welcome Hill, (SY503953) and Horn Park quarry, (ST458022).

4.4 The 'Cole Syncline'

In the area of Bruton (Somerset), a small syncline, possibly a remnant of an original basin of deposition, has preserved an isolated patch of Aalenian and Lower Bajocian rocks some way to the north of the main Sherborne outcrops, (Richardson, 1916) - see Fig. 1. At the Lusty quarry, Bruton, (ST679344), a thin, hard limestone bed, very like the 'Irony Bed' to the south, has yielded a solitary specimen of <u>Teloceras</u>, (Richardson, 1916, p.496, bed 4). This specimen, now in Reading University collections, (3134), undoubtedly shows the Blagdeni subzone age of this bed. Unfortunately it has not been possible to find any further ammonites <u>in citu</u>.

4.5 The Doulting District

The Doulting Conglomerate bed, of the Doulting district, near Shepton Mallet, (Somerset), - see Fig. 1 - a thin bioclastic limestone (- 0.40m.), with numerous iron stained oncolites, was once well exposed in the Doulting rail-way cutting, (ST645425- Richardson, 1907, p.390, bed vi). This bed can now only be seen in temporary exposures and in fragments excavated by rabbits and badgers! From such unpromising sources the following basal Subfurcatum Zone fauna has been recorded, (Parsons, 1975) - <u>Cadonites deslongehamosi</u> (d'Orb.), <u>Stephanoceras</u> sp., <u>Teloceras banksi</u> (Sow.), <u>Strenoceras</u> (<u>Strenoceras</u>) <u>subfurcatum</u> (Zieten), <u>Orthorarantiana</u> sp. and <u>Leptosphinetes</u> aff. <u>davidsoni</u> (S.B.). This is the most northerly, fully authenticated, Subfurcatum Zone fauna yet found in the British Isles. 5. THE HORIZONS OF SMITH'S, SOWERBYS' AND BUCKEAN'S TYPES

One of the prime aims of this work was to determine the stratigraphic range of the numerous ammonite species which have been described from the Humphriesianum/Subfurcatum Zones of southern England. These fall essentially into two groups, firstly those described by earlier workers, such as W. Smith (1817) and J. and J. de C. Sowerby, (1812-46) at the beginning of the nineteenth century and secondly those described by S. Buckman, (1831, 1883, & 1909-30). I was fortunate in being able to locate topotypes of most of these species and thus to determine their type horizon, often for the first time.

5.1 Smith's and Sowerbys' specimens

the

5.1.1 Stephanoceras calix (N. Smith, 1817)

The lectotype of this, the earliest species described from these beds in England, is said to have come from Sherborne; its matrix would suggest a provenance from beds equivalent to 4b/c at Oborne Wood. This specimen, (DMMH, C671) refigured by Cox (1930, P1.12, fig.10) is wholly septate, and, contrary to Cox's opinion, is very different to the following species.

5.1.2 <u>Stephanoceras brodiei</u> (J. Sow.)

The holotype of this species, (BMUM,43905), was said to have been found on the Isle of Portland, by J. Brodie, (Sowerby, J., 1822 in 1812-46), but the matrix would indicate a provenance from the 'iron-shot' Inferior Oolite of the Sherborne area. Contrary to Cox's opinion (Cox, 1930, p.303) this specimen is not conspecific with the previous species, <u>S. calix</u>, although in terms of preservation it is extremely similar. The matrix would suggest beds 4b/c at Oborne Wood as the probable type horizon; somewhat similar topotypes have been acquired from bed 4c.

5.1.3 Stephanoceras humphriesianum (J. de C. Sow.)

The lectotype of this species (Buckman & Secretary, 1908, Pl. vii, fig. la & b) came from the Sherborne area, but there is insufficient matrix on this sectioned specimen, (BMNH,43908a) to enable an accurate idea of the original bed to be obtained. However several topotypes, with a similar excellent style of preservation have been found in Oborne Wood bed 4c, the undoubted type horizon.

5.1.4 <u>Teloceras banksi</u> (J. Sow.)

The holotype of this species (BMNH,43910) has a highly characteristic matrix, and comparable specimens have only been/found at Oborne Wood and Frogden quarry ibed 6a, the undoubted type horizon.

5.1.5 Teloceras blagdeni (J. Sow.)

The type specimen of this species (BENH,43908) has a unique mode of preservation, as may clearly be seen in Buckman's refiguring of it (Buckman & Secretary, 1908, Pl. III, fig.1); the back of the ammonite has been planed off, and a later limestone bed, containing Perisphinctid ammonites, 'welded' on. This mode of preservation is typical of bed 5a Oborne Wood, where numerous specimens of <u>T. blacteni</u> have been found, many of which, especially at the top of the bed, are flat lying. These have been eroded in half, and had the overlying limestones, cemented to the eroded surface. This then is unquestionably the original type horizon of this species.

5.2 Buchman's Specimens

5.2.1 'Type Ammonites'

Buckman in a series of papers from 1881 onwards described and figured a large number of ammonite species from these two zones. Whilst Buckman's early stratigraphic work is a classic of its kind. his later published work often had little or no factual basis. In 'Type Ammonites' in particular, (1909-30 Buckman) figured more than 35 new species from the Dorset Humphriesianum/Subfurcatum Zones, of which only a small fraction were founded on stratigraphically well localised material. The 'hemerae' to which Buckman assigned these ammonites were based more on differences in matrix and inferred phylogenetic links, than on actual stratigraphic evidence. Whilst this sorry tale is true of most of the hemerae erected in 'Type Ammonites', it is particularly relevant to those here included in the Humphriesianum Zone (see Table 1), since many of these hemerae are some of the most inadequate members of Euckman's polyhemeral scheme. An analysis of the ammonite species included by Buckman in his hemerae, shows that those hemerae now relegated to the Humphriesianum Zone were pure artifacts of Buchman's imagination, bearing no relationship to the actual stratigraphic distribution of the ammonites.

Most of the specimens figured by Buckman from the Sherborne area came from the old Frogden quarry. Oborne Wood section has shown an identical succession to this quarry, and it has provided topotypes of virtually all the species described from this area. The stratigraphic horizons of these topotypes is given in Table 2 : the more doubtful topotypes are indicated by a question mark, since these indicate specimens which either may not be from the type horizon of the holotype, or which show some differences in preservation or matrix. Several other localities in the Dorset area yielded ammonites which

AMMONITE SPECIES	VOLUME	PLATE	oBO	INI) }	ND	FR	aab	EN B	ED	Nº's
	NQ.	NQ.	<u>6</u> a	c	b	a	5b	5a	4c	ъ	a
Cadomoceras simulacrum	5	458	T								
Plectostrigites symplectus	5	471	Т								
Flexoxyites flexus	5	525		T							
Oppelina pulchra	6	670		т	?			-			
Poecilomorphus primiferous	7	756								т	
Stegoxyites parcicarinatus	5	474								'n	
Cadomites hom.logaster	5	543	{		Т						
C. septicostatus	5	432			?						
Gibbistephanus gibbosus	7	780							?		
Epalaxites formosus	3	151							т		
E. latansatus	3	159								т	
Pheulostephanus paululus	7	754								Т	
Teloceras banksii J.Sow.	6	660				r					
T. multinodus Quenst.	7	788					x				
Chondroceras delphinus	5	431	•							т	
C. gervillii J.Sow.	7	724	*							x	
C. grandiforme	4	357						1	a de la composition de la comp	т	
C. wrighti	4	415			:					Т	
Baculatoceras baculatum Qu.	6	581	x							÷.,	1.6
Caumontisphinctes aplous	3	241					T			1.6	
C. bifurcus	3	192			T						
C. polygyralis	3	163			T						$\frac{1}{2} = \lambda^{2}$
Strenoceras apleurum	3	239		т							
Rhabdoites rhabdodes	4	374	т								
Lep <u>tosphinctes</u> cleistus	3	161			Т						
L. coronarius	3	202			?						
L. davidsoni	3	201			?		?				
L. leptus	3	160	?		?						
Sonninites alsoticus Haug	5	528					and a		ad Series Production	x	ang kalèn n Ng kalèn n

Table 2,

The type horizons of some of those species of ammonites described by Buckman (1909-30), from Frogden Quarry and the Oborne District, Dorset.

T = topotype

? = possible topotype, original horizon uncertain

X = original horizon of Buckman's specimen

Buckman figured in 'Type Ammonites', namely Milborne Wick, Clatcombe, Louse Hill, and Burton Bradstock. Only at the first two of these localities are the beds sufficiently expanded to yield any reliable stratigraphic information, hence only these will be discussed here.

(a) Clatcombe Farm

Buckman acquired a large quantity of material from the old quarries in the area immediately to the north of Sherborne, around Clatcombe farm. Whilst the Humphriesianum Zone rocks of this district are virtually identical to those of Oborne, the Subfurcatum Zone is much more condensed; it is represented by one thin bed (bed 6, section 4.1.1). The specimens described from this latter horizon, <u>Caumontisphinetes phaulus</u> and <u>C. nodatus</u> S.B., could thus be Banksi-Polygyralis subzones in age, as there is no evidence for the Baculatum subzone in this bed. The specimen of <u>Poscilomorphus regulatus</u> figured by Buckman (1909-30, Plate 746) from this district must have come from the equivalent of bed 5b at Clatcombe Farm, which is on exactly the same horizon as bed 4b Chorne Wood. Topotypes of all three of these species have been collected from Clatcombe.

(b) Milborne Wick

Whilst Humphriesianum Zone fossils are extremely common in this district, Buckman described only one 'species' from here; <u>Poecilo-</u><u>morphus angulinus</u> S.B. (1909-30, Plate 757); of which topotypes have been collected from bed 5, Milborne Wick lane section.

5.2.2 The Monograph

It is fortunate that the Monograph of the Inferior Colite Ammonites (Buckman, 1837-1907), ceased publication before Buckman had a chance to 'get to grips' with the majority of Eajocian ammonite

groups. Only a small number of ammonites from the Humphriesianua Zone were figured in this work, notably members of the genera Poecilomorphus and Dorsetensia, and they all originate from the Romani subzone. Topotypes of Poecilomorphus evolutus (Buckman, 1887-1907, Pl. 22, figs. 21 & 22) have been collected from the 'Irony Bed' of Louse Hill, whilst topotypes of P. asper (loc. cit., figs. 3 & 4) and P. cappilaceous (loc. cit., figs 11 & 12) have been collected from bed 4b, Oborne Wood. Specimens similar to the syntypes of Dorsetensia regrediens (Haug), (op. cit., P1.52, figs. 8-24) have been collected from beds 4b-c, Oborne Wood, whilst a specimen of the fine ribbed variant of this species (loc. cit., figs. 11 & 12), has been collected from bed 5, Milborne Wick. Similarly topotypes of D. complanata (op. cit. Pl.53, figs 3-5), D. pulchra (op. cit. Pl.52, figs. 25-27), D. subtecta (op. cit. Pl.54, figs. 4 & 5), D. liostraca (op. cit. P1.55, fig. 3) and D. tecta (on. cit. P1.56, figs. 2-4) have been collected from beds 4a-b, Oborne Wood.

6. CONCLUSIONS

6.1 The Geographical distribution of the Humphriesianum and Subfurcatum Zones.

Buckman (1923, in 1909-30, p.54), considered that the present patches of Humphriesianum/Subfurcatum Zone sediments represented relicts of once more extensive deposits. His argument was that subsequent erosion, especially during the 'Vesulian Transgression', (basal Garantiana Zone), had removed most of the rocks of this age in southern England. That reworking and erosion was at least partly responsible for the present restricted distribution of these Zones is confirmed by their fragmentary and conglomeratic nature. However

Frogden ammonite quarry species The and within the Subfur Oborne graphic poor ribu tum Zone 50 rocks 30 the 30 main



these rocks, where preserved, invariably consist of thin, 'ironshot' limestones, highly fossiliferous and with abundant evidence (fat-all d al.,1972) of a slow rate of deposition. There is thus every reason to suspect that these beds were not of any great thickness to start with. Their present distribution is thus as much due to initial fest-ided

deposition, as to subsequent erosion. In these circumstances, it is certain that the original area of deposition was considerably larger than the present restricted outcrop pattern would suggest.

It is now certain that Morley-Davies's concept of a northerly overlap of the Humphriesianum/Subfurcatum rocks by the basal Garantiana Zone transgression, (Norley-Davies, 1930, p.232; Arkell, 1933, p.235), is untenable. The Doulting Conglomerate, of Subfurcatum Zone age, outcrops well to the north of the Sherborne district, where the main, isolated patches of the Subfurcatum/ Humphriesianum Zone rocks occur. This northerly occurrence of Subfurcatum Zone rocks thus rules out any possibility of the more southerly outcrops having been over lapped. The more convincing explanation is Buckman's original one : that the existing outcrops are relicts of once more continuous deposits, broken up and most commonly removed by subsequent erosion.

6.2 The Stratigraphic distribution of ammonites in the Humphriesianum/ Subfurcatum Zones.

The only sections in Southern England which show thick enough deposits of these Zones to warrant a study of the stratigraphic distribution of their ammonites are Frogden Quarry and Oborne Wood, (sections 4.12-Jhere). The exposures in the Subfurcatum Zone rocks at these two localities are identical, thus the ammonite records from

the ammonite Oborne species Wood The within ő strat the graphi Humphri â bu ũn Ë 20 ñc 0 Ē, the 10 ö To main 05



these beds have been combined in text figure 5. The Humphriesianum Zone beds are now only exposed at Oborne Wood as this part of the section at Frogden is now grassed over; thus the distribution of the more common ammonite species from the Humphriesianum Zone of this former locality is shown in text figure 6. Lastly the overall distribution of the major genera within the Humphriesianum/Subfurcatum Zones of the Sherborne district is shown in text figure 7. To summarise this data, all the subzones are clearly recognisable in terms of the distribution of species, although the base of the Romani subzone, the top of the Humphriesianum subzone and the base of the Banksi and Polycyralis subzones are perhaps the most clearly delimited. At higher taxonomic levels, the greatest turnover of genera and subgenera, comes at the base of the Humphriesianum Zone, and at the base of the Polygyralis subzone of the Subfurcatum Zone. Many taxa however, particularly those within the Haplocerataceae, are wide ranging.

6.3 The Standard Zonal Scheme

6.3.1 The Zone of <u>Stephanoceras</u> (<u>Stephanoceras</u>) <u>humphriesianum</u> (J. de C. Sow.)

Author:-Oppel, 1856.Type area:-Schwabian Albe, south Germany.Type horizon:-The beds above the Elau Kalke and below the
Subfurcatum Oolithe.Fauna:-At its base it is characterised by the abundance
of Dorsetensia, whilst Stephanoceras and Teloceras

are abundant throughout.

Subfur ammonite ó p, tuin Sonc Zon The and 3 subgenera tigraphi 50 0 within Sherb dj. 20 bution llumphricsianum ...nd よってアン o σ 3 the ŝ Uorset. major





WERE AND AND A WAR AND A WAR

(i) The subzone of <u>Dorsetensia</u> romani (Oppel)

Author:-	Haug, 1891, restricted Nuller, 1941.
Type area:-	The Basse Alpes, south-east France.
Type horizon:-	The 'Couches à amonites ferrugineuses de Beaumont'
	(Haug, 1891, p.70) near Digne, south-east France
	(Pavia & Sturani, 1968, p.312).
Fauna:-	This horizon is characterised by a fauna, which

This horizon is characterised by a fauna, which includes abundant <u>Dorsetensia</u> - <u>D. romani</u>,

<u>D. eduardiana</u> (d'Orb.), <u>D. complanata</u> Buckman and <u>D. subtecta</u> Buckman - as well as <u>Chondroceras evolvescens</u>, <u>Sphaeroceras brongniarti</u>, <u>Teloceras blagdeniformis</u>, <u>Epalaxites portitor</u>, <u>Poecilomorphus cycloides and Stegozyites parcicari-</u> <u>natus</u>.

Typical British horizon:- Oborne Wood, bed 4b.

(ii) The sul	ozone of <u>Stephanoceras</u> (S.) <u>humphricsianum</u>
Author:-	Oppel, 1856, restricted Muller, 1941.
Type area:-	The Schwabian Albe.
Type horizon:-	To be designated.
Fauna :-	This horizon is characterised by the abundance of
	Stephanoceras s. str., including S. crassicostatum
	(Qu. emend. Renz), S. humphriesianum, S. pyritosum,
	(Qu. emend. Renz) and S. mutabile (Qu. emend. Renz).
Typical British horizon:-	Oborne Wood, bed 4c.

(iii) The subzone of <u>Teloceras</u> (<u>Teloceras</u>) <u>blazdani</u> (J. Sow.) Author:- Six, 1879, restricted Spath, 1936.

Type area:-	Ardennes, north-east France.		
Type horizon:-	To be designated.		
Fauna:-	This is a range Zone, based on the abundant		
	occurrence of Teloceras blagdeni and other members		
	of this genus.		
Typical British horizon:-	Bed 5a Oborne Wood.		
6.3.2 The Zone of <u>Strenoceras</u> (<u>Strenoceras</u>) <u>subfurcatum</u> (Zieten)			
Author:-	Terquem and Jourdy, 1869.		
Type area:-	Departement de la Moselle, France.		
Type horizon:-	Marnes a Longwy (in part).		

Characterised at its base by the appearance of Fauna:-Leptosphinctes and Caunontisphinctes, the main feature of this Zone is the abundance of Garantiana and Strenoceran.

The subzone of Teloceras (Teloceras) banksi (J. Sow.) (i)

Author:-	Buckman, 1910, restricted Sturani 1971.
Type area:-	Sherborne area, North Dorset.
Type horizon:-	Buckman's bed 5, Frogden quarry.
Fauna:-	This horizon is marked by the appearance of
	Leptosphinctes and Caumontisphinctes and by the
	presence of the last Stephanoceratids which were
	typical of the Humphricsianum Zone, such as <u>Teloceras</u>
	and Stenhanoceras s. str.

Typical British horizon:-

Bed 5b Frogden quarry.

115

(ii) The subzone of <u>Caumontisphinctes</u> (<u>Caumontisphinctes</u>) polygyralis S. Buckman

Author:- Sturani, 1971.

Type area:- Digne, Basse-Alpes, South-east France.

Type horizon:- The limestone/marl sequence, particularly of the 'Ravin du Feston', (Pavia & Sturani, 1968), beds 119-138, (Pavia 1973).

Fauna:-

This horizon is marked by the abundance of, the dimorphic pair <u>Caumontisphinetes</u> (C.) <u>polysyralis</u> and <u>C. (Infraparkinsonia) phaula S.B., Leptosphinetes</u> <u>davidsoni</u>, and <u>Orthograntiana haugi</u> Favia, as well as by the first appearance of the <u>Genus Strenoceras</u>.

Typical British horizon:-

Bed 6b Frogden Quarry.

(iii) The subzone of <u>Strenoceras</u> (<u>Garantians</u>) <u>baculata</u> (Qu.)
Author:- Kumm 1952, restricted Gabilly <u>et al</u>. 1971.
Type area:- Bielefeld area, North-west Germany.
Type horizon:- 'Untere Strenoceraten Schichten', (Althoff 1923).
Fauna:- This horizon is marked by the abundance of <u>Garantiana</u>

baculata and <u>Strenoceras subfurcatur</u>, as well as by the appearance of the Spiroceratid camonites,

App<u>soroceras</u> <u>baulatum</u> (Qu.) in particular.

Typical British horizon:- Bed 6d, Frogden Quarry.

7. REFERENCES

Althoff, W., von, 1914,

Die geologischen Aufschlüsse Bielefelds. Bericht. Katurwiss. ver. Bielefeld (1911-13), 3: 195-225.

, 1928,

Übersicht über die Gliederung der mesozoischen Schichten bei Bielefeld. <u>Idem, 5</u>: 1-

Arkell, N., J., 1933, (reprinted 1970),

The Jurassic System in Great Britain. Oxford University Press; xii + 681 pp. 41 pls.

1954, in

Arkell, N. & Playford, P.E. 1954,

The Bajocian Ammonites of Western Australia. Phil. Trans. Roy. Soc., (B), 237: 547-605, pls 27-40.

1956,

The Jurassic Geology of the World. Oliver & Boyd, Edinburgh; xv + 806 pp., 46 pl.

Bentz, A. 1924,

1928,

Die Garantienschichten von Norddeutschland mit besonderer Berucksichtigung des Brauneisensolithhorizontes von Harzburg. Jahrb. Preuss. Geolog. Landesanst., <u>45</u> : 119-193, pl. 4-9.

Ü ber Strenoceraten und Garantianen insbesondere aus dem Mittleren Dogger von Bielefeld. <u>Idem, 49</u>: 138-206, pl. 14-19. 「おいいの」ののないないないないので、

Bomaford, G., 1948,

New sections in the Inferior Colite. Proc. Geol. Assoc., <u>59</u>: 148-150.

Erasil, L., 1895,

Observations sur le Bajocien de Normandie. Bul. Lab. Geol. Fac. Sci. Caen, <u>2</u>: 222-248.

Buckman, S. S., 1881,

A descriptive catalogue of some of the species of ammonites from the Inferior Colite of Dorset. Quart. J. geol. Soc. Lond., <u>37</u>: 588-608.

1883,

Some new species of ammonites from the Inferior Colite. Proc. Dorset Nat. Hist. Archaeol. Soc., 4: 137-146, pls 1-4.

1887-1907,

A Monograph of the Ammonites of the Inferior Oolite Series. Palaeontogr. Soc. : cclxii + 456 pp., 127 pls.

1891,

On certain Ammonite-zones of Dorset and Somerset. Report Brit. Assoc. Adv. Sci. (1891), section C, (5) : 655-6.

1893.

The Bajocian of the Sherborne district; its relation to subjacent and superjacent strata. Quart. J. geol. Soc. Lond., <u>49</u>: 479-522.

1898,

On the Grouping of some Divisions of so-called 'Jurassic' Time. Idem., 54 : 442-462. Buckman, S. S., 1909-30,

Yorkshire Type Ammonites - continued as Type Ammonites. London & Thame, vols. 1-7, : 790 pls.

1910,

Certain Jurassic (Lias-Oolite)strata of South Dorset. Quart.

J. geol. Soc. Lond., <u>66</u> : 52-89.

____, & 'Secretary', 1908,

Illustrations of Type Specimens of Inferior Colite ammonites in the Sowerby Collection. Palaeontogr. Soc.; 7 pls.

Cox, L. R., 1930,

On British fossils named by William Smith. Ann. Mag. Nat. Hist.,

33 : 287-304, pl.12.

Cross, J.E., 1875,

The geology of N.W. Lincolnshire. Quart. J. geol. Soc. Lond.,

31 : 115-129.

Dorn, P., 1935,

Die Hammatoceraten, Sonninien, Ludwigien, Dorsetensien und Witchellich des süddeutschen insbesondere fränkischen Doggers. Palaeontographica, <u>82</u>(A) : 1-124, pls. 1-29.

Gabilly, J., Contini, D., Nouterde, R., & Rioult, N., 1971,

Les Zones du Bajocien. C.R.S. Soc. geol. Fr., : 85-3.

Gatrall, M., Jenkyns, H., & Parsons, C., 1972,

Limonitic concretions from the European Jurassic, with particular reference to the 'snuff-boxes' of Southern England. Sedimentology, <u>18</u> : 79-103.

Gosselet, M. J., 1881 in 1880-3,

Esquisse Géologique du Nord de la France. 2e fasicule -Terraine Secondaires. Lille (1881); pp. 169-278, 20 pls.

Haug, E., 1891,

Les Chaînes Subalpines entre Gap et Digne. Bull. Serv. Carte géol. Fr., 3, (21) : 1-199.

____ 1893,

Etude sur les Ammonites des Etages moyen du System Jurassique.

Bull. Soc. geol. Fr., 3e ser. 20 : 277-333, pls. 7-10.

Hudleston, W. H., 1886,

Excursion to Sherborne and Bridport. Proc. Geol. Assoc., 9: 187-199.

1887-96,

A monograph of the British Jurassic Gasteropoda; 1. The Gasteropoda of the Inferior Colite. Palaeontogr. Soc. : 514 pp., 44 pls.

Huf, N., 1968,

Uber Sonninien und Dorsetensien aus dem Bajocium von Nordwestdeutschland. Beih. geol. Jb., <u>64</u> : 126 pp., 51 pls.

Kellaway, G.A., & Wilson, V., 1941,

An outline of the Geology of Yeovil, Sherborne and Sparkford Vale. Proc. Geol. Assoc., <u>52</u>: 131-174.

Kumm, A., 1952,

Das Mesozoikum in Niedersachsen : Der Dogger. Geol. u. Lagerstatten Niedersachsens 2 ; Abt. 2: 329-509. Geological Highlights of the West Country. Butterworths, London: 296 pp.

Mascke, E., 1907,

Die Stephanoceras-Verwandten in den Coronatenschichten von Norddeutschland. Inaugural-Dissertation, University of Gottingen : 38 pp.

Morley-Davies, A.M., 1930,

The Geological life work of Sidney, Savory, Buckman. Proc. Geol. Assoc., 41 : 231-4.

Morton, N., 1965,

The Bearreraig Sandstone Series (Middle Jurassic), of Skye and Raasay. Scot. J. Geol., <u>1</u>: 189-216.

_ 1971,

Some Bajocian ammonites from western Scotland. Palaeontology, 14: 266-293, pls. 40-51.

Muller, S.W., 1941,

Standard of the Jurassic System. Bul. Geol. Soc. Amer., <u>52</u>: 1427-1444.

Oppel, A., 1856-8,

Die Juraformation Englands, Frankreichs und des südwestlichen Deutschlands. Jb. Ver. vaterländ. Naturkde. Württ. <u>12-14</u>: 1-837.

Pavia, G., 1973,

Ammoniti del Baiociano Superiore di Digne. Boll. Soc. Fal.

Italy, 10 : 75-142, pls. 13-29.
Pavia, G. & Sturani, C., 1968,

Étude biostratigraphique du Bajocien des Chaînes Subalpines aux environs de Digne (Basses-Alpes). Boll. Soc. Geol. It., <u>87</u>: 305-316.

Parsons, C.F., 1972,

A new temporary section in the Inferior Colite of South Dorset. Proc. Dorset Nat. Hist. Archaeol. Soc., <u>93</u> : 117-8.

____ 1975,

Ammonites from the Doulting Conglomerate Eed (Upper Bajocian, Middle Jurassic) of Doulting, Somerset. Palaeontology, <u>18</u>(1): 191-205, 1pl.

Richardson, L.R., 1907,

The Inferior Colite and Contiguous Deposits of the Bath-Doulting District. Quart. J. geol. Soc. Lond., <u>63</u> : 383-423.

1915,

Report of an excursion to Bridport, Beaminster and Crewkerne. Proc. Geol. Assoc., <u>26</u> : 47-78.

1916,

The Inferior Colite and Contiguous Deposits of the Doulting-Milborne Port district (Somerset). Quart. J. geol. Soc. Lond., <u>71</u>: 473-520, pls. 40-3.

1928-30,

The Inferior Colite and Contiguous Deposits of the Burton Bradstock - Broadwindsor district, Dorset. Proc. Cotteswold Nat. F. Club, <u>23</u>: 35-68, 149-185, 253-264.

1931,

The Inferior Colite and Contiguous Deposits of the Sherborna District, Dorset. <u>Idem., 24</u> : 35-65, pls. vi-xi.

Senior, J.R., Parsons, C.F., & Torrens, H.S., 1970,

New sections in the Inferior Oolite of south Dorset. Froc.

Dorset Nat. Hist. Archaeol. Soc., 91 : 114-9.

```
Six, M.A., 1879,
```

Comptes rendues des excursion dans les Terraines Secondaries de l'Aisne et des Ardennes. Ann. Soc. geol. Nord (Lille), <u>6</u>: 400-427.

```
Smith. W.. 1817.
```

Stratigraphical System of Organised Fossils. Williams, London, xi + 118 pp.

Sowerby, J., & Sowerby, J. de C., 1812-46,

The Mineral Conchology of Great Britain. London : 1287 pp. 648 pls.

Spath, L.F., 1936,

On Bajocian ammonites and belemnites from eastern Persia (Iran).

Palaeont. Indica, (N.S.) 22 : 1-21, 1 pl.

Stahlecker, G., 1935,

Stratigraphie und Tektonik des Braunen Jura im Gebiet des Stuifen und Nechberg. Inaugural-Dissertation, University of Tübingen; 62 pp.

Sturani, C., 1971,

Ammonites and Stratigraphy of the <u>Posidonia alpina</u> beds of the Venetian Alps. Mem. Ist. Geol. Mineral. Padova, <u>20</u>: 190 pp., 16 pls.

Torrens, H.S., 1969, (Editor),

International Field Symposium on the British Jurassic, Excursion Guide 1 : Dorset and South Somerset. University of Keele : 71

pp.

Terquen, 0., & Jourdy, E., 1869,

Monographie de l'étage Bathonien dans le Département de la Moselle. Mém. Soc. géol. France, (2), <u>9</u>:

Waagen, W., 1867,

Über die Zone des <u>Ammonites sowerbyi</u>. Geogn.-Paläont., <u>1</u>: 511-607, pls. 24-29.

Westermann, G.E.G., 1954,

Monographie der Otoitidae (Ammonoidea). Beih. Geol. Jb., <u>15</u>: 1-364, 33 pls.

_____ 1967,

Jurassique moyen (Alpes exclues), Lexique Strat. Int. vol. 1; Europe, Fasc. 5; Allemagne. Paris : 197 pp.

Whicher, J. & Palmer, C., 1971,

Preliminary description of a temporary exposure in the Middle Inferior Oolite, near Sherborne, Dorset. Proc. Dorset Nat. Hist. Archaeol. Soc., <u>92</u>: 110-9.

Noodward, H.B., 1894,

The Jurassic rocks of Britain; vol 4, the Lower Colitic rocks of England. Nem. Geol. Surv. G.B. : 628pp. そうして、このでは、「ない」では、こので、「ない」では、「ない」のでは、「ない」ので、「ない」で、「いい」ので、「ない」ので、「ない」ので、「ない」ので、「ない」ので、「ない」ので、「ない」ので、「ない」

1C. A STRATIGRAPHIC REVISION OF THE garantiana AND parkinsoni ZONE

(UPPER BAJOCIAN, MIDDLE JURASSIC) ROCKS OF SHERBORNE, DORSET

C.F. Parsons

SUMMARY : A new formal lithostratigraphic scheme for the Stroud Formation of the Inferior Oolite Group, in the Sherborne area of Dorset, is erected. The development of the zonal and subzonal scheme for the <u>parkinsoni</u> and <u>garantiana</u> Zones is discussed in the light of new evidence from Dorset. All the currently accepted subzones of the <u>garantiana</u> Zone (<u>dichotoma</u>, <u>subgaranti</u>, <u>tetragona</u> and <u>acris</u>) are recognized in Britain for the first time. All the existing exposures of <u>garantiana/parkinsoni</u> Zone rocks in the Sherborne area are described, their ammonite faunas discussed and correlated with other areas of southern England and Europe.

The Sherborne district of north Dorset has a considerable historical significance as the type area for the <u>garantiana</u> Zone and the <u>truellei</u> subzone of the <u>parkinsoni</u> Zone (Upper Bajocian, Middle Jurassic). The need for a detailed stratigraphic revision of the beds of this age has long been recognized (Richardson, 1932, p.82). The aim of the present work is to undertake a revision of the ammonite faunas from all the existing exposures in the area of <u>garantiana/parkinsoni</u> Zone rocks, (see Fig.1). These rocks, which have been previously included in the 'Upper Inferior Colite', are a continuation of the Stroud Formation of the Inferior Colite Group, a unit recognized in the Cotswolds, but which has been shown to be laterally persistent across much of southern England (Parsons, 1979a). The exposures



Figure 1.

A sketch map of the Sherborne district, showing the location of the main exposures described in the text. Locality 1 = Bradford Abbas railway cutting,3 = Louse Hill Quarry, the rest will be found on Figure 7. in the immediate Sherborne area (localities 4-11, Figs.1 & 7), are dominated by extensive faces of <u>garantiana</u> Zone beds, and the <u>parkinsoni</u> Zone has only been recognized <u>in situ</u> at two small, isolated exposures. The latter unit is undoubtedly widely represented, although inaccessible at the top of the high, vertical quarry faces, thus only the <u>garantiana</u> Zone can be treated in any detail. Because of the historical, and international significance of the area, the development of the european zonal and subzonal divisions of the <u>garantiana</u> and <u>parkinsoni</u> Zones have been discussed in detail, and in part re-interpreted in the light of the new Dorset evidence. All of these Zones and subzones are essentially 'concurrent-range biozones' (Holland, <u>et al</u>., pp.13-14; Hedberg, 1976, pp.56-8), although the overall ranges of several taxa in the Sherborne area, are still to be precisely determined.

A new formal, hierarchical, lithostratigraphic terminology for the Stroud Formation in the Sherborne area has been erected (see Figure 2). Although exposures are now poor in the upper part, the new terminology has not been restricted to the <u>garantiana</u> Zone beds, and the whole Formation has been divided into Members and Beds of mainly local significance. There is a broad dichotomy of facies, between the thin, ferruginous, highly 'condensed' beds found in the west (localities 1-3, Fig.1), and the thicker, sparsely glauconitic, 'expanded' facies found to the east. Although the exact point of transition between the two cannot be seen, it must be extremely sharp and clear cut. The change in thickness is certainly more abrupt than that shown by Gatrall <u>et al</u>. (1972, fig.1), since the supposed intervening <u>garantiana</u> Zone, ferruginous, intraclastic limestone recorded (ST617,165), by them (op. cit., fig.3), is now known to be of upper 「日本のなる」というと



Figure 2.

The proposed lithostratigraphic terminology for the ^{subdivisions} of the Stroud Formation found in the Sherborne area of north Dorset.

parkinsoni Zone age (Whicher, 1969). The garantiana Zone/truellei subzone beds at this point are still highly attenuated, (op. cit.). The northern and eastern boundaries of the 'expanded facies' are less certain. The lack of ammonites from the small and sporadic exposures in the north around Charlton Horethorne and Blackford, confuses our understanding of the lateral passage into the 'Hadspen stone' (see Gatrall, et al., 1972, fig.3). To the east the Sherborne and Combe Limestone Members appear to pass, virtually unchanged beneath the Bathonian outcrop, and have been detected at the subsurface in the Stowell borehole (Pringle, 1909, 1910).

LITHOSTRATIGRAPHY

The 'Upper Inferior Oolite' of the Sherborne district (Milborne Wick-Sandford Lane; see fig.1), has in the past been divided between three, broad, informal units : the 'Sherborne building-stone' (Richardson, 1932), the 'Rubbly beds' (Buckman, 1893) and 'Crackment Limestones' (Osborne-White, 1923). These units, with some modification and subdivision, largely form the basis for the formal members defined below (see fig.2). To the west of Sherborne, the above beds rapidly attenuate, and are replaced by thin, highly fossiliferous, 'condensed' beds.

Vertical limits of the Formation.

The base of the Stroud Formation

The lower boundary of the Stroud Formation (largely equivalent to the 'Upper Inferior Oolite' <u>sensu</u> Arkell, 1933), is here drawn at a well defined disconformity, with an iron-stained conglomerate, at the base of the Acanthothiris Beds of the Sherborne Limestone Member. These latter

beds rest on 'iron-shot' <u>subfurcatum</u> Zone beds at Frogden quarry (Parsons, 1976a, p.125), whilst at Sandford Lane, they have cut down to the <u>sauzei</u> Zone top of the 'fossil-bed' (Parsons, 1974, p.164). Although the <u>subfurcatum</u> Zone beds (= the '<u>cadomensis</u> beds', Hudleston, 1886), have previously been included in the 'Upper Inferior Oolite' (Kellaway & Wilson, 1941; Wilson <u>et al.</u>, 1959, p.96), they clearly lie beneath the above mentioned disconformity. In addition the lithology and general facies of these beds has more in common with the equally 'condensed' <u>humphriesianum</u>/ <u>sauzei</u> Zone rocks, than with the overlying, more 'expanded', <u>garantiana</u> Zone beds.

The upper boundary of the Stroud Formation

The top of the Stroud Formation in the Sherborne district is less easily defined than its base. Richardson (1932) took this boundary at the top of what is now called the <u>bomfordi</u> subzone, and included the 'Limestone beds' (= Crackment Member) in the overlying Fuller's Earth 'series'. However, this is rather an arbitrary boundary, based more on palaeontological, rather than lithological criteria, and is difficult to trace in most sections. Subsequently, most authors (Kellaway & Wilson, 1941; Arkell, 1951-9; Fowler, 1957; Wilson <u>et al.</u>, 1959; Torrens, 1969) have included these beds in the Inferior Oolite Group, since the main mappable feature is produced by the top of the Crackment Limestones weathering out against the Fuller's Earth Clay. Although slightly problematic, the precise boundary can be taken below the '<u>knorri</u> clay', which acts as a basal marker bed for the Fuller's Earth Formation. The Lenthay Bed is thus included in the top of the Crackment Limestone Member (cf. Torrens, 1969, fig.2).

The Sherborne Limestone Member

The beds previously known as the 'Sherborne Euilding-stone' (Richardson, 1932) or 'Building-stone' (Buckman, 1893), and including the 'stone used for lime' (op. cit.), are here included in a single, new, formal unit : the Sherborne Limestone Member of the Stroud Formation. At its type locality, here designated as Sandford Lane quarry, it consists of 3.0m of massive, sparsely glauconitic and colitic, sandy, recrystallized biomicrite, with subsidiary marl partings (Buckman, 1893, p.492; Richardson, 1932, p.75, beds 4-6; see fig.3). Relatively unfossiliferous, with only sparse clusters of <u>Sphaeroidothyris</u>, these beds are at their thickest in the region of Redhole Lane quarry, where over 7.5m have been recorded (Richardson, 1932, p.73; see fig.5). Although generally massive, variation in the relative thickness of the marl partings allows some lithostratigraphic subdivision to be made.

Acanthothiris Beds

The lowest beds within the Sherborne Limestone Member, consisting of brown, sandy biomicrites, interbedded with thick, brown marls, are here formally named the Acanthothiris Beds, after the common occurrence of <u>A. obornensis</u> Buckman & Walker. At their type locality, Frogden quarry, they are 1.45m thick and consist of five, brown, sandy limestone courses (c.20cm thick), interbedded with brown sandy marls, underlain by a brown, ferruginous, rubbly limestone and marl, with a basal conglomerate, totalling 0.30m. A very similar sequence has been recorded from Oborne Wood (Farsons, 1976a, p.128), whilst the limestone courses are slightly thicker at Sandford Lane, where the beds are again 1.45m thick.

Building-stone Beds

The thick, massive biomicrites, which were formerly worked as a

※二,法言法出國法書中原,例将

'free-stone' are here formally named the Building-stone Beds, following Buckman's (1893, p.496), original usage. Although at their thickest in the region of Redhole Lane quarry, where more than 6.0m have been recorded (Richardson, 1932, p.73), only Sandford Lane, here designated the type locality, now shows a complete sequence of these beds, which are 1.5m thick (op. cit., p.76, bed 4).

Redhole Lane Beds

The thin bedded limestones previously known as the 'stone used for lime' (Buckman, 1893, p.496), which are less massive than the underlying Building-stone <u>sensu stricto</u>, are here formally named the Redhole Lane Beds. At their type locality, Redhole Lane quarry, they are made up of 1.5m of thin bedded biomicrites, with subsidiary marks and well burrowed, especially down towards the basal mark parting (Richardson, 1932, p.73, beds 4-5). These beds become thicker and more massive towards the east (e.g. Oborne Lane section - see fig.7), whilst to the west they become more modular and marky, so that at Sandford Lane (see fig.3), they are indistinguishable from the overlying Combe Member .

Combe Limestone Member

The beds previously known as the 'Rubbly beds' are here formally named the Combe Limestone Member, after the excellent exposures on either side of the 'Combe' (Combe quarry = Sandford Lane, Buckman, 1893, p.492). The term 'Rubbly beds' (<u>op. cit.</u>), is rejected as it has a prior use in the Portland Beds (Blake, 1880), as well as for other beds in the Inferior Oolite of the Eath district (Richardson, 1907). Redhole Lane quarry is here chosen as type locality, in order to prevent any uncertainty in relation to the boundary with the underlying Redhole Lane Beds. The maximum thickness of こうしんですい あちからか あまう ゆうかんちゃうかん

日本のですが

this unit is unknown, although in excess of 5.6m have been recorded from Redhole Lane (Richardson, 1932, p.73) whilst 13m can probably be assigned to this Member in the Oborne Borehole (Wilson <u>et al.</u>, 1959, p.96). Taking the first appearance of <u>Parkinsonia s. lat.</u> as a datum level, the base of this unit appears to be diachronous. Thus in the east at Milborne Wick (see fig.6), the first <u>Parkinsonia</u> is only 0.15m from the base, at Redhole Lane 2.2m (see fig.5), whilst at Sandford Lane, the most westerly exposure, they appear 3.0m above the base. This is partly illustrated in figure 7, where the first appearance of <u>P. rarecostata</u> is plotted. A degree of lithostratigraphic subdivision is possible, since the upper beds tend to be more massive.

Quarr Lane Beds

These beds, named after the excellent exposures either side of Quarr Lane, Sherborne, at Blackberry Lane and Redhole Lane quarries, are the typical 'Rubbly beds' <u>5</u>. <u>str</u>. At their type locality, here designated as Redhole Lane quarry, they consist of 5.0m (Richardson, 1932, p.73), of hard, grey, nodular, sparsely colitic, recrystallized biomicrite, highly bioturbated and interbedded with soft brown marl, to give an overall rubbly and nodular appearance. Fossils are rare, apart from distorted ammonite body-chambers, and the beds usually have a brown, iron-stained appearance, which separates them from the Crackment Limestone Member.

Goathill Beds

The <u>parkinsoni</u> Zone limestones previously recorded from Goathill (ST674122; Torrens, 1969, p.317) as 'Rubbly beds' are more evenly bedded than the bulk of the Combe Member. At this, the type section, and an exposure 100m to the south by the side of the path, the Goathill Eeds

consist of +2.4m of brown, sparsely colitic, fossiliferous biomicrites, evenly bedded, with subsidiary brown, marl partings. These beds are consistently more evenly bedded and fossiliferous, with common echinoids (<u>Holectypus</u>, <u>Pygomalus</u> and <u>Pygorhytis</u>), than the subjacent Quarr Lane Beds, and are very like the Burton Limestone on the Dorset coast (Parsons, 1975a). Unfortunately there is currently no good exposure of the base of these beds, although it may be represented at both the Milborne Port 'station cutting' (Richardson, 1916, pp.510-1) and at the Redhole Lane quarry, where 0.70m of more regularly bedded limestone has been recorded from the top of the section (Richardson, 1932, p.73). No complete exposure of these beds has been seen, thus their complete thickness is unknown, although the 3.0m of grey, shelly limestone overlying the Quarr Lane Beds (= 'Rubbly beds') in the Oborne borehole (Wilson, <u>et al.</u>, 1959, p.96) are probably their equivalent.

Crackment Limestone Member

The existing informal unit, the 'Crackment limestones' (Osborne-White, 1923; = 'the Limestone beds'; Buckman, 1893, p.486; Richardson, 1932, p.49), is here redefined as a formal Member, with the retention of the original type locality; Crackment or Crachment Hill (ST667183); selected by Arkell (in Donovan & Hemingway, 1963, p.103-4). Here only an upper part of the unit is visible, where it typically consists of thin limestones, alternating with relatively thick, brown clay/marl bands. These beds are very badly exposed at present, and most existing stratigraphic information has come from temporary sections (Arkell, 1951-9, p.10; Fowler, 1957; Torrens, 1969). Evidently the proportion of clay increases towards the top, and the Lenthay Bed, here defined as the topmost bed of the Crackment 「「「かななの」のである

Limestone Member, is separated from the bulk of the limestone below by 2.1m of clay (Torrens, 1969, p.315). Although lack of exposure makes any estimate of the total thickness of this unit difficult, in the region of 12m have been recorded from the Oborne borehole (Wilson <u>et al.</u>, 1959, p.96).

'Condensed' beds of Nether Compton

The lithostratigraphic units described above are applicable to the area between Sandford Lane and Milborne Wick. However, west of Sherborne many of these units, particularly the Sherborne and Combe Members, thin rapidly (Gatrall <u>et al.</u>, 1972, figs. 1 & 3), until at Halfway House, Nether Compton, they are represented only by some very thin, ferruginous, highly 'condensed' beds.

The Astarte Bed

The beds representing the whole of the <u>garantiana</u> Zone in much of north and south Dorset, are extremely thin (generally less than 0.20m), and of highly variable lithology. At its type locality, here designated as the Rock Cottage quarry, the Astarte Bed consists of 0.13-0.17m of moderately hard, light-grey, highly fossiliferous obliomicrite, with yellow/orange shell replacement, iron and manganese staining, sparse pisoliths and common, often serpulid encrusted <u>Astarte (Neocrassina)</u> and belemnites (Richardson, 1932, p.62). However, other lithologies are common, such as ferruginous conglomerates at Louse Hill and highly limonitic marls with 'snuff-boxes' (q.v. Gatrall <u>et al.</u>, 1972), at Bradford Abbas railwaycutting. Following previous usage all these are here included in the Astarte Bed (Buckman, 1893, p.487, = 'Shell bed', 'Rotten bed'; Richard「あい」からいの高の事の製肉の

son, 1932, p.48), including the overlying, impersistent marl bed, which is mainly the result of the weathering of the Astarte Bed <u>s. str.</u>, such as at Bradford Abbas (Buckman, 1893, p.485, bed 4 ='Marl bed' or 'Dirt bed').

Halfway House Bed

The horizon previously known as the Halfway House 'fossil-bed' (Buckman, 1893, p.487, bed 4; Richardson, 1932, p.62) is here formally named the Halfway House Bed. At its type locality, Rock Cottage quarry, it consists of 0.30m of soft, light-grey/off-white, biomicrite, with yellow shell replacement. At other localities it is thicker, with 'snuff-boxes' (e.g. Bradford Abbas railway-cutting, 0.70m), whilst elsewhere (e.g. Clifton Maybank, ST577140), it is more ferruginous, with abundant ammonites. Although precise correlation is difficult, this bed is probably a 'condensed' equivalent of parts of the Combe Member.

'Chapel beds'

West of Sherborne the upper parts of the Combe Member (mainly Goathill Beds) pass into a ferruginous, grey-brown, sandy, relatively unfossiliferous, thin bedded, sparsely crinoidal, iron-stained, intraclastic limestone, with brown, sandy marl partings. In the western part of the new road-cutting, adjacent to the old Chapel quarry, Halfway House (Wilson <u>et al.</u>, 1959, p.95), c.3.0m of these beds are visible. They have not been formally defined here, as their exact stratigraphic position is still problematic, although they are certainly, at least in part, of immediately post-<u>truellei</u> subzone age (Whicher, 1969, p.326). Their overall distribution is restricted. They first appear 3km west of Sherborne (ST617165; <u>loc. cit.</u>), and are also visible at Louse Hill (Richardson, 1932, p.67) and Rock Cottage quarry (op. cit. p.62), but are absent from the Bradford Abbas railwaycutting, where the Crackment Limestone Member rests directly on the Halfway House Bed. However, the presence of <u>Parkinsonia parkinsoni</u> (Buckman <u>non</u> Sow.), in the bottom of the Crackment Limestone at the latter locality suggests that a lateral facies change may have occurred, and that the base of the Crackment Limestone Member may be the equivalent here, of the 'Chapel beds' to the east.

BIOSTRATIGRAPHY

The age range of the Stroud Formation in the Sherborne area is similar to that of the 'parkinsoni zone' as originally defined (Oppel, 1856 in 1856-8, p.344). Although in his early papers on this area, Buckman (1881, 1883) utilized this zone in its original, broad sense, he rapidly recognized the need for its subdivision. Thus in 1891, Buckman used the 'cadomensis' (= subfurcatum Zone), 'truellei' and 'zigzag zones' at least in part to fulfil this need. In a subsequent paper (Buckman, 1894, tab.1), he clearly delimited these zones and compared them with the zonal schemes of earlier authors. A significant feature of these papers, relates to the introduction of the 'truellei zone', which was based on "the bed at Halfway House, which yields the large Parkinsoniae" (Buckman, 1891, p.656). The Halfway House Bed (= 'fossil-bed'), of the present area must thus be taken as the type horizon for this unit (Arkell in Donovan & Hemingway, 1963, p.355). Buckman (1893), in his detailed description of the Sherborne area introduced a further subdivision, in the form of a garantiana hemera between his niortensis (= previous 'cadomensis zone') and truellei hemerae, whilst he was later to use a schloenbachi hemera between his truellei and zigzag hemerae (Buckman in Richardson, 1908; Buckman, 1910, p.78). Although

135

		euoz tu	partauq		euo7	anofinanas			
SCHEME USED HEAE	zizzag Zona	boaford 14	true11e1		t e t r e t e t		dichotoma	subfurcatum Xone	
		•u02 Ţ	Larkinson		euoz wuwta	Rutan	**		
PARSONS 1976		boaford	truelle4						
TOPRENS 1969		to a la l	Line (1)		garanti ana	euoz			
ARKELL 1951-94 1956		Parkinsoni	8 107		Gar en tíana	e uo 7	ч		
		•aoz žnosn'izsq							
SPATH 1936	zigzag at	schloen- bachi subzone	truel) et		garahtia. Da seze			niortene- ie *.E.	
ARKELL 1935		Schloenbach1 Zone	Truelled Zone				a anti-para anta anti- anta anti- anta anti-		
RUCKNAN 1910		<u>echlounbechi</u> heacra	truelled heasts						
скили 1891. 1895 -			ue (1891) ane (1891)		8			rtensie GC	

1. N. . . N. .

「たる」を見たるとなった。

Table 1. Development of the classification of the garantiana and parkinsoni Zones in Great Britain.

Buckman (1909-30) introduced further, but less well defined hemerae, only the above mentioned units were used by Arkell (1933), as the basis for the zonal scheme, which he used in his description of the British 'Upper Inferior Oolite'. Spath (1936, p.15), returned to the original, wide definition of the 'parkinsoni zone', with the use of Buckman's hemerae as subzones. Arkell (1951-9, pp.8-9; 1956, p.31), subsequently preferred the use of a more restricted, but undivided <u>parkinsoni</u> Zone for World correlation, since he considered the <u>truellei</u> subzone to have no more than local significance. Up until recently (Parsons, 1976b, p.48), no attempt has been made to subdivide the British <u>garantiana</u> Zone. The development of the <u>garantiana</u> and <u>parkinsoni</u> Zones in Britain is summarized in Table 1.

The garantiana Zone

Buckman (1893, p.483), introduced the <u>garantiana</u> hemera in his description of an area, which largely coincides with that presently under discussion. Its introduction as a zone must also date from this publication (Arkell, 1951, in 1951-9, p.9), rather than from its subsequent use in a list of zones (Buckman, 1913 in 1909-30). The original theoretical differences between chronological (= 'hemerae') and strictly stratigraphical units (= zones), were misunderstood by many authors (see the discussion of Buckman's 1893 paper), and this clear cut division was in any case destroyed by Buckman's (1913 in 1909-30; 1915) later confused interchange of hemerae and zones (<u>contra</u>, Dean <u>et al.</u>, 1961, p.440). Additionally, it could be argued that Buckman's original hemerae (Buckman, 1893, pp.481-2; see Arkell, 1933, pp.20-1), have more in common with the present standard Zones (Holland <u>et al.</u>, 1978, p.7) or chronozones (Hedberg, 1976, p.67), than with the poorly defined 'biozones' then in use. The Sherborne

district must thus be taken as the type area for the garantiana Zone (Arkell in Donovan & Hemingway, 1963, p.144).

In his description of the garantiana hemera, Buckman (1893), recorded very few ammonites : "Parkinsonia garantiana, P. praecursor, P. rarecostata, Perisphinctes triplicatus, Oppelia sp. and Ancylyoceras sp.". Whilst these taxa must now be interpreted in a very broad sense, his records of "P. rarecostata" (op. cit., pp.496, 497), suggest that the garantiana Zone must overlap with the range of a species, which would now be referred to Parkinsonia s. str. This is confirmed by his subsequent assignation of "evolute Parkinsonae" and "Garantiana" to the ... "General Ammonite Facies" .. of the garantiana hemera (Buckman, 1910, Tab.1), at a time when the definition of Bajocian ammonite genera was closer to that of the present day. The recognition that members of the P. rarecostata/acris/subarietis group make up an essential part of the garantiana Zone fauna, as originally defined, confirms that the first appearance of the genus Parkinsonia cannot be taken as defining the base of the parkinsoni Zone (contra Mouterde, 1953; Westermann, 1967; Pavia & Sturani, 1968; Gabilly et al., 1971). Thus the ammonite fauna showing the overlap of S. (Garantiana) and early Parkinsonia; the acris zone (Kumm, 1952) or subzone (Westermann, 1967); is essentially synonymous with much of the garantiana hemera, and must be included in the top of the garantiana Zone (Parsons, 1976b, p.48). Fortunately the definition, and subsequent recognition, of the lithological units, which Buckman (1893), originally assigned to the garantiana hemera is clear cut. Thus it has proved possible (see below), to rectify some of the gaps in our knowledge of the garantiana Zone's ammonite faunas, in its type area. It is evident that Buckman (op. cit.), recognized, from its first inception,

the possibility of subdivision, since he divided the <u>garantiana</u> hemera between two of his labelled horizons ("E" & "F"; e.g. <u>op</u>. <u>cit</u>., p.496, section xi, beds 1 & 2). These horizons coincide with the lower part of the Combe Member (lower 'Rubbly beds' = "E") and the Sherborne Member ('Building-stone' = "F"), which do indeed show some differences in their ammonite faunas (see below). However, this early lead was not followed up until relatively recently (Parsons, 1976b, p.48), and most of the more detailed stratigraphic work, upon which the present subzonal scheme is based, has been undertaken in the thick clay sequences of north Germany.

The first person to recognize, and so name the garantiana Zone in Germany was Mascke (1907), and it is interesting to note that he closely followed Buckman (1893), and drew the base of his 'Parkinsonia-Zone', at the base of the truellei zone. However, many subsequent german authors rejected the use of Buckman's hemerae/zones, and relied on a less formal terminology. Thus initially the garantiana Zone beds were divided between the lower half of the 'Parkinsoni-Schichten' and the upper part of the 'Subfurcaten-Schichten' (Wetzel, 1911; Althoff, 1914). Further subdivision followed, with the recognition between these two of the 'Perisphinctesschichten! (Althoff, 1920, p.4) or 'Bigotiten-Schichten' (Wetzel, 1954, p.549), which was to be named the tetragona zone (Kumm, 1952, p.334) or subzone (Westermann, 1967, tab.1). Other refinements included the renaming of the upper 'Subfurcaten-Schichten', first as the 'Pseudogarantien-schichten' (Bentz, 1928; Althoff, 1928, pp.8-9), and then as the dichotoma Zone (Kumm, 1952, p.334) or subzone (Westermann, 1967, tab.1), whilst the 'untere Parkinsonien-schichten' (Althoff, 1928), was redefined as the acris zone (Kumm, 1952, p.334), or subzone of the parkinsoni zone (Wester-

	Parkinsoni z.	Gar	rantiana Zone	
WESTERMANN 1967	<u>acris</u> subzone	tetracolla subzona	garantiana subzone dichotoma eubzene	
Kirt2M 1952	Acris Zone	('Bigotiten Schichten') Totragona Zono	Dichotoma Zone (para) J	and a subscription of the second s
RENTZ 1928 Althorf 1928	Untere Parkinsonien Schichten'	'Perisphinkten Schichten'	'Pseudogara- ntien Schichten'	
BUNTZ 1924	'Untere Parkinsonien Schichten'	e contraction de la contractio	Subfurcaten Schichten' (?pors)	
ALTHUFF 1920	'Untere Parkinsoni Schichten'	'Perisphincten Schichten'	• Subfurkaten Schichten• (rors)	
WEZEL 1911 ALTHOFF 1914	• Untere Parkinsonien Schichten		'Subfurcaten (Bifurcaten) Schichten' (pars)	
MASCKE 1907		Gerentiane Zone	.	

Table 2. Development of the subdivision of the garantiana Zone beds in Germany.

Sec.

mann, 1967, tab.1). An additional subdivision, the garantiana subzone was erected by Westermann (loc. cit.) between the dichotoma and tetragona subzones, but its relationship to either pre-existing or subsequent schemes is difficult to establish (Hinkelbein, 1975, p.154). This german terminology, the development of which is summarised in Table 2, formed the basis of the subzonal scheme used in the Basse-Alpes of south-east France (Pavia & Sturani, 1968; Pavia, 1973). The main modification consisted of a new subzone, the subgaranti, between the dichotoma and tetragona horizons (Pavia & Sturani, 1968, p.314), which was later renamed the trauthi subzone (Favia, 1973, p.87), because of the previous misidentification of the specimens of 'Hlawiceras'. However, this was an unnecessary change, since S. (G.) subgaranti undoubtedly occurs (contra Pavia, 1973) at this horizon (Althoff, 1928, p.8; Kumm, 1952, p.390; and herein), although it ranges up into the acris subzone (op. cit.). This wide range does not rule out its use (Hedberg, 1976, pp.52, 60-1), and it must have priority over the trauthi subzone. It has been suggested (Sturani, 1971, p.48), that the subgaranti subzone was introduced as a replacement for Westermann's (1967), garantiana subzone. This is difficult to substantiate, since the latter was so poorly defined.

The four subzones adopted by Pavia (1973), including the <u>acris</u> subzone (see above and table 1), can be recognized in the Sherborne area. However, it should be stressed that this subzonal scheme is still very provisional. The lower limits of several subzones are difficult to establish and many important taxa appear to range throughout the bulk, if not all of the Zone (e.g. <u>S</u>. (<u>G</u>.) <u>garantiana</u>, <u>S</u>. (<u>G</u>.) <u>alticostata</u>, <u>S</u>. (<u>G</u>.) <u>depressa</u>, <u>S</u>. (<u>G</u>.) <u>subgaranti</u>, <u>S</u>. (<u>G</u>.) <u>suevica</u>, <u>S</u>. (<u>G</u>.) <u>tetragona</u>; Althoff, 1928, pp.7-9;

Kumm, 1952, pp.390, 430). Some of these problems are discussed in more detail below, and there is little doubt that more work on the ammonite distributions is needed before the subzonal scheme for the <u>garantiana</u> Zone is fully stabilised.

The parkinsoni Zone and its base

Since the original zone (Oppel, 1856) has been considerably restricted, its limits must be determined in relation to the ranges of the subsequently erected, adjacent zones. Its top is defined by the base (see Sturani, 1967, pp.10-1) of the Bathonian zigzag Zone (Oppel, 1865), thus only the lower boundary of the parkinsoni Zone need be discussed here. As detailed above, the original upper limit of the garantiana hemera was drawn at the base of the truellei zone/hemera. This boundary has been implicitly accepted by all British workers (Richardson, 1916, 1932; Arkell, 1933, 1951-9, 1956; Torrens, 1969, etc.), since the bulk of the deposits assigned by Buckman and later workers to the garantiana hemera, and subsequently correlated with the garantiana Zone, are of acris subzone age (= Astarte Bed, south and much of north Dorset; Hadspen Stone of Somerset; Upper-Trigonia-grit of the Cotswolds). If the acris subzone were to be included in the parkinsoni Zone, then the restricted garantiana Zone would be almost unrepresented in southern England, including its type area (Parsons, 1976b, p.48). Thus the truellei zone (Buckman, 1891) is here defined as the basal subzone of the parkinsoni Zone. As originally defined (Buckman, 1891, p.656), this horizon was based on the 'fossil-bed' at Halfway House, yielding "large Parkinsoniae" ... Ammonites more recently collected from this horizon (Torrens, 1969, p.308; Whicher, 1969, p.330; see below), confirm Buckman's (1910, p.55), subsequent assignation of "compressed Parkinsoniae"

(of the "<u>P.-parkinsoni</u> type", <u>op. cit.</u>, p.78), and "<u>Strigoceras</u>" to the <u>truellei</u> hemera. The <u>truellei</u> subzone is thus synonymous with the <u>densicostata</u> subzone (Pavia & Sturani, 1968, p.315), and must have priority. The fact that <u>S. truellei</u> has a wide range beyond that of its subzone, does not discredit its use as an index, as it occurs abundantly in the assemblage-biozone (Holland <u>et al.</u>, p.13), for which it acts as a label.

The horizon between the zigzag and truellei hemera was assigned by Buckman (1907, p.424), to the 'pre-zigzag/post-truellei' hemera or schloenbachi hemera (Buckman, 1908, 1910, tab.1). This horizon was said to be characterised by Parkinsonia "massive forms, with stout, somewhat squared whorls" ... (Buckman, 1910, p.78), which Buckman (1907, p.424), assigned to the 'P. schloenbachi, Schlippe' group. These specimens; a typical representative of which was figured by Buckman (1921 in 1909-30, pl.493); were said by Arkell (1951-9, p.9) to have been misidentified, and were renamed by him as P. bomfordi Arkell (op. cit., pp.149, 157). Thus this horizon has subsequently been known as "horizon a Parkinsonia bomfordi Ark. (= P. schloenbachi S. Buck. non Schlippe)".... (Gabilly, 1964, p.73) or bomfordi subzone (Sturani, 1967). This is an occasion, where there can be little dispute (cf. Dean et al., 1961, p.439), in applying Arkell's rule relating to the priority of zonal indices, which have been misidentified by their original author (Arkell, 1946, p.3, 'rule 4'). Since they are largely identical in all but their corrected name, the bomfordi subzone can take its authorship from Buckman's introduction of the schloenbachi hemera/zone.

The presence of a further subzone, the parkinsoni subzone, between

the <u>truellei</u> and <u>bomfordi</u> subzones has been suggested (Gabilly, 1964, p.73, fig.1; Torrens, 1969, p.302). Whilst evidence for this unit has up to now only been slight, there are some indications (see below) in its support, which will have to be verified by further, more detailed stratigraphic work. Until such work has been undertaken, there is still insufficient evidence to support its use.

DESCRIPTION OF SECTIONS

Since one of the main aims of this work was to undertake a biostratigraphic revision, only those sections which have recently yielded relevant ammonites are described here (see Fig.1), starting at the most westerly and working east.

Bradford Abbas

Although all of the exposures in the Bradford Abbas district described by Buckman (1893, pp.485-6) and Richardson (1932, pp.54-62) are now obscured, a good section can still be made out in the railway-cutting (ST594145). Here above an impersistent, limonitic Astarte Bed (0-0.20m), are seen the Halfway House Bed (0.35-0.70m); with <u>Parkinsonia (Okribites</u>) cf. <u>pseudoparkinsoni</u> Wetzel, CP2881; overlain by more than 2m of the Crackment Limestone Member, with <u>P. (O.) parkinsoni</u> (Buckman <u>non</u> Sow.), CP2994, in the basal 10cm. Rare specimens of <u>P.(?P.) pachypleura</u> Buckman from the upper part of the latter beds indicate the presence of the lower part of the <u>zigzag</u> Zone, although it is difficult to determine its base. Halfway House

Some of the sections in this area have been obliterated, but several.

important exposures are still available for study (Torrens, 1969, pp.308-10, fig.1). Approximately 3.0m of 'Chapel beds' are visible in a cutting adjacent to the old Chapel quarry (Richardson, 1932, p.65), whilst more extensive exposures are available at the new road-cutting (ST602163; Whicher, 1969, p.329) and at the Rock Cottage quarry (ST601165). The latter section (Buckman, 1893, p.487; Richardson, 1932, pp.62-3), which is more weathered and amenable to collecting, shows 0.20m of Astarte Bed, overlain by the Halfway House Bed (0.30m) and +1.5m of 'Chapel beds'. The Halfway House Bed, which has yielded the most ammonites : numerous <u>Parkinsonia (P.) dorsetensis</u> (Wright), topotypes; <u>P. spp.; Prorsisphinctes</u> spp; <u>Cadomites</u> sp. and <u>Strigoceras truellei</u> (d'Orb.), (Torrens, 1969, p.308; Whicher, 1969, p.330): is the type horizon of the <u>truellei</u> subzone.

Louse Hill quarry

At this section (ST609151), which has been previously described by Buckman (1893, pp.488-9) and Richardson (1932, pp.67-70), 0.13m of intermittently conglomeratic Astarte Bed is overlain by +0.20m of brown, sandy, crinoidal limestone (= 'Chapel beds'). Richardson (1932, p.67) followed Buckman (1893, p.488) in correlating the latter bed with the <u>garantiana</u> hemera. However, there is little evidence for this correlation, and it is more likely that they are <u>parkinsoni</u> Zone, 'Chapel beds', similar to those seen in a section further east (ST617165; Whicher, 1969, pp.325-9, bed 7). Here 2.1m of undoubted <u>parkinsoni</u> Zone 'Chapel beds', with <u>Aulacothyris</u> cf. <u>cucullata</u> Buckman, overlay a 0.10m bed, which probably represents a 'condensed' equivalent of both the Astarte and Halfway House Heds (<u>loc</u>. cit.). The Astarte Bed at Louse Hill has yielded an extensive, although



the corro relative lation. An out سو dis ine 1 Soc ribut 1 ولموق . n دغنو 0 ž 0 0 B H the the Sandford amnoni te Lane faunas quarry and their showing

Figur

0

J.

##*`*

rather fragmentary, <u>garantiana</u> Zone, <u>subgaranti</u> subzone ammonite fauna : <u>Strenoceras</u> (<u>Garantiana</u>) <u>platyrryma</u> (Buckman), CP3311-2; topotypes; <u>S.</u> (<u>G.</u>) cf. <u>wetzeli</u> (Trauth), CP3313-4; <u>S.</u> (<u>Pseudogarantiana</u>) <u>minima</u> (Wetzel), CP3316; S.(P.) <u>postrema</u> (Wetzel, 1937, pl.10, fig.11), CP3317; <u>S.</u> (<u>P.</u>) cf. <u>dichotoma</u> (Bentz), CP3315; <u>Spiroceras annulatum</u> (Deshayes), CP3681-2 and <u>S. cylindricum</u> (Baugier & Sauzé), CP3680.

Sandford Lane

There are several exposures on either side of the 'Sandford Lane', Sherborne, of which the most important (ST628178), the Sandford Lane or Combe quarry (Buckman, 1893, pp.492-5; Richardson, 1932, pp.75-7) is found on the west side. A similar section (see Fig.7) is found directly opposite, on the east side, whilst a slightly less well exposed section is found further north (ST629181). The main, or Sandford Lane quarry, is one of the most important Bajocian sections in northern Europe. Apart from the importance of the sauzei/laeviuscula Zone ammonite faunas from the Sandford Lane Bed (= 'fossil-bed'; Parsons, 1974, pp.163-8), this locality is the only one to show a complete section through the Sherborne Limestone Member, as well as an important exposure in the Quarr Lane Beds of the Combe Limestone Member. The distribution of its garantiana Zone ammonite faunas is shown in Figure 3. These specimens were accurately located in relation to a datum level; a thin, marl band (Richardson, 1932, p.76, bed 2); which appears to be laterally persistent and can be traced in most sections (see Fig. 7). All of these records are based on material in the author's collection, apart from the S. (P.) dichotoma, which is based on a specimen in the Buckman collection, BMMM.C80969. This was recorded as coming from the 'Building-stone', and its matrix and preservation is not



Figure An outl ne

\$

the di stribution 05 ۲Ø 0 0 <u>المراجع</u> ts no ammonite of the Blackberry fauna Lane quarry, showing

out of keeping with the Building-stone Beds of the Sherborne Limestone Member.

Blackberry Lane

This section (Richardson, 1932, p.74), is well exposed in a long face on the west side of Quarr and Blackberry Lanes. The present section (Fig.4), was measured at the southern end (ST635174), where the top of the Redhole Lane Beds is visible. These beds disappear at some point between here and Sandford Lane. At the latter locality they are represented by the basal part of the Quarr Lane Beds, which are slightly better bedded than the bulk of this deposit.

Redhole Lane

This large quarry, which is now a caravan park, has been previously described by Buckman (1893, p.496) and Richardson (1932, p.73). The lower part of the main quarry has now been filled in, so that a large part of the Building-stone Beds is obscured. The ammonites from the Quarr Lane Beds recorded in Figure 5, have mainly come from the long, low faces found to the south of the main quarry. A specimen from this quarry, cited as '<u>Parkinsonia praecursor</u>' by Buckman (1893, p.496), BMNH.C9209, is close to <u>S</u>. (<u>G</u>.) <u>garantiana</u> (Douvillé, 1916, pl.1, fig.1 <u>non</u> d'Orb). Ammonites similar to this specimen have been recorded in Figures 3-5 as <u>S</u>. (<u>G</u>.) sp. A. The specimens recorded in Figure 5 as <u>S</u>. (<u>G</u>.) sp. B are fine ribbed variants of <u>S</u>. (<u>G</u>.) <u>trauthi</u> (Bentz), [<u>op. cit.</u>, cf. pl.2, fig.8; =? <u>S</u>. (<u>G</u>.) <u>trauthi</u> (Pavia, 1973, pl.19, fig.3 ?<u>non</u> Bentz)].

Oborne

There are several important sections in this district, including



Frogden quarry (ST642185; Buckman, 1893, p.500; Richardson, 1932, pp.78-9; Parsons, 1976a, pp.125-7) and the Oborne Lane section and quarry (ST656186; Buckman, 1893, p.502; Richardson, 1932, p.81). The only relevant ammonite, which has come from the lower part of the Acanthothiris Beds at a temporary section at Oborne Wood (ST648188; Parsons, 1976a, pp.128-133, bed 8b), is <u>Cadomites</u> (<u>C</u>.) cf. <u>deslongchampsi</u> (d'Orb.), CP2812. The <u>Teloceras</u> sp. recorded from the basal part of the Sherborne Limestone Member at the Oborne Lane section (<u>op. cit.</u>, p.133) has proved to be derived. This has been confirmed by a specimen of <u>Stephanoceras</u> (<u>S</u>.) sp. recently collected from the same horizon, which has a strongly 'iron-shot' matrix in its camerae.

Milborne Wick

East of Oborne, the Stroud Formation is poorly exposed. The conglomeratic base of the Sherborne Limestone Member is visible at the Milborne Wick Lane section (ST663205; Buckman, 1893, pp.502-3; Richardson, 1916, pp.516-7; Osborne-White, 1923, p.13; Parsons, 1976a, pp.133-5), and a similar horizon must have been exposed nearby, at the site of the Poyntington reservoir excavations (Kellaway & Wilson, 1941, p.154). These latter beds were originally described (<u>loc. cit.</u>) as being a sandy limestone facies of the <u>subfurcatum</u> Zone. However, the specimens variously recorded as <u>Strenoceras niortensis</u> (D'Orbigny)', (<u>loc. cit.</u>), or <u>'S</u>. <u>bajocense</u> (Defrance)', (Wilson <u>et al.</u>, 1959, p.75), have been misidentified, and prove to be <u>S</u>. (<u>Pseudogarantiana</u>) cf. <u>dichotoma</u>, IGS.GK311 & 313. There is no evidence for the <u>subfurcatum</u> Zone in this immediate area.

The most important exposure in the district is at Barrow Hill (ST672210), and although briefly mentioned in previous accounts (Woodward,

?tetragona ? acris	SUBZONE
	Bigotites aff. admirandus B. sp.nov. A Parkinsonia rarecostata P. subarietis P. pseudoparkinsoni Spiroceras sp. Orthogarantiana inflatocoronata S.(Garantiana) cf. suevica S.(G.) cf. alticostata S.(Pseudogarantiana) postrema S.(P.) cf. minima S.(P.) sp. cf. sp. nov. B
?Redhole Quarr Lane Beds	

relative An outline ine section of distribution 05 of its ammonito the Barrow Hill fauna. quarry, showing

the

Figure

<u>б</u>

1894, p.80; Richardson, 1916, p.511; Osborne-White, 1923, p.16), it has never been described in detail. Here an old, shallow quarry, shows a long, rather slipped face in predominantly rubbly limestones (= Quarr Lane Beds). At the extreme western end, below an impersistent marl band, the rocks are more evenly bedded, and probably represent the top of the Redhole Lane Beds. Because much of the section has no obvious datum level, some of the ammonite records shown in Figure 6, are probably slightly less accurately localized than is the case elsewhere. A specimen in the Buckman collection, BMNH.C80967, from Barrow Hill; a complete, small macroconch, with mouth-border, akin to S. (G.) suevica (Wetzel), (cf. Quenstedt, 1886, pl.71, fig.18); probably came from a similar horizon to that of the related specimen recorded in Figure 6. The specimens of S. (Pseudogarantiana) sp. nov.B, recorded both in the latter and Figure 5, are of an undescribed, relatively large, inflated microconch species, which is found in the acris subzone, Astarte Bed of south Dorset (Parsons, 1975a, p.9). The specimens recorded here as P. subarietis are probable microconchs (cf. Wetzel, 1911, pl.14, figs.3-4, pl.15, fig.1), whilst the specimen from Sandford Lane (Fig. 3), is larger and is an undoubted small macroconch (op. cit., cf. pl.14, fig.6). The specimens recorded as Bigotites sp. nov. A. in Figures 3-4 & 6, represent a problematic group, of medium sized, relatively compressed macroconchs. They have strong constrictions, well marked interdigitation and interruption of the ribs on the venter, no tubercles and only rounded, relatively weak ribs on the internal mould. Unfortunately the state of preservation in most cases is not good, but enough details are shown to suggest that these ammonites form an intermediate link between Bigotites and Parkinsonia, particularly P. rarecostata Buckman.

The only other relevant section in this area, is the old roadcutting, near the now disused Milborne Port railway station (ST675209). Although rather poorly exposed and overgrown, a similar section to that recorded by Richardson (1916, pp.510-1) can still be made out. The lower half consists of rubbly limestones, probably equivalent to the top of the Quarr Lane Beds, whilst the upper 1.5m is more evenly bedded, massive, oolitic and has yielded a specimen of <u>Parkinsonia</u> (<u>P.</u>) sp. from the top 10cm. They are thus parkinsoni Zone Goathill Beds.

Goathill

There have been several sections in this area of the Crackment Limestone Member (Torrens, 1969, pp.318 & 321), but they are not now well exposed. A section in a silage pit (ST674172), on the line of the Poyntington Fault, showed on the east side c.3m of Crackment Limestones, whilst on the west wide 2.3m of Goathill Beds were preserved (op. cit., pp.317-8). A section in the Goathill Beds, almost identical to the latter, can be seen just to the south, in the bank next to the path (ST674171). The ammonites from both these sections [Parkinsonia (P.) cf. subplanulata Wetzel (cf. Sturani, 1967, pl.8, fig.2), CP3489; P. (O.) cf. parkinsoni Buckman non Sow., HT4881; P. (Gonolkites) sp., HT4737], suggest a correlation with the parkinsoni Zone and possibly the bomfordi subzone.

Temporary sections in Sherborne

Castle View

Large areas of shallow, highly disturbed exposures in the Quarr Lane Beds were (July 1968 - Feb.1969) visible during the excavation of foundations for this housing estate (ST646172). Ammonites were abundant and included : S. (G.) platyrryma, CP2093; S. (G.) cf. inflatocoronata (Wetzel), CP1181; <u>S.</u> (<u>P.</u>) <u>minima</u> (Wetzel), CP1185; <u>S.</u> (<u>P.</u>) cf. <u>dichotoma</u>, CP1183 (specimen with square cross-section, similar to those recorded in Figure 5); <u>S.</u> (<u>?P.</u>) <u>pompeckji</u> (Wetzel), CP2096; <u>S.</u> (<u>P.</u>) sp. nov. A., CP2095 (a small undescribed, fine ribbed microconch, with a round whorl cross-section, see Figures 3 & 5); <u>Orthogarantiana inflata</u> Bentz, CP1184; <u>O. cf. uncinata</u> (Qu.), CP1186; <u>Parkinsonia</u> cf. <u>gracilis</u> Wetzel (1937, pl.12, fig.4), CP2094; <u>Prorsisphinctes pseudomartinsi</u> (Siemiradzki), CP1179; <u>Bigotites</u> cf. <u>tuberculatus</u> (Nicolesco), CP1182; <u>Sphaeroceras</u> cf. <u>globosum</u> Buckman, EMNH.C80371; <u>Cadomites</u> (<u>C.</u>) sp., CP1180, <u>Oppelia</u> (<u>O</u>.) cf. <u>flexa</u> (Buckman), CP1178. This assemblage, of <u>garantiana</u> Zone, and probably <u>subgaranti- tetragona</u> subzone age, is important because of its high diversity : it contains forms not found <u>in situ</u>.

Newell

A relatively small section was produced (March, 1978) during the excavation of an access road and drains for a new house overlooking Newell, Sherborne (ST634170). The basal exposure in +1.20m of thin bedded limestones : with <u>Parkinsonia</u> (P.) <u>depressa</u> (Qu.), inner whorls cf. Pavia, (1973, pl.23, fig.3), CP3335, (a similar specimen, IGS.GSM62717 has come from a temporary section at Lenthay); <u>P. (P.) cf. homfordi</u>, CP3488; <u>P.</u> (<u>Gonolkites</u>) sp., CP3337 and <u>Prorsisphinctes</u> sp., CP3336: was overlain by a massive bed 0.50m thick. These beds, which are probably on the same horizon as those exposed at Goathill (see Fig.7), are of <u>bomfordi</u> subzone age. Above came 2.10m of thin bedded limestones with marls, capped by 0.80m of thin limestones interbedded with clay bands. The latter undoubtedly represent a lower part of the Crackment Limestone Member. A similar section in the Goathill Beds, yielding abundant Parkinsonia, was
HALFWAY NEWELL SANDFORD BLACK- REDHOLE OBORNE OBORNE MILBORNE GOAT HILL (11) WOOD(7) LANE WICK LANE BERRY LANE HOUSE (5) (8) LANE(9) (2)(4) LANE & GOAT HILL FROGDEN (5a) & BEDS TITIT BARROW (6) TIT 0 HILL (10) WEST EAST 0 MB $\sim \sim \sim$ ~~~ m Thursday Pourse ~~~ $\sim \sim$ • ^ * LIMESTONE $\sim \sim \sim$ (10) $\sim n$ \sim ~~~ ~~~ QUARR ~~~ ~ h a~~~ \sim CHAPEL nen \sim \sim 5 LANE first Prarecostata-nm \sim ~~ در م $\sim \sim$ $\sim \sim$ BEDS $\langle \cdot, \cdot, \cdot \rangle$ Mbr $\langle \ddots \rangle$ \sim $\sim \sim$ ~~~ 35. V 1 3~~~ fra · M 5 S_{n} ∞ ~~~ _____ \sim $\sim \sim$ him ASTARTE BED ----***** ~ SHERBORN \sim REDHOLE $\sim \sim$ LANE $\sim \sim$ ŝ BEDS $\sim \sim \sim$ $\sum_{n} \sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{i=1}^{n} \sum_{j=1}^{n$ \sim m BUILDING STONE Z BEDS TITI Jana E in ALC: III ATTAL ONE 1111 (9)2m **alauk**i ACANTHO-CT 3637. BEDS

Figuro >

narkinsoni 100 relation 05 the

Zone main

rocks 3 11 the Shorb exposures õ 1713 ō

of Ó Ö

carantiana

and

probably seen at a temporary section at Sheeplands, Sherborne (Kellaway & Wilson, 1941, p.156).

DISCUSSION OF AMMONITE FAUNAS

garantiana Zone

dichotoma subzone

The dichotoma subzone is based on the first appearance and acme of S. (P.) dichotoma (Pavia & Sturani, 1968, p.314). However, all the in situ specimens allied to this species have in the Sherborne area come from relatively high horizons (see Fig.5), although specimens very close to its type (Bentz, 1928, pl.19, figs.2a-b), but unfortunately less well localized, are known. These ammonites from Sandford Lane (BMUH.C80969) and Poyntington reservoir (IGS.GK311 & 313, see above), from both their matrix and location, undoubtedly come from the lower part of the Sherborne Limestone Member. The previous confusion of the latter specimens with members of S. (Strenoceras) s. str. (e.g. Kellaway & Wilson, 1941), only tends to confirm both their stratigraphic position at the base of the garantiana Zone (cf. Althoff, 1928; Kumm, 1952) and their phylogenetic position (Wetzel, 1954, fig.7), as the earliest members of S. (Pseudogarantiana), and transitional to S. (Strenoceras). It thus seems reasonable to correlate the lower part of the Sherborne Limestone Member (Acanthothiris & Building-stone Beds) with the dichotoma subzone, although the scarcity of ammonites prevents any precise definition of its limits.

subgaranti subzone

The first abundant <u>garantiana</u> Zone ammonites occur at the base of the Quarr Lane Beds at Sandford Lane, and in their partial lateral equivalent,

建筑中国合同市场建筑的

the Redhole Lane Beds, at Redhole Lane quarry (see Figs.3 & 5). The most significant members of this fauna, <u>S</u>. (<u>G</u>.) <u>subgaranti</u> and <u>S</u>. (<u>G</u>.) <u>trauthi</u>, belong to a group previously included in <u>G</u>. (<u>Subgarantiana</u>) Bentz 1928, (= <u>Hlawiceras</u> Buckman 1921). However, the separation of <u>Subgarantiana</u> from <u>Garantiana s</u>. <u>str</u>. on both morphological and stratigraphical grounds (cf. Wetzel, 1954, fig.7) appears artificial, and the two are closely associated throughout much of their vertical anges. Separate subgeneric status for two such similar macroconchs would thus seem unnecessary.

The original definition of the subgaranti subzone (Pavia & Sturani, 1968, p.314) was based on the first appearance and abundant occurrence of S. (G.) subgaranti. The associated ammonite fauna from the Basse Alpes is less diagnostic, as it has a wider range, although subsequent additions to the typical fauna include S. (G.) platyrryma (Gabilly et al., 1971, p.88). Unfortunately the specimens of S. (G.) subgaranti from the Basse Alpes (Pavia & Sturani, 1968), were later re-identified as S. (G.) trauthi, with a consequent change in subzonal index (Pavia, 1973). However, if the specimen of the latter taxon figured by Pavia (1973, pl.19, fig.3) is typical of the rest of his material, then there must, even now be doubts as to the correct identification of these specimens. The type of S. (G.) trauthi (Bentz, 1928, pl.18, fig.2), is fairly coarse ribbed, and on the evidence of topotype material from Louse Hill, it is possibly conspecific with S. (G.) platyrryma (Buckman, 1921). Pavia's (loc. cit.) specimen is finer ribbed, rather like a specimen figured by Douville (1916, pl.2, fig.8), and is very similar to ammonites recorded here (Fig.5), from an horizon probably to be correlated with the tetragona subzone. Taking into

account these uncertainties in identification, the basal boundary of this subzone is best defined by the first appearance of any 'Subgarantiana' : i.e. members of the S. (G.) platyrryma/subgaranti/trauthi group: rather than by that of any one individual species. S. (G.) subgaranti can be retained as index species, since it undoubtedly occurs at this horizon, in Dorset (Fig. 3), in the upper half of the 'dichotoma-schichten' (= garantiana subzone, Westermann, 1966; subgaranti subzone, Sturani, 1971), in north Germany (Althoff, 1928; Kumm, 1952; Wetzel, 1954) and in Spain (Hinkelbien, 1975, figs.11-12). It is difficult to establish if the first appearance of S. (G.) subgaranti/trauthi in the Sherborne area, as recorded here (Figs. 3 & 5), in fact marks their absolute range. Unfortunately their appearance coincides with that of the general ammonite fauna, which may mark a switch to an ammonite rich biofacies, rather than a biostratigraphic event. A more accurate boundary between the dichotoma and subgaranti subzones, if it can be established at some future date with the aid of the rare ammonites to be found at this horizon, is likely to be in an upper part of the Building-stone Beds. However, it must be stressed that there is a possibility, based on their general morphology and some stratigraphic evidence (Hinkelbein, 1975, fig.12), that the S. (G.) platyrryma/trauthi group represent the macroconch counterpart to the microconch S. (P.) dichotoma. If this were to be confirmed at other exposures, it would totally destroy the basis for the separation of the subgaranti and dichotoma subzones.

tetragona subzone

The base of this subzone is taken here at the marl band (Richardson, 1932, p.76, bed 2), which appears to be laterally persistent, and is

probably approximately isochronous over the short distances involved (see Fig.7). At Sandford Lane (Fig.3), its base is marked by the first appearance and abundant occurrence of large Perisphinctids : Prorsisphinctes and Bigotites (B.sp.A), which seem to be transitional to Parkinsonia. Although it is possible that we are again dealing with a change in biofacies, since the large Perisphinctids are noticeably absent from earlier beds, their appearance does seem to be a biostratigraphic event, which can be traced in both north Germany (= 'Bigotiten/Perisphinctes-schichten'; Althoff, 1928, etc.) and France (Gabilly et al. 1971). However, it should be noted that there are areas where the larger Perisphinctids are absent (e.g. east Spain, Hinkelbein, 1975) or rare (e.g. south-east France, Pavia, 1973). In the latter area (op. cit., tab.2-3), the limits of the tetragona subzone were determined by the absolute range of specimens allied to S. (G.) tetragona (Wetzel). This would appear to be very unreliable, as this species has been recorded in both the acris subzone (Althoff, 1928; Kumm, 1952, p.430), and at an horizon now to be correlated with the dichotoma/subgaranti subzones (Althoff, 1928, p.8). In north Dorset there are less problems in its recognition, since an influx of large Perisphinctids usually mark its base (see Fig.3), whilst specimens closely allied to the subzonal index only occur towards the top of the subzone (see Fig.5). It should be noted that many of the taxa recorded in the latter figure probably have a wide range down into the subgaranti subzone (e.g. S. (G.) subgaranti itself, as in Fig.3). The sharp cut-off at the probable subgaranti/tetragona subzone boundary is at least in part due to collection failure, as ammonites are much rarer in the Redhole Lane Beds.

acris subzone

The base of the acris subzone has been defined by the first appear-

ance of Parkinsonia s. lat. (Kumm, 1952, p.430; Pavia & Sturani, 1968, p.314), and this is relatively clear cut in most Dorset sections (see Figs. 3, 5-6). One area of uncertainty has come to light in the study of the Barrow Hill section (Fig.6). It is evident that many of the microconch Bigotites are very similar to Parkinsonia, and have stronger, sharper ribbing, with less well marked constrictions (cf. Pavia, 1973, pl.29, fig.3), than their macroconch partners. Taking into account that macroconch Bigotites have been found ('B.sp.A'), which are transitional to P. (Parkinsonia), it is logical to expect their microconch partners, to be even closer to, and virtually indistinguishable from, Parkinsonia s. lat. Such indeed, does appear to be the case. The specimens of P. (Okribites) subarietis Wetzel and P. (O.) pseudoparkinsoni Wetzel recorded in Figure 6, are microconchs which appear some distance below the first macroconch Parkinsonia s. str., P. (P.) rarecostata. It is possible that these are the microconch. partners of macroconchs which would be best assigned to Bigotites, rather than Parkinsonia (P.). If it should be confirmed from other sections, that the microconch P. (Okribites) makes its appearance before the macroconch P. (Parkinsonia), then in order to prevent any possible confusion with the earlier Bigotites microconchs, the base of the acris subzone should be redrawn below the first macroconch Parkinsonia s. str. Only in the case of Figure 6 would any future change be required. Here the boundary would have to be moved up to below the first P. (P.) rarecostata. Unfortunately the upper limits of this subzone cannot be discussed, as no fossiliferous exposures of an unbroken contact with the truellei subzone have been observed.

<u>parkinsoni</u> Zone

truellei subzone

This subzone, which is defined by the first appearance of the macro-

conch, P. (Parkinsonia) parkinsoni (J.Sow.), [? = P. (P.) dorsetensis(Wright)], is well represented by the condensed Halfway House Bed. However, I have not been able to recognize it in the more expanded facies of the Sherborne district, nor establish its contact with the underlying <u>garantiana</u> Zone. This boundary must fall within the upper part of the Quarr Lane Beds, where the previous records of 'P. <u>dorsetensis</u> and <u>Strigoceras truellei</u>' (Buckman, 1893, p.496), suggest that the <u>truellei</u> subzone fauna can only be a short distance above the last <u>in situ</u> elements of the top of the precipitous faces at Sandford Lane and Redhole Lane quarries, from which it has proved impossible to collect any <u>in situ</u> specimens. Future work, possibly with the aid of extending ladders, may solve this problem.

There is a level at the top of the <u>acris</u> subzone, where the last <u>S</u>. (<u>Garantiana</u>) and <u>S</u>. (<u>Pseudogarantiana</u>) have died out, and only <u>Parkinsonia</u> are common (e.g. <u>P</u>. <u>rarecostata</u>, <u>P</u>. <u>subarietis</u>). This horizon, (? = "horizon a <u>Parkinsonia orbignyana</u> Wetz."; Contini, 1970, p.154; Gabilly <u>et al.</u>, 1971, p.88), is poorly defined, and must be included in the top of the <u>garantiana</u> Zone, since it falls below the base of the <u>truellei</u> subzone, here defined as the lowest unit of the <u>parkinsoni</u> Zone.

bomfordi subzone

The two exposures in the Sherborne area yielding Parkinsonid ammonites, described here, are stratigraphically isolated, and their contact with adjacent faunas cannot as yet be established. These faunas, with inflated macroconch Parkinsonids (e.g. P. (P.) <u>bomfordi</u>, P. (<u>Gonolkites</u>) spp.), are very similar to those from the middle of the Burton Limestone in south

Dorset (Parsons, 1975a), and are thus of <u>bomfordi</u> subzone age. The topmost fauna of the latter, characterised by <u>Planisphinctes</u> (Sturani, 1967, pl.1), which has been found 10-20cm below the <u>zigzag</u> Bed in south Dorset (Stony Head & Burton Bradstock, pers. obs.), has not been observed in north Dorset. However, it is possibly the same age as the <u>Decotraustes</u> <u>nodifer</u> Fauna' (Torrens, 1974, p.584), recorded from the Crackment Limestone Member at Bradford Abbas (Arkell, 1951-9, p.10), and now correlated with the top of the <u>parkinsoni</u> Zone.

Similarly little can be deduced as to the nature of the contact between the <u>bomfordi</u> and <u>truellei</u> subzones in the Sherborne area, although the record of <u>P</u>. (<u>Okribites</u>) <u>parkinsoni</u> (Buck. <u>non</u> Sow.), from the base of the Crackment Limestone at Bradford Abbas railway cutting (see above) is perhaps significant. This 'species' is characteristic of the lower part of the <u>bomfordi</u> subzone in south Dorset (pers. obs.) and in the mid and north Cotswolds (Parsons, 1976b, p.55). There is thus some evidence for the recognition of a '<u>parkinsoni</u> subzone' (see above), although it would need confirmation from more detailed work on an ammonite rich sequence, such as the Burton Limestone of south Dorset.

CORRELATION WITH OTHER AREAS

Because of the discontinuous nature of the Sherborne exposures in the <u>parkinsoni</u> Zone, only the <u>garantiana</u> Zone will be dealt with here in any detail.

Southern England

Upper Bajocian rocks are found between the south coast at Burton Bradstock, and Hook Norton in Oxfordshire. North of this point, with the

exception of the poor ?garantiana/?subfurcatum Zone fauna from the Inner Hebrides (Morton, 1971; Parsons, 1975b, p.192), no upper Bajocian ammonites have been recovered.

The present study has done little to improve the correlation of the parkinsoni Zone beds; such as the Cotswold Clypeus-grit Member (Parsons, 1976b, p.55), the Dundry Freestone and 'Coralline beds' of Dundry Hill, Bristol (Parsons, 1979b, p.149), the Doulting beds of the Mendips (Parsons, 1975b, pp.203-4) and the Burton Limestone of south Dorset (Parsons, 1975a, p.12) and south-west Somerset (Parsons & Torrens in Cope et al., 1969, p.A26); which are broadly equivalent of the lower part of the Crackment Limestone, the Goat Hill Beds and an upper part of the Quarr Lane Beds. On the other hand, the finer subdivisions of the garantiana Zone, which have been recognized here, have confirmed the wide extent of the stratigraphic hiatus, which is present over much of southern England (Parsons, 1976b, p.48). Thus the Cotswold Upper-Trigonia-grit Member (Parsons, 1976b, pp.52-5), the Dundry Maes Knoll Conglomerate (Parsons, 1979b, p.149) and the Astarte Bed of south Dorset (Senior et al., 1970) and south-west Somerset (Seavington St. Mary; Parsons & Torrens, in Cope et al., 1969, p.A26), have all yielded a combination of S. (Garantiana) and Parkinsonia, mainly P. rarecostata. They are thus garantiana Zone, acris subzone in age, as with much of the Quarr Lane Beds of the Combe Limestone Member. A similar age is indicated for the 'Hadspen Stone' (Richardson, 1916, p.486) of Somerset, a relatively thick deposit of ferruginous, intraclastic limestone. Here at the Horsecombe Bottom (ST656315) and Limekiln quarries (ST655314), (Richardson, 1916, pp.504-6) an acris subzone fauna : of Parkinsonia pseudoparkinsoni, CP1128; S. (Garantiana) aff. garantiana,

CP3266; <u>S.</u> (<u>Pseudogarantiana</u>) <u>minima</u>, CP3281 and <u>Prorsisphinctes</u> sp., CP3308: has been found, thus confirming Richardson's (<u>op. cit.</u>, pp.505, 508-9), records of <u>Parkinsonia</u> of the <u>P. rarecostata</u> group. It is thus evident that the major Upper Bajocian unconformity, present across much of southern England (= 'Vesulian Transgression'), is both isochronous and of immediately pre-<u>acris</u> subzone age. Similarly, only a small area around Sherborne now preserves any beds of early <u>garantiana</u> Zone age at the surface, since elsewhere they are absent, due to subsequent erosion and/or non-deposition. There are however, indications that more extensive and complete <u>garantiana</u> Zone sequences are preserved at the subsurface elsewhere in Dorset.

Europe

It is only recently that there has been a revival of interest in the detailed stratigraphy of the Upper Bajocian (Favia & Sturani, 1968; Pavia, 1973). Unfortunately in the latter works, there was a great reliance on the earlier german studies (Bentz, 1928; Westermann, 1957), since the sequence between the <u>subfurcatum</u> Zone and the <u>bomfordi</u> subzone in the Basse Alpes has a rather sparse and often poorly preserved ammonite fauna. The pre-war work on the north German <u>garantiana</u> beds, summarized in Kumm (1952) and Westermann (1967), suffered from relatively poor localization of material, which was increasingly split into an excessive number of 'species' (Bentz, 1928; Wetzel, 1954). These faunas are in urgent need of a modern revision, and as consequence the long species lists (Althoff, 1928; Kumm, 1952) are very difficult to interpret. Taking this into account, together with the fact that the faunas of the <u>subgaranti</u> and <u>tetrapona</u> subzones are rarely separated, it is not surprising that only a

two-fold subdivision of the <u>garantiana</u> Zone has been commonly attempted. These divisions, which have rested on the first appearance of <u>Parkinsonia</u> (= <u>acris</u> subzone), underlain by a fauna dominated by <u>Pseudogarantiana</u> (= <u>dichotoma</u>, <u>subgaranti</u> & ? <u>tetragona</u> subzones), have been widely recognized throughout Europe, and indeed were accepted by the present author (Parsons, 1975b, 1976b, 1979b), prior to this study. Areas where these subdivisions can, or have been recognized include; Württemberg, south Germany (Dietl, Flaig & Gluck, 1978), north-east of the Massife Central (Mouterde, 1953) and the Jura, France (Contini, 1970), north-east Spain (Westermann, 1955), south-east Spain (Hinkelbein, 1975) and northern Italy (Sturani, 1964, 1971).

STRATIGRAPHIC CONCLUSIONS

1. Following both original definitions and subsequent British usage, the <u>garantiana/parkinsoni</u> zonal boundary is here drawn below the <u>truellei</u> subzone. The <u>acris</u> subzone must thus be included in the <u>garantiana</u> Zone.

2. All the currently accepted subzones of the <u>garantiana</u> Zone (<u>dichotoma, subgaranti, tetragona & acris</u>), have been recognized in its type area, the Sherborne district.

3. As it provides the only complete section through the <u>garantiana</u> Zone beds in the district, Sandford Lane quarry (ST629181), is here desigmated as the type locality for this Zone (see Figure 3).

4. Only two of the subzones of the <u>garantiana</u> Zone, the <u>acris</u> and <u>dichotoma</u> subzones, are both relatively well defined and widely recognized in other areas of Europe. It is possible that the <u>subgaranti</u> (= <u>trauthi</u>

subzone, Pavia, 1973) and <u>tetragona</u> subzones, will not survive any future, detailed revision based on complete, ammonite rich sections.

5. Although a continuous sequence through the <u>parkinsoni</u> Zone is probably present in the Combe Limestone Member of the Sherborne area, only the <u>bomfordi</u> subzone has been recognized in this work.

6. It is possible that an additional subzone (= the 'parkinsoni' subzone) can be recognized between the <u>truellei</u> and <u>bomfordi</u> subzones, but this must be confirmed by future, more detailed work, probably in the south Dorset area.

7. The "Upper Bajocian Unconformity" (= site of "Vesulian transgression") is demonstrably isochronous (pre-acris subzone age), and only in the Sherborne district are any rocks of early <u>garantiana</u> Zone age preserved at the surface.

8. The area containing the rocks of <u>dichotoma-tetragona</u> subzone age is small and very sharply defined. The lateral changes in thickness and facies are very abrupt (see Figure 7), and some penecontemporaneous structural control must be postulated (? = fault bounded 'graben').

ACKNOWLEDGEMENTS AND LOCATION OF MATERIAL

The bulk of the material recorded here is in the author's collection (CP.), whilst other specimens are located in the British Museum (Nat. Hist.), London (EMNH.) and the Institute of Geological Sciences, London (IGS.). I should like to thank the curators of these museums for access to, and the loan of material in their care. I should also like to thank Dr. H.S. Torrens, University of Keele, for his help during the course of my work, and for the loan of specimens in his collection (HT.).

REFERENCES

ALTHOFF, W. 1914. Die geologischen Aufschlüsse Bielefelds. <u>Bericht</u> . <u>Naturwissens. ver. Bielfeld</u> . 3, 195-225.	
,1920. Über die Coronatenschichten-Aufschlüsse der Jahre 1913-1918 in Grube I im Tal von Bethel bei Bielefeld. Bericht. Naturwissens. ver. Bielfeld. 4, 3-10.	
,1928. Übersicht über die Gliederung dermesozoischen Schichten bei Bielfeld. <u>Bericht. Naturwissens. ver. Bielfeld</u> . 5, 1-20.	
ARKELL, W.J. 1933. The Jurassic System in Great Britain. Oxford Univer- sity Press. xii + 681pp.,41 pls.	•
, 1946. Standard of the European Jurassic. <u>Bull. geol. Soc.</u> <u>Amer. 57, 1-34.</u>	
, 1956. <u>Jurassic Geology of the World</u> . Oliver and Boyd, Edinburgh, xiv + 806pp., 46 pls.	
BENTZ, A. 1924. Die Garantienschichten von Norddeutschland mit besondere Berucksichtigung des Brauneisen-oolithhorizontes von Harzburg. Jb. Preuss. Geol. Landesanst. 45, 119-193, pls. 4-9.). S
ELAKE, J.F. 1880. On the Portland rocks of England. Q.Jl.geol.Soc.Lond. 36, 189-236.	
BUCKMAN, S.S. 1881. A descriptive catalogue of some species of ammonites from the Inferior Colite of Dorset. Q. Jl. geol. Soc. Lond. 37 588-608.	3 7,

, 1889. On the Cotteswold, Midford and Yeovil Sands, and the division between Lias and Colite. Q. Jl. geol. Soc. Lond. 45, 440-474.

i i se depens

la programa

BUCKMAN, S.S. 1891. On certain Ammonite-zones of Dorset and Somerset, <u>Rep. Brit. Assoc. Adv. Sci.</u> (1891), C,5, 655-6.

> <u>, 1893.</u> The Bajocian of the Sherborne district; its relation to subjacent and superjacent strata. <u>Q. Jl. geol. Soc.</u> Lond. 49, 479-522.

, 1894. "The top of the Inferior Oolite" and a correlation of "Inferior Oolite" deposits. <u>Proc. Dorset Nat. Hist. Antiq.</u> <u>F.C. 14, 37-43.</u>

, 1907. On the correlation of the Bath-Doulting Strata with those of Dorset. Q. Jl. geol. Soc. Lond. 63, 424-6.

<u>Ammonites</u>, London and Thame, Vols. 1-7, 790 pls.

, 1910. Certain Jurassic (Lias-Oolite) Strata of South Dorset; and their Correlation. <u>Q. Jl. geol. Soc. Lond.</u> 66, 52-89.

- CONTINI, D. 1970. L'Aalenian et le Bajocien du Jura franc-comtois. Ann. Sci. Univ. Besançon. (3), Géologie, fasc. 11, 204pp.
 - COPE, J.C.W. et al. 1969 International Field Symposium on the British Jurassic, Excursion 1, Guide for Dorset and South Somerset (Editor H.S. Torrens), University of Keele, 71pp.
 - DEAN, W.T., DONOVAN, D.T. and HOWARTH, M.K. 1961. The Liassic Ammonite Zones and Subzones of the North-West European Province. <u>Bull.</u> <u>Brit. Mus. (N.H.), Geology</u>, 4, 435-505, pls. 63-75.
 - DIETL, G., FLAIG, R. and GLUCK, E. 1978. Zur Stratigraphie des Ober-Bajocium (Braunjura *f/E*-Grenzschichten) am Plettenberg bei Balingen, Württemberg. <u>Stuttgarter Beitr. Naturk.</u>, ser. B, 40, 16pp.
- DONOVAN, D.T. and HEMINGWAY, J.E. 1963. Lexique Stratigraphique International ; Fascicule 3a, Part 3ax, Jurassic. Paris, 394pp.
- DOUVILLE, R. 1916. Études sur Les Cosmocératidés. <u>Mém. Serv. Explic.</u> <u>Carte. géol. dét. France</u>, 75pp., 24pls.

FOWLER, J. 1957. The Geology of the Thornford Pipe-Trench. Proc. Dorset Nat. Hist. Arch. Soc. 78, 51-7.

- GABILLY, J. 1964. Le Jurassique Inférieur et Moyen sur le littoral Vendéen. <u>Trav. Inst. Géol. Anthropol. Préhist. Fac. Sci.</u> <u>Poitiers</u>, 5, 65-107.
 - du Jurassique en France : Bajocien. C.R.S. Soc. géol. Fr., 85-8.
- GATRALL, M., JENKYNS, H.C. and PARSONS, C.F. 1972. Limonitic concretions from the European Jurassic, with particular reference to the "Snuff-boxes" of Southern England. Sedimentology, 18, 79-103.
- HEDBERG, H.D. 1976. International Stratigraphic Guide: A guide to stratigraphic classification, terminology and procedure. Wiley, New York, xvii + 200pp.
- HINKELBEIN, K. 1975. Beiträge zur Stratigraphie und Paläontologie des Juras von Ostspanien. VIII. Stratigraphie und Fazies im Mittel jura der zentralen Iberischen Ketten. <u>N. Jb. Geol. Paläont. Abh.</u> 148, 139-184.
- HOLLAND, C.H. et al. 1978. A guide to stratigraphical procedure. <u>Geol.</u> <u>Soc. Lond. Spec. Rep. No.</u>10, 18pp.
- HUDLESTON, W.H. 1886. Excursion to Sherborne and Bridport. Proc. Geol. Ass. 9, 187-199.
- KELLAWAY, G.A. and WILSON, V. 1941. An outline of the geology of Yeovil, Sherborne and Sparkford Vale. <u>Proc. Geol. Ass</u>. 52, 131-174, pls. 8-10.
- KUMM, A. 1952. Das Mesozoikum in Niedersachsen : Der Dogger. <u>Geol. u.</u> Lagerstätten Niedersachsens. 2 (2), 329-509.
- MASCKE, E. 1907. Die Stephanoceras-Verwandten in den Coronatenschichten von Norddeutschland. Inaug. Dissert. Univ. Göttingen. 38pp.
- MORTON, N. 1971. Some Bajocian ammonites from western Scotland. Palaeontology. 14, 266-293, pls.40-51.
- MOUTERDE, R. 1953. Études sur le Lias et le Bajocien des bordures nord et nord-est du Massif Central Français. <u>Bull. Serv. Carte Géol.</u> <u>Fr.</u> 50, No.236, 63-521, pls.1-14.

OFPEL, A. 1856-8. Die Juraformation Englands, Frankreiche und des südwestlichen Deutschlands. Jb. Ver. vaterländ. <u>Naturkde</u>. Württ. 12-14, 1-837.

- OPPEL, A. 1865. Geognostische Studien in dem Ardeche Dept. Pal. Mitt. Mus. k. Bayer. Staates. 5, 305-322.
- PARSONS, C.F. 1974. The sauzei and 'so called' <u>sowerbyi</u> Zones of the Lower Bajocian. <u>Newsl.Stratigr.</u> 3, 153-180.

___, 1975a. The Stratigraphy of the Stony Head Cutting. Proc. Dorset Nat. Hist. Arch. Soc. 96, 8-13.

____, 1975b. Ammonites from the Doulting Conglomerate Bed (Upper Bajocian, Jurassic) of Somerset. <u>Palaeontology</u>. 18, 191-205, pl.36.

____, 1976a. A stratigraphic revision of the humphriesianum/ <u>subfurcatum</u> Zone rocks (Bajocian Stage, Middle Jurassic) of Southern England. <u>Newsl. Stratigr. 5, 114-142</u>.

____, 1976b. Ammonite evidence for dating some Inferior Oolite sections in the north Cotswolds. <u>Proc. Geol. Ass</u>. 87, 45-63, pls.1-2.

, 1979b. A stratigraphic revision of the Inferior Oolite of Dundry Hill, Bristol. <u>Proc. Geol. Ass</u>. 90, 131-151, pls. 1-2.

- PAVIA, G. 1973. Ammoniti del Baiociano Superiore di Digne (Francia SE, Dip. Basses-Alpes). <u>Boll. Soc. Paleont. Ital.</u> 10, 75-142, pls.13-29.
 - , and STURANI, C. 1968. Étude biostratigraphique du Bajocien des Chaines Subalpines aux environs de Digne (Basses-Alpes). <u>Boll. Soc. Geol. Ital.</u> 87, 305-316.
- PRINGLE, J. 1909. On a Boring in the Fullonian and Inferior Oolite at Stowell, Somerset. <u>Geol. Surv. G.B.</u>, <u>Sum.</u> <u>Prog.</u> 1908, 83-86.

, 1910. On a boring at Stowell, Somerset. <u>Geol. Surv. G.B.</u>, <u>Sum. Prog.</u> 1909, 68-70.

QUENSTEDT, F.A. 1886, in 1886-7. <u>Die Ammoniten des Schwäbischen Jura</u>. <u>2. Der Braune Jura</u>. Schweizerbart, Stuttgart, pp.441-815, pls. 55-90. RICHARDSON, L.R. 1907. The Inferior Colite and Contiguous Deposits of the Bath-Doulting District. Q. Jl. geol. Soc. 63, 383-423.

> , 1916. The Inferior Oolite and contiguous deposits of the Doulting-Milborne Port district (Somerset). Q. <u>Jl. geol.</u> <u>Soc. Lond.</u> 71, 473-520, pls.40-43.

, 1932. The Inferior Colite and contiguous deposits of the Sherborne district, Dorset. <u>Proc. Cottes. Nat. F.C.</u> 24, 35-85, pls. 6-11.

, and PARIS, E.T. 1908. On the Stratigraphical and Geographical Distribution of the Inferior Oolite Echinoids of the West of England. <u>Proc. Cottes. Nat. F.C.</u> 18, 72-82, pl.8.

SENIOR, J.R., PARSONS, C.F. and TORRENS, H.S. 1970. New sections in the Inferior Oolite of south Dorset. Proc. Dorset Nat. Hist. Arch. Soc. 91, 114-119.

- SPATH, L.F. 1936. On Bajocian ammonites and belemnites from Eastern Persia (Iran). <u>Palaeont. Indica n.s.</u> 22, <u>mem.</u> 3, 1-21, pl.1.
- STURANI, C. 1964. Ammoniti Mediogiurassiche del Veneto : Faune del Baiociano terminale (zone a Garantiana e a Parkinsoni). <u>Mem. Ist. Geol. Mineral. Univ. Padova.</u> 24, 43pp., 4pls.

, 1967. Ammonites and Stratigraphy of the Bathonian in the Digne-Barreme area. <u>Boll. Soc. Paleont. Ital.</u> 5, 3-57, pls. 1-24.

, 1971. Ammonites and stratigraphy of the "Posidonia alpina" Beds of the Venetian Alps. Mem. Ist. Geol. Mineral. Univ. Padova. 28, 1-190, pls. 1-16.

TORRENS, H.S. 1969. Field Meeting in the Sherborne - Yeovil District. Proc. Geol. Ass. 80, 301-324.

> , 1974. Standard zones of the Bathonian. <u>Mem. B.R.G.M., Fr.</u> 75, 581-604.

WESTERMANN, G.E.G. 1955. Biostratigraphische Untersuchungen im Jura südlich der Sierra de la Demanda (Nord-Spanien). <u>Geol. Jb</u>. 70, 515-534.

> , 1967. Lexique Stratigraphique International; Fascicule 5f2 Jurassique Moyen (Alpes exclues). C.N.R.S., Paris, 197pp.

- WETZEL, W. 1911. Faunistische und stratigraphische Untersuchung der Parkinsonien-schichten des Teutolburger Waldes bei Bielefeld. Palaeontographica 58, 139-278, pls. 11-20.
 - _, 1954. Die Bielefelder Garantianen, Geschichte einer Ammonitengattung. Geol. Jb. 68, 547-586, pls. 11-14.
- WHICHER, J. 1969. New sections in the Inferior Oolite, near Halfway House, Sherborne, Dorset. <u>Proc. Geol. Ass</u>. 80, 324-330.
- WHITE, H.J. OSBORNE. 1923. The Geology of the country south and west of Shaftesbury. <u>Mem. Geol. Surv. G.B.</u> iv + 112pp.
- WILSON, V., WELCH, F.B.A., ROBBIE, J.A. and GREEN, G.W. 1959. Geology of the country around Bridport and Yeovil. <u>Mem. Geol. Surv.</u> <u>G.B. xii + 239pp. 7 pls.</u>
- WOODWARD, H.B. 1894. The Jurassic Rocks of Britain. IV : Lower Oolitic Rocks of England. <u>Mem. Geol. Surv. G.B.</u> 628pp.

· 这些新国家的情绪和考虑了。这些新闻的表达,这些新闻

the second s

Ngar S

2. A STRATIGRAPHIC REVISION OF THE BAJOCIAN ROCKS OF CERTAIN AREAS OF ENGLAND

Much of the detailed stratigraphy of the most important area of Bajocian rocks in England ; the Sherborne district; has already been discussed in section 1. Here I am restricting myself largely to the application of the revised zonal scheme, to three specific areas. First there is Dundry Hill, near Bristol, which with its abundant Lower Bajocian ammonite faunas, figured prominently in the works of d'Orbigny,Oppel and Buckman. Secondly the stratigraphy of the Hiddle and Upper Inferior Oolite of the Cotswold Hills is discussed, since this region was both a key area in the development of Middle Jurassic palaeontology, and was an important factor in the development of Buckman's polyhemeral concept. Lastly the Scarborough Formation of north-east Yorkshire, with its poorly known ammonite faunas, is discussed and revised.

2A. A STRATIGRAPHIC REVISION OF THE INFERIOR OOLITE

OF DUNDRY HILL BRISTOL

CONTENTS

1. INTRODUCTION

- 2. THE PRESERVATION, DEPOSITIONAL ENVIRONMENT AND LITHOSTRATIGRAPHY 172 OF THE DUNDRY INFERIOR OOLITE
 - (a) Preservation and depositional environment
 - (b) A revision of the Lithostratigraphy
- 3. DESCRIPTION OF SECTIONS
 - (a) South Main-road Quarry
 - (b) Rackledown Quarry
 - (c) Barns Batch Spinney
 - (d) Grove Farm

4.

- (e) Old sections
- THE AALENIAN-BAJOCIAN AMMONITE FAUNAS FROM DUNDRY
- (a) The murchisonae Zone
- (b) The concavum Zone
- (c) The discites Zone
- (d) The laeviuscula Zone
- (e) The sauzei Zone
- (f) The humphriesianum Zone
- (g) The garantiana Zone
- (h) The parkinsoni Zone
- 5. CORRELATION OF THE DUNDRY INFERIOR OOLITE
 - (a) The Cotswolds
 - (b) The Bruton district

THE HORIZON OF BUCKMAN'S AND SOWERBYS' TYPES 6. 205 (a) [Sowerbys' type specimens (b) Buckman's specimens ACKNOWLEDGEMENTS 210 REFERENCES 211 我是你们的,你就能没有能到我都能能能好吧。" an destant an 的复数使用的复数 化拉拉拉拉拉拉拉拉拉拉 等于是理论之后,如于现于是解释的中国,而此不是是实际上的原本的问题。

<u>ABSTRACT</u>: The Lithostratigraphy of the Inferior Oolite Group (Aalenian and Bajocian, Middle Jurassic) of Dundry Hill, Bristol, Avon, is revised and several new lithostratigraphic units are erected: the Barns Batch, Grove Farm and Elton Farm Limestones. The depositional environment and present pattern of preservation of these beds are discussed and four exposures are described in detail. An analysis of the Aalenian-Bajocian ammonite faunas is made and they are correlated with their equivalents elsewhere in southern England, particularly those recently collected from the Bruton district of Somerset. Finally the type horizons of the Sowerbys' and Buckman's ammonite species are discussed, and the majority assigned to their correct stratigraphic position.

1. INTRODUCTION

The Inferior Colite (Aalenian-Bajocian stages) of Dundry Hill, near Bristol, Avon, is one of the World's most famous fossiliferous deposits. The quarries on this hill, which were opened during the exploitation of the famous Dundry Freestone ("Upper Inferior Colite"), often showed sections in the highly fossiliferous and 'condensed' "Lower and Middle Inferior Colite" (Aalenian-Lower Bajocian). It was the abundant ammonites from these horizons, which attracted the attention of the early palaeontologists and stratigraphers, such as d'Orbigny (1842-51), Oppel (1856-8), the Sowerbys (1812-46) and Wright (1860). Other sections of the fauna, such as the brachiopods, gastropods and bivalves are equally prolific and they figured in the early work of numerous palaeontologists, such as S.S. Buckman, Moore, Stoddart, Tawney and S. Woodward. However,

STAGE	ZONE		SUBZONE			
	n an	incori	Bomfordi			
	Park.		Truellei			
UPPER	Gara	ntiana	Acris			
	n an	an a	Dichotoma			
BAJOCIAN			Baculata			
nge sprjagsterne. Se	Subfu	ırcatum	Polygyralis			
	and a start of the second start		Banksi			
			Blagdeni			
	Humpl	nriesianum	Humphriesianum			
LOWER			Romani			
	Sauze	Sauzei				
	Laev	inscula	Laeviuscula			
BAJOCIAN			Ovalis			
	Disci	l tes				
	Conca	Concavum				
	a Print and Angeler and Ang Angeler and Angeler and Ange Angeler and Angeler and Ang	Nurchisonae	Bradfordensis			
AALENT AN	Nurch		Murchisonae			
			Haugi			
			Scissum			
	Opali	Inum	Opalinum			

Table 1.

JZAHAT The standard Zones and Subzones of the Aalenian and Bajocian Stages. Bajocian Stages.

as early as the 1890's these famous exposures were becoming obscured, as the quarries closed down, owing to the depletion of the Freestone. Our knowledge of the stratigraphy of these beds has thus rested on the classic work of S.S. Buckman and E. Wilson (1896 and 1897), who reopened many already disused quarries and carefully described those still in work. Whilst Buckman was later to figure numerous Dundry ammonites as new species and genera in 'Type Ammonites' (Buckman, 1909-30), all subsequent accounts of the stratigraphy have rested on the earlier work. Thus the South Main-road quarry, perhaps the most famous Dundry section has been re-described several times by Tutcher (1903, and in Reynolds <u>et al.</u>, 1930) and Macfadyen (1970), but these accounts have only been slight modifications of Buckman and Wilson's work.

The sections now available for study on Dundry have deteriorated even more since ^Buckman's day. However, there are strong reasons for giving a modern description of these rocks. The ammonite faunas from the Dundry <u>laeviuscula</u> and <u>sauzei</u> Zones (see Table 1) are unparalleled in Europe for their abundance and diversity: they are thus of extreme importance in elucidating both, the stratigraphy of this period and the phylogeny and systematics of the ammonites, particularly the Stephanoceratidae. Whilst Buckman (1909-30) figured a number of ammonites from Dundry, often little is known of their stratigraphic range. The following account is thus primarily an attempt to revise the litho- and biostratigraphy of the Dundry Inferior Colite and to determine the correct stratigraphic position of as many as possible of the previously recorded ammonites.

NOTE: In the following, specimen numbers preceded by, BUGM. refer to ammonites in the collections of the Department of Geology, University of

Bristol; by J. to the Oxford University Museum and by CP to the author's collection. In the faunal lists the citing of a single specimen number refers to the specimen, upon which the identification is based, and is no indication of the number of individuals collected.

- 2. THE PRESERVATION, DEPOSITIONAL ENVIRONMENT AND LITHOSTRATIGRAPHY OF THE DUNDRY INFERIOR OOLITE
- (a) Preservation and depositional environment

In terms of depositional environment and existing pattern of preservation, the Inferior Colite on Dundry Hill falls into two divisions (Cleak, 1930). Firstly there is the Stroud Formation (="Upper Inferior Oolite"), a series of bioclastic and colitic limestones of variable thickness, which are present over all the hill. Secondly there is the "Lower and Middle Inferior Oolite", a series of thin, "condensed", more or less 'iron-shot' limestones, which are found only on the western half of the hill. In general facies and lithology, the Stroud Formation is very similar to its equivalents in the Cotswolds, whilst the "Lower and Middle Inferior Oolite" have no similarity to their Cotswold equivalents, but rather to the north Dorset deposits of the same age. These differences are not due to any physical barrier separating the Dundry and Cotswold areas during the Aalenian-Bajocian, as suggested by Buckman (1889, 1901), but of course to original variations in depositional environment. The Lower Bajocian rocks of Dundry show the closest similarity in facies to Dorset, with the prevalence of limonite coliths, 'snuff-boxes' (complex laminated limonite concretions, q.v. Gatrall et

<u>al</u>., 1972), limonite-incrustation, thick shelled bivalves and abundant ammonites. These characters all point to a very slow rate of deposition, dominated by ferruginous, authigenic minerals, in a shallow sea. This area of shallow water was probably caused by the presence of a broad shelf, or similar palaeo-topographic feature, to the west of the Cotswold basin, which acted as a 'clastic trap'. The Stroud Formation on the other hand also shows signs of a relatively strong current regime, which however, was insufficient to remove the fragmented byproducts of a high rate of primary carbonate production.

Separating these two very different facies is a prominent 'hardground'; a very flat, bored, sparsely oyster-encrusted surface. Resting on this 'hard-ground' is usually a conglomerate, consisting of limonite coated lithoclasts, derived from subjacent beds. Taking this evidence of erosion into account, Buckman's explanation of the varied preservation of the Aalenian-Lower Bajocian rocks is probably the right one (Buckman & Wilson, 1896, p.695). He attributed it, as in the Cotswolds, to minor flexuring of the rocks, prior to the onset of the Upper Bajocian transgression. Thus a small "down-warp" in the region of the main Bristol road resulted in the local preservation of the Elton Farm Limestone, whilst to the west erosion cut down to the ovalis bed and to the east as far as the Upper Lias (Arkell, 1933, fig. 34). Although this is the best overall explanation of the present patterns of preservation, there is some evidence of an original primary control of deposition. Thus the upper part of the Elton Farm Limestone (Brown Iron-shot and Witchellia beds) shows signs of a change in lithology, (sparser coliths, fewer fossils), as it thickens from the South Main-road quarry, towards

	Stroud Formation or "Upper I.O."	Coralline beds (6.0m+) Dundry Freestone Member Maes Knoll Conglomerate Member (0 - 0.68m)	(0 - 6.0m)
INFERIOR		GAP	
OOLITE	"Middle I.O."	Elton Farm Limestone Member (0 - 1.9m)	2200 - 2442 - 246 - 446 - 472 - 474 - 474 - 475 - 474 - 474 - 474 - 474 - 474 - 474 - 474 - 474 - 474 - 474 - 4
GROUP		Grove Farm Limestone Member (0 - 1.0m)	:
an Santa Managaran Managaran	"Lower Inferior Colite"	SLIGHT GAP	
		Barns Batch Limestone Member (0 - +3.0m)	

Table 2.

The lithostratigraphic subdivisions of the Inferior Oolite Group recognised on Dundry Hill. Rackledown. The possibility must remain that these isolated beds originally filled a shallow negative area on the sea-floor. This in no way contradicts Buckman's original concept of an overall secondary control in preservation, although this "penecontemporaneous erosion" may have in part accentuated some pre-existing minor differences in bed thickness.

(b) A revision of the Lithostratigraphy

The existing informal lithostratigraphic terms for the Dundry Inferior Oolite, introduced haphazardly by Buckman and Wilson are highly ambiguous. This is particularly true of the individual beds in the Thus the term 'White Iron-shot', which was used Elton Farm Limestone. by Buckman for two horizons, is here shown to include three. The division between Buckman's so called 'Iron-shot' and the 'Upper White Ironshot! must be arbitrary as the boundary between them is usually of a gradational nature; no division between these two should be attempted solely on the basis of lithology. However, this does not totally preclude the use of matrix in determining the position of unlocalised specimens, as the topmost, dark, purple stained, part of the Brown Ironshot bed is totally characteristic, and restricted to this horizon. It is thus suggested that, in order to reduce confusion, the following named units should be used. These have been based on the degree of lithological similarity between components, and the boundaries taken at sharp lithological changes and unconformities - see Table 2.

In accordance with the recommendations of the Geological Society of London (Harland <u>et al.</u>, 1972), the Dundry Inferior Oolite should be divided into a series of formal hierarchical units. In terms of its overall lithology, present pattern of preservation and relationship with nearby equivalent beds, the Dundry Inferior Oolite can be divided into three sections, which could be accorded the rank of Formations. However, the area of outcrop, particularly of the "Lower and Middle Inferior Oolite" is so small, that it is thought inadvisable to erect new formational names for these beds. The "Upper Inferior Oolite" is merely an extension of the equivalent Cotswold beds, hence there is no problem in including it in the Stroud Formation of the Inferior Oolite Group (Parsons, 1980). The lower beds are more problematic, as there are no pronounced changes in lithology, and they would probably be best placed in a single new Formation (cf. 'Oborne beds', Parsons, 1980, p. 222 here). This is not the place for the erection of such a new unit, thus as an interim measure these beds are included in the existing informal subdivisions - see Table 2. The Inferior Oolite has been divided into six Members, which are described below, starting from the base.

(i) The Barns Batch Limestone Member

The lowest Inferior Oolite now exposed on Dundry Hill is to be seen at Barns Batch Spinney. These beds form a series of hard oolitic limestones, whose massive nature enables them to be easily traced around the hill. The upper surface of this unit, the Barns Batch Limestone, is marked by a prominent 'hard-ground', which is very flat, limonite-encrusted, bored by <u>'Lithophaga</u>' and surmounted by a conglomerate. These beds at Barns Batch Spinney, the type locality, are +1.65m thick (beds 1 - 3, section 3c) and span the <u>murchisonae / haugi</u>

Subzones of the <u>murchisonae</u> Zone - see Table 1 for zonal scheme used here. It is impossible to exactly define the lower limits of these beds, as no exposure is now available showing <u>opalinum</u> Zone rocks, (Buckman & Wilson, 1896, p.677). Since there is an unconformity at the base of these <u>opalinum</u> Zone rocks at Castle Farm (<u>loc. cit.</u>), where they rest on blue, argillaceous <u>aalensis</u> Subzone rocks (<u>loc. cit.</u>); bed 24 at that section may be taken as defining the base of the Barns Batch Member, which are there 2.40m. thick (<u>loc. cit.</u>). At Rackledown Farm, these beds are slightly thicker, as they must be in excess of 3.0m. thick (<u>op. cit.</u>, p.692).

The upper bed of this unit is one of the most distinctive of those on the hill, and it may be informally separated as the <u>Pleurotomaria</u> bed (bed 3a, section 3c) in recognition of the large numbers of this group of gastropods recorded from it, (Buckman & Wilson, 1896, Table Va).

(ii) The Grove Farm Limestone Member

The beds called by Buckman the 'Nodular beds', (Buckman, 1892, in 1887-1907, p.293) or the 'limestone and marl' beds, (Buckman, 1901, p.154) are here named the Grove Farm Limestones, after the exposures in the region of Grove Farm, West Dundry. These beds consist of nodular, slightly 'iron-shot' limestones, interbedded with soft marls. They are best seen at Barns Batch Spinney, which may be taken as the type locality, where they are 0.9m. thick, (beds 4-6, section 3c), and span the <u>concavum</u> Zone and the basal part of the <u>discites</u> Zone. The basal beds of this unit (e.g. Barns Batch, bed 4), which are rich in <u>Sphaeroidothyis</u>, may be informally separated as the <u>eudesi</u> beds, (Buckman & Wilson, 1896, p.672).

BUCKMAN 1892	BUCKMAN 1893	BUCKMAN & WILSON 1896	BUCKMAN 1901	TUTCHER 1903	TUTCHER 1929	TUTCHER 1930	SCHEME USE HERE	ED
IRON-SHOT BED	IRON-SHOT BED	THE IRON-SHOT	THE IRON-SHOT	THE FOSSIL BED	THE DUNDRY IRON-SHOT	BROWN IRON-SHOT	BROWN IRON-SHOT BED	(1)
	WHITE IRON-SHOT BED	UPPER WHITE IRON-SHOT Witchellia bed	UPPER WHITE IRON-SHOT			UPPER WHITE IRON-SHOT	Witchellia BED	RM LIMESTONE
							LIMONITIC BED	ON FA
WHITE BED BELOW THE IRON-SHOT		LOWER WHITE IRON-SHOT	LOWER WHITE IRON-SHOT			LOWER WHITE IRON-SHOT	ovalis BED	ELT
							BIVALVE	

TABLE 3 Development of the informal lithostratigraphic subdivisions of the Dundry Iron-shot beds.

(iii) The Elton Farm Limestone Member

The beds called by Buckman the 'Iron-shot', (Buckman, 1892 in 1887-1907, p.282) and 'White Iron-shot', (Buckman, 1893, p.508), which was subsequently separated as 'Upper and Lower White Iron-shots', (Buckman & Wilson, 1896, pp.678 & 682), are too thin and poorly defined to warrant formal status as litho-stratigraphic units. The terms 'Ironshot! and 'Lower and Upper White Iron-shot! were introduced by Buckman for the purpose of recording the ammonite fauna; there is little lithological basis for these divisions. Thus, for example, there is usually a gradational boundary between the 'Iron-shot' and the 'Upper White Iron-shot! at both South Main-road quarry and Rackledown, making it impossible to delimit a formal boundary between these two. There has been considerable confusion over the interpretation of the term 'White Iron-shot . Buckman first introduced the term 'White bed below the Iron-shot! (Buckman, 1892, in 1887-1907, p.293; Wilson, 1893) for the horizon which he afterwards called the 'Lower White Iron-shot', (Buckman & Wilson, 1896, p.681, bed 9). Buckman later (1893, p.508) introduced the term 'White Iron-shot' for beds equivalent to all those below the Brown Iron-shot! and above the 'Limestone/marl beds'. The restricted term 'Lower White Iron-shot' was subsequently solely used for the fissilobatum/ovalis horizon, (Buckman & Wilson, 1896, pp.678 and 676, beds 4-8), whilst the 'Upper White Iron-shot' was restricted to the 'Witchellia bed' (op. cit. p.681, bed 6). Unfortunately this refined usage overlooked the middle bed of the original 'White Iron-shot' (loc. cit., bed 8), which in patches is as densely 'iron-shot' as the true Brown Iron-shot sensu stricto. The time has thus come for some attempt to be made to simplify this muddled litho-stratigraphy, which is

summarised in Table 3.

Since there are no grounds for giving formal names to all the thin, highly fossiliferous beds of the 'Middle Inferior Oolite', these may all be grouped together within one formal unit, the Elton Farm Limestone. This is named after the famous exposures in these beds found close to Elton Farm at the South Main-road quarry. The latter quarry may be taken as the type locality of this unit and beds 7-10 (section 3a herein) as the type horizon, which is some 1.7m. + thick. These beds are of discites-sauzei Zones in age. For the smaller subdivisions of this formal unit the following informal bed names may be used, the ovalis bed (Parsons, 1977, p.101) instead of 'Lower White Iron-shot' (= fissilobatum/ovalis horizon, Buckman & Wilson, 1896, p.681) as for bed 8 at Barns Batch spinney, the Limonitic bed (= bed 9, South Main-road), the Witchellia bed solely for the one bed with the soft, white, pasty matrix (= 'Upper White Iron-shot' sensu Buckman & Wilson, 1896), that is bed 10a South Main-road quarry, since this is closest to Buckman's original usage (Buckman, 1893), and the Brown Ironshot bed for 10b, South Main-road, and its equivalents.

(iv) The Maes Knoll Conglomerate Member

At the base of the Upper Inferior Colite on Dundry, there is an extensive 'hard-ground', marked by a prominent flat, bored and cysterencrusted surface. Directly above this 'hard-ground' at various localities there is usually a conglomerate consisting of limonite- and serpulid-encrusted lithoclasts, set in an 'iron-shot' limestone matrix. This conglomerate was first described by Buckman and Wilson, (1896, p.686 & Table IV) and this formal lithological name follows their usage

and Buckman's subsequent use of the term 'the conglomerate bed' (Buckman, 1901, p.154), which was later amended to the 'Maes Knoll Conglomerate-bed' (Richardson, 1907, p.420). The type locality of this unit can be taken as Maes Knoll, where according to Buckman and Wilson (1896, p.684) the total thickness of 'iron-shot' limestone present is 0.68m. Only the basal part of these beds at this locality is a true conglomerate, but all the 'iron-shot' limestones below the Dundry Freestone may be included in this formal unit. The ammonites found in these beds would indicate that they are <u>acris</u> Subzone, <u>garantiana</u> Zone in age.

(v) The Dundry Freestone Member

The presence of this limestone on Dundry Hill was the reason for the extensive quarrying activities which have been carried out since the Roman period. The term Dundry Freestone has been suggested as a formal name (Worssam in Donovan & Hemingway, 1963, p.120) and Dundry Hill designated its type locality (<u>loc. cit.</u>). This unit consists of a series of massively bedded freestones at their thickest at the west end of the Hill. One of the few existing sections of this unit on Dundry Down (ST 552666), which shows 4.6m of massive bioclastic limestone, overlain by 1.3m of more thinly bedded limestone, can be taken as its type locality. No exposure now shows a complete section through these beds, it is thus difficult to determine their exact stratigraphic Position, although available ammonite evidence would point to a basal <u>Parkinsoni</u> Zone age.

(vi) The Coralline beds

These beds were so called by Buckman and Wilson (1896, pp.672 &



Figure 1.

A sketch map of Dundry Hill, showing the main localities cited in the text. 1 - South Main-road quarry, 2 - Rachledown quarry, 3 - Barns Batch Spinney, 4 - Grove Farm, 5 - Castle Farm, 6 - North Main-road quarry, 7 - Malnut Farm quarry. 679 <u>et seq</u>.) on account of the numerous coral fragments present. Since no complete exposures can be seen of this unit, the total thickness is unknown but it must be in excess of 6.0m. (Donovan, 1958, p.132); no formal type locality is thus designated here, since more information is needed on the lateral and vertical extent of these beds.

3. DESCRIPTION OF SECTIONS

Since the main aim of this work was to carry out a stratigraphic revision of the Inferior Oolite of Dundry Hill, only those sections which have yielded ammonites at all abundantly will be described here. This naturally precludes the sections in the Dundry Freestone and Coralline Beds, since these two have produced little or no ammonite evidence as to their age. The following sections are thus all those now showing exposures in the richly fossiliferous

"Middle and Lower Inferior Oolite", Aalenian and Lower Bajocian Stages, that is: the 'West End' (Castle Farm-Grove Farm), Barns Batch Spinney, South Main-road and Rackledown quarries, see Text fig.1. for their location. Although all these exposures were described by Buckman and Wilson, (1896 & 1897), the last 80 years of research on the ammonite faunas has revealed a need for their revision. In these redescribed sections, all the beds have been renumbered using a single set of numbers based on a hypothetical complete sequence (see Text fig.3), but where applicable Buckman and Wilson's numbers follow the new ones in brackets.

(a) The South Main-road Quarry

This section (ST567655), which according to Buckman and Wilson,
(1896, p.691) was very similar to the famous North Main-road quarry, has in recent years been largely obscured. During the period 1955-58, the Bristol Naturalist Society cleared a section down to the <u>murchisonae</u> Zone limestones. Unfortunately the uncontrolled action of fossil collectors soon filled the quarry in again, until little but the Upper Inferior Oolite was visible. Thanks to the recent work (Sept. 1979) of the Nature Conservancy Council the quarry has again been re-excavated:

The Coralline Beds

13. (1)

A series of white, bioclastic limestones, much of which are cambered and slipped material, mixed with clay and rubble.

seen to 1.5m.

The Dundry Freestone

A compact, well bedded limestone, oolitic and slightly 'iron-shot', especially towards the base. The probable lateral equivalent of the thicker Dundry Freestone to the west.

0.48 - 1.0m.

brown marl, resting on flat surface

The Maes Knoll Conglomerate

11. (3)

An ironstained conglomerate, consisting of limonite- and serpulidencrusted pebbles, set in a slightly 'iron-shot' limestone matrix. The erosion plane, on which this bed rests, is limonite stained, bored and sparsely oyster-encrusted.

Prorsisphinctes sp., CP3787

0.20 - 0.36m.

The Elton Farm Limestone

very flat surface

10b. (4) The Brown Iron-shot bed

A densely 'iron-shot' limestone, with shiny, brown limonite ooliths set in a purple tinged matrix. This, the 'Iron-shot' of Buckman, (Buckman & Wilson, 1896) is the horizon long famous for its fossils, particularly the abundant ammonites. This bed seems to thicken towards the south, and is marked at its base by a very poor parting, due to the presence of large numbers of bivalves; Liostrea and Ctenostreon.

Emileia (E.) bulligera Buckman, CP 3796 E. (E.) polyschides (Wag.) BMNH. C80497 E. (E.) polymera (Wa.) CP.2423 E. (Otoites) cf. sauzei (d'Orb.) CP 2448 E. (Otoites) aff. fortis (West.) CP.2425 Labyrinthoceras aff. meniscum (Waagen) CP.2402 Stephanoceras (Kumatostephanus) perjucundus Buckman BMNH.C80496 Stephanoceras (Skirroceras) bayleanum (Oppel) CP.3771 S. (S.) macrum (Weisert non Qu.), CP 3769 S. (Normannites) sp. Sonninia (S.) corrugata (Buckman non Sow.) CP 3792 S. (S.) aff. felix (Buckman), CP.2422 S. (Papilliceras) mesacantha (Waagen), CP 3809 ?Witchellia (?W.) hebridica (Morton) CP.2442 <u>?W. (Pelekodites) cf. sulcata</u> (Buckman) CP.2451 Protoecotraustes spiniger (Buckman) (Torrens Col.)

arbitrary boundary

10a. (5) The <u>Witchellia</u> bed

0.22 - 0.28m.

Total for bed 10 =

0.40 - 0.52m.

- Irregular, limonite coated surface

9. (?6) The Limonitic bed

사람들은 이 것은 영국에서는 소설적 수전에 관광했다. 관

A very nodular, hard and 'iron-shot' limestone. The irregular nature of this bed, due to bioturbation, together with the limonite incrustation gives the false impression that it is conglomeratic. The upper surface is thickly coated with limonite, and with some incipient 'snuff-boxes', (q.v. Gatrall, <u>et al.</u>, 1972, p.85). There are many fossils present, but they tend to be in either the hard limestone nodules and thus difficult to extract, or in the limonitic partings and hence distorted and badly preserved.

Stephanoceras (Skirroceras) sp.

Emileia (Otoites) sp. nov. A., CP.2404 Trilobiticeras (T.) cricki Parsons, CP.2403 Mollistephanus (M.) aff. mollis Buckman, CP.2405 Shirbuirnia stephani (Buckman), BMNH. C80498 Witchellia (W.) aff. romanoides (Douvillé), CP.2435 W. (W.) pavimentaria (Buckman), CP.2433 S. (Euhoploceras) cf. acanthera (Buckman), CP.2429 Lissoceras aff. semicostulatum Buckman, CP.2443

0.20 - 0.25m.

8 a + b. (?7) The <u>ovalis</u> bed

A finely 'iron-shot' limestone, crystalline and with a pinkish matrix. The top of the bed (8b) tends to be nodular and limonite stained, whilst the base (8a) is more crystalline and massive with sparser limonite ooliths. There is a rich fauna, particularly of bivalves, as in the same horizon at Barns Batch, which has a similar lithology.

Sonninia (Fissilobiceras) ovalis (Buckman ex. Qu.), CP.2445

<u>Witchellia (W.) romanoides</u>, CP.2437 <u>W. (W.) aff. connata</u> (Buckman), CP.2441 <u>Trilobiticeras (T.) cricki</u>, CP.2971

0.60m.

irregular parting and pockets of orange/red, sandy marl

7.

A hard, grey-brown, nodular, slightly iron-shot limestone.

Sonninia (Euhoploceras) marginata Buckman (loose, but possibly from here by matrix)

0.20m.

Collections made by the Bristol Naturalists show that horizons lower than that currently visible, were exposed in 1955. The following specimens are from the Stenhouse-Ross collection in Oxford University Museum, and indicate the presence of the <u>discites/concavum</u> Zones in the marly Grove Farm Limestone and the <u>murchisonae</u> Zone in the top of the Barns Batch Limestone. There is also an extensive collection of similar material in the Bristol City Museum.

Explanation of Plate 1.

Figure 1.

A general view of the South-Main-road quarry, Dundry Hill, showing the highly disturbed and cambered state of the beds.

Rigure 2.

South-Main-road quarry, Dundry, showing a close-up of the Dundry 'Iron-shot' beds.



Bed 9' (8) <u>Ludwigia</u> (<u>Ludwigia</u>) sp. J16767 (in an iron-shot matrix, with limonite incrustation, pisoliths and small 'snuffboxes' see Gatrall <u>et al.</u>, 1972).

The bed numbers used above are those given by Stenhouse-Ross, whilst those given in brackets are the probable Buckman and Wilson (1896) equivalents.

(b) Rackledown Quarry

This is an old quarry (ST572654), which, even in Buckman's day was not good for detailed stratigraphic collecting, as cambering and tufa coating of the faces made it difficult to interpret. Since then, dumping of rubbish, as well as more natural deterioration has done nothing to help the visiting geologist. It is impossible to see a continuous section of those beds still visible in April 1972; the following is thus a composite description taken from several separate exposures:

The Coralline Beds

A tumbled mass of white coloured, bioclastic limestone, rich in broken corals; probably not seen anywhere in situ.

1.0m.+

The Elton Farm Limestone

10b/ (3) The Brown Iron-shot bed

A densely 'iron-shot' limestone with a dark, almost purple matrix. Visible as a steeply dipping cambered block at the right-hand side of the short quarry face, opposite the road, this bed may also be seen <u>in situ</u> on the hill slope to the south. The exact thickness of this bed is difficult to determine, since there is a transition to bed 10a. Emileia (E.) cf. bulligera Buckman, CP.1476 E. (E.) cf. vagabunda, Buckman, CP.2140 Stephanoceras (Kumatostephanus) sp. S. (Skirroceras) aff. bayleanum (Oppel) Bradfordia sp. CP.2138

Strigoceras (Cadomoceras) ellipticum, (Buckman) CP.2139

Sonninia (S.) aff. propinguans (Bayle), CP.2137

c.0.34m.

10a/ (3) The Witchellia bed

A slightly less densely 'iron-shot' limestone than the bed above, with a lighter, more white coloured matrix.

Shirbuirnia aff. trigonata (Buckman ex. Qu.) CP.1479

Sonninia (Fissilobiceras) aff. fissilobata (Waagen.), CP.1480

Witchellia (W.) laeviuscula (Sow.), CP.2129

W. (Pelekodites) aurifer (Buckman), CP.2135

Various other unlocalised specimens, undoubtedly from this horizon, by matrix include:-

W. (W.) spinifera Buckman, CP.2132

W. (W.) cf. sutneri (Branco) CP.2130

c.0.50m.

Total for bed 10 = 0.90m.

- Break in section - which continues in the long southern face

8/(?4 pars) The ovalis bed

A rubble of broken up and weathered 'iron-shot' limestone, which grades into the top of bed 7.

Trilobitioeras (Emileites) aff. malenotatus Buckman, CP.2167 T. (E.) liebi (Maubeuge), CP.1979

Witchellia (W.) romanoides CP.2133

seen to 0.25m.

7/ (4)

A massive, well bedded 'iron-shot', crystalline limestone, with a pink matrix. This is the lowest bed seen in situ in the quarry.

Hyperlioceras (Hyperlioceras) sp.

seen to 0.7m.

Break in section, the Grove Farm -Limestone is not exposed

The Barns Batch Limestone

3/ (5) The <u>Pleurotomaria</u> bed

In the field, on the hill slope to the south of the quarry, the flat iron stained top to this unit is clearly visible. As at Barns Batch, this finely 'iron-shot', massive limestone, has a conglomeratic upper surface, rich in limonite pisoliths, 'snuffboxes' and limonite laminae.

Ludwigia (L.) crassa (Contini non Horn), EMNH. C80495

seen to 0.30m.

(c) Barns Batch Spinney

This section was described by Buckman and Wilson (1896, p.689), but it appears to have been subsequently ignored. Although it now only consists of a shallow pit (see Plate 2, figs.1 and 2) in a clump of trees (ST557659), after some excavation in April 1972, it revealed the following details: The Coralline Beds

13 (1)

White, rubbly, bioclastic limestones, which mainly consist of disturbed and slipped material, although it may be seen in situ in the pit next to the road (ST554659).

0.3m. +

A planed, limonite-stained surface

The Elton Farm Limestone (pars)

9. (pars 2)

A thin, 'iron-shot', conglomeratic limestone, which thins rapidly towards the west. Much of the fauna from this bed is rolled and worn looking and has been derived from the bed below, since the indigenous forms are much better preserved.

Emileia (E.) sp. nov. A., CP.1463
Derived <u>Witchellia</u> (W.) cf. <u>romanoides</u>, CP.1461
W. (W.) aff. <u>sayni</u> Haug, CP.1458
<u>Trilobiticeras</u> (<u>T.</u>) <u>cricki</u>, CP.1452
<u>T. (Emileites) liebi</u>, CP.1448

0.0 - 0.1m.

- An irregular, limonite-stained surface -

8a - b (pars 2)

A cream-grey, rubbly limestone, with numerous small limonite ooliths (8b), which grades down into a harder limestone, with a pinker, more crystalline matrix (8a). This horizon is highly fossiliferous, with the fauna mainly concentrated at 0.1 - 0.15m. and 0.25m. below the top. In the latter case the brittle fracture

of the rock makes it more difficult to extract.

Witchellia (W.) albida (Buckman), CP.1455 W. (W.) cf. connata (Buckman), CP.1457 W. (W.) romanoides, BMNH. C80494 W. (W.) cf. sutneri, CP.1424 W. (Pelekodites) pelekus Buckman, CP.2523 <u>W. (P.) macra, CP.2525</u> S. (Fissilobiceras) ovalis, CP.1459 S. (F.) gingensis (Waagen), CP.2685 S. (?Euhoploceras) sp. Bradfordia cf. inclusa, CP.2508 Toxamblyites sp., CP.1418 Strigoceras (S.) compressum (Buckman), CP.1419 Docidoceras cf. cylindroides Buckman, CP.1481 Emileia (Otoites) douvillei Parsons, BMNH. C79429 Trilobiticeras (T.) cricki, BMNH. C79426 T. (Emileitas) liebi, BMNH. C80493

0.25 - 0.45m.

7 (3)

A hard, massive, crystalline, 'iron-shot' limestone, with a pink matrix, and which, except for a few belemnites, is poorly fossiliferous. This bed is roughly separated into two courses, by an irregular sandy parting, either side of which this bed is sandier and more nodular, due to intense bio-turbation.

Hyperlioceras sp.

0.3 - 0.48m.

A brown sandy marl 0.05m.

The Grove Farm Limestone (6-4)

6 (4)

A hard, nodular, crystalline limestone, with minor marl intercalations, which becomes softer and whiter towards the base. Divided roughly in half by a poor parting, this bed is extensively bio-turbated, and although fairly fossiliferous, most of the shell material has been leached out, to leave crushed and distorted internal casts.

'Braunsina' cf. projecta Buckman, CP.1497

Hyperlioceras (H.) walkeri Buckman, CP.1496

<u>Н. (Н.)</u> врр.

化化化性 化化化学 意义的变形 建立的

?Darellia (?D.) aff. polita Buckman, CP.1495

<u>S. (Euhoploceras) cf. acanthodes</u> Buckman, (from here by matrix), CP.1660

0.3m.

0.2m.

- A marl parting - -

5 (pars 5)

A series of hard, pink-gray, slightly 'iron-shot', limestone nodules, set in a soft marl. There are numerous poorly preserved ammonites present, mainly distorted, internal casts of bodychambers.

<u>Graphoceras</u> (G.) <u>formosum</u> (Buckman), CP.1499 <u>G.</u> (G.) spp. 4 (6) The eudesi bed

A pink-grey, limestone, which is harder and more massive than the bed above. Divided into three courses by two partings; the middle course contains abundant <u>Sphaeroidothyris eudesi</u> (Oppel), whilst the basal course is darker coloured, more crystalline, contains derived pisoliths and has yielded one ammonite.

'Braunsina' aff. rotabilis (Buckman), CP.2158

0.4m.

CP.2161

The Barns Batch Limestone (3-1)

A prominent, flat, limonite-encrusted surface

3b (pars 7) A thin, laminated limonite layer

0.02 - 0.04m.

3a (pars 7) The Pleurotomaria bed

A hard, pink-grey, 'iron-shot' limestone. This bed is particularly 'iron-shot' towards its very irregular top, where it is also highly conglomeratic, bored by 'Lithophaga', and where it contains large pisoliths, serpulid- and limonite-encrusted ammonites and small 'snuff-boxes' (q.v. Gatrall, <u>et al.</u>, 1972). Particularly hard and massive towards its base, this bed, together with those below, forms a distinctive topographic feature, in contrast to the softer, more marly limestones above.

From the top (- Brasilia (Brasilina) cf. tutcheri (Buckman), (CP.2156

- Ludwigia (Ludwigina) sp.

From the middle (- Ludwigia (L.) murchisonae (J. de C.Sow.),

Brasilia (Brasilina) sp.

Explanation of Plate 2.

Figure 1.

A general view of the Barns Batch Spinney section, on Dundry Hill, showing its degraded state.

Figure 2.

이 같은 집에 있는 것이

A close-up of the ovalis bed at Barns Batch Spinney. The hammer is resting on the flat, planed off top of the latter bed.



Figure 1.



From the base - ?B. (?Brasilia) cf. wilsoni (Buckman) CP.2155 0.5m.

2 (<u>pars</u> 7)

A hard, pink-grey, crystalline, sandy, sparsely oolitic, very nodular limestone, with some marl layers. This horizon is well bio-turbated, probably by the <u>Pleuromya</u> to be found in life position.

Ludwigia (L.) obtusa (Buckman ex. Qu.), CP.2159

L. (L.) obtusiformis (Buckman), CP.2162

0.28 - 0.30m.

1 (pars 7)

A more massive, buff coloured, sandy limestone, with small limonite flecks.

seen to 0.85m.

(d) Grove Farm

This exposure at the 'West End' of Dundry, (ST551671), is quite near to the location of the excavations which Buckman and Wilson (1896, pp.676-8) had made at Castle Farm. The present section was measured in July 1975, from a series of natural scarps on the north slope of Dundry Hill, overlooking the road; hence a less detailed subdivision of the beds was possible, than that undertaken by Buckman. Taking this into account, the section described here is very similar to that described from Castle Farm (loc. cit.):

The Maes Knoll Conglomerate

11 (3)

A conglomeratic, brown, 'iron-shot', highly fossiliferous limestone.

Discacyathus sp.

Trautscholdia spissa (Buckman)

0.0 - 0.15m.

The Elton Farm Limestone (8-6)

A limonite-stained, very flat surface

8a-b. (4-7) The ovalis bed

A cream-grey, 'iron-shot', limestone, divisible into two courses by a poor parting. The bottom course (8a), is harder and more nodular, with a more crystalline matrix and sparser iron coliths, than the top course (8b), which is softer, more rubbly and which contains more fossils, particularly bivalves. Most of the fossils recorded here come from -0.10m. below the top of the bed.

<u>S. (F.) ovalis</u>, CP.2846, at -0.40m. <u>Witchellia (W.) romanoides</u>, CP.1409 <u>W. (W.) cf. sutneri</u>, CP.1407 <u>W. (W.) aff. connata</u>, CP.1408 <u>S. ('Euhoploceras') sp.</u> <u>Docidoceras aff. cylindroides</u>, CP.1417 <u>Trilobiticeras (T.) cricki</u>, CP.1414

T. (Emileites) sp.

Strigoceras (S.) sp.

0.48m

0.43m.

---- A sandy, marl parting ------

7a-b (9-11)

A hard, crystalline, sandy, slightly 'iron-shot' limestone, divided into two courses by a sandy marl parting, and which has

abundant, badly preserved bivalves, particularly in the lower course (7a).

- - A sandy, marl parting -

The Grove Farm Limestone

6 (13)

A grey, nodular, slightly 'iron-shot' limestone, intermixed with a soft marl. The common fossils are preserved as distorted internal casts.

Hyperlioceras (H.) deflexum Buckman, BMNH. C80492 H. (H.) spp.

?Darellia (?D.) aff. polita, CP.1406

0.12-0.20m.

+ 0.45m.

5 (14) A soft, grey coloured limestone.

4 (15-18) Massive, slightly 'iron-shot' limestone, with subsidiary marl partings.

Graphoceras (G.) pulchra (Buckman), CP.2839

(e) Old sections

Several other sections on Dundry have in the past exposed the Elton Farm Limestone. The most famous Dundry locality is undoubtedly the North Main-road quarry (locality 6 on Text fig.1), which according to Buckman and Wilson (1896, p.692) showed a very similar sequence to that of the South Main-road quarry. However, as early as the end of the nineteenth century, this guarry had gone 'out of work' and become obscured, (Buckman, 1901). The old road-cutting close to the above section has yielded similar faunas from the upper Elton Farm Limestone, (Buckman, 1901, p.158), and these have found their way into the Bristol University Geological collections. More recently (1977) a fresh cutting has been produced during the widening of the road, and this has confirmed the accuracy of Buckman and Wilson's section (1896, p.692). There have been several exposures on the hill slope to the west of Dundry village, from below the church, past Clement's Yard, (op. cit., p.679) to Grove Farm. The main beds, the ovalis bed and the Grove Farm Limestone appear to have been identical to those exposed at the Grove Farm and Barns Batch exposures. To the east, exposures in the Elton Farm Limestone have been recorded from East Dundry village, (op. cit., 1896, p.682) and from slightly further east at Walnut Farm. A shallow roadside quarry at this latter locality (number 7 in Text fig.1), has produced several ammonites, now in the Bristol University collections, including a rare specimen of Mollistephanus (BUGM. 3338). Beyond Walnut Farm the Aalenian/Lower Bajocian rocks are rapidly overstepped by the Maes Knoll Conglomerate, until they disappear from the field rubble about a half a mile east of East Dundry.

4. THE AALENIAN/BAJOCIAN AMMONITE FAUNAS FROM DUNDRY

AND

The stratigraphic distribution of the main ammonite genera within the Dundry Inferior Oolite is shown in Text fig.2, whilst the distribution of the more important ammonite species found at the South Main-road quarry has already been published elsewhere (Parsons, 1974, fig.4). Whilst these two diagrams give the main pattern of the ammonite distributions, a more detailed appraisal of the ammonite faunas from Dundry, and their relationship to other assemblages from Southern England is given below, starting from the base of the Aalenian rocks.

(a) The murchisonae Zone

The lowest ammonites found <u>in situ</u> during the course of this work were <u>Ludwigia</u> spp. from the basal half of the Barns Batch Limestone, (bed 2, section c). These specimens are not a typical of the <u>murchisonae</u> Subzone of the <u>murchisonae</u> Zone, although a slightly greater age, the <u>haugi</u> Subzone, (= <u>Ancolioceras</u> hemera, S. Buckman, 1910), is possible. The ammonites from the upper Barns Batch Limestone, the <u>Pleurotomaria</u> bed, are more definitive, as the combination of bullate ribbed <u>Ludwigia</u> (<u>L</u>. (<u>L</u>.) <u>crama</u>) and early <u>Brasilia</u>, (7<u>B</u>. (?<u>B</u>.) <u>wilsoni</u>), is characteristic of the <u>murchisonae</u> Subzone <u>sensu stricto</u>, as with the Paving bed of north Dorset, (Buckman, 1893, p.485). There must be a considerable stratigraphic break above this horizon, as no representatives of the <u>bradfordensis</u> Subzone have been found. The one previous record of <u>'Lioceras</u> cf. <u>bradfordensis</u> (Buckman & Wilson, 1896, p.677) is based on the misidentification of a specimen of ?<u>Ancolioceras</u> 5p. (Buckman m.s. note in his own copy of the latter paper).

(b) The concavum Zone

The graphoceratid fauna from the lower part of the Grove Farm Limestone is not well preserved, but it is comparable with the <u>concavum</u> Zone fauna from the lower half of the Bradford Abbas "fossilbed", Bradford Abbas, near Sherborne, north Dorset, (Parsons, 1974, pp.170-1).

(c) The <u>discites</u> Zone

The graphoceratids from this horizon are also not very well preserved, but they are of interest, since they come from the thickest, ammonite rich development of <u>discites</u> Zone rocks in England. The Lower-Trigonia-grit of the Cotswolds, although often thicker, never yields as many ammonites as the equivalent Dundry beds. As yet it has proved impossible to demonstrate any stratigraphic subdivision of the Dundry <u>discites</u> faunas. This is disappointing, as subzonal division of this Zone has been suggested for equivalent horizons in Germany, (Bayer, 1969, p.35), so perhaps further collecting from these beds is necessary. The main ammonite fauna collected from this horizon (bed 6, section C), is exactly equivalent to those collected from the top of the Bradford Abbas "fossil-bed", (Buckman, 1893) and from the 'Snuff-box' bed of south Dorset, (Parsons, 1972).

(d) The laeviuscula Zone

The <u>ovalis</u> bed of the lower Elton Farm Limestone yields a fauna which is characteristic of the <u>ovalis</u> Subzone of the <u>laeviuscula</u> Zone (Parsons, 1974, p.169). This assemblage, which includes well preserved specimens of <u>Witchellia romanoides</u> (Douville), <u>Emileites liebi</u> (Maubeuge)

and various species of Emileia and Trilobiticeras, (Parsons, 1977, p.116), is very similar to that recorded by Douvillé (1885) from southern France. The one major stratigraphic error made by Buckman and Wilson (1896) in their fine work, concerned the correlation of this bed with the north Dorset succession. The ovalis bed is not to be correlated with the basal half of the Sandford Lane "fossil-bed", Sandford Lane, near Sherborne, Dorset, (Buckman, 1893, p.492) as suggested by Buckman, (Buckman & Wilson, 1896, pp.708-9), but with the sandy limestones found beneath this horizon, (Buckman, 1893, p.493, The fauna which is characteristic of the basal part of the bed 8). Sandford "fossil-bed", including S. (Euhoploceras) acanthera, Shirbuirnia spp. and Mollistephanus spp., comes in fact from a slightly higher horizon; the Limonitic bed, (see Parsons, 1974, pp.166-9, and fig.4, for details of the distribution of ammonites within the laeviuscula Zone), which is thus lower laeviuscula Subzone in age.

The fauna from the <u>Witchellia</u> bed, which includes well preserved members of the <u>Witchellia glauca</u> - <u>falcata</u> group, as well as specimens of <u>Frogdenites</u>, is characteristic of the upper <u>laeviuscula</u> Subzone, like that from the 'Green grained marl' of Oborne, north Dorset, (Buckman, 1893, p.500, bed 9; Parsons, 1976b, p.132, bed 3).

(e) The sauzei Zone

The ammonite fauna collected from the Brown Iron-shot bed, (upper part of the Elton Farm Limestone), is exactly equivalent to that recently collected from the top half of the Sandford "fossil-bed", north Dorset, (Parsons, 1974, p.166; Buckman, 1893, p.492), it is thus <u>sauzei</u> Zone in age. This assemblage, which includes fine specimens of <u>Kumatostephanus</u>, <u>Labyrinthoceras</u> and a variety of species of <u>Emileia</u>, was, during the nineteenth century, one of the most famous and well collected, european ammonite faunas and numerous specimens from this horizon have found their way into most national museum collections.

(f) The humphriesianum Zone

There is a possibility that higher stratigraphic horizons than that of the sauzei Zone may have been preserved in small pockets of sediment beneath the 'Vesulian transgression', at the centre of the gentle syncline between South Main-road quarry and East Dundry. This would not come as a surprise as humphriesianum Zone rocks have been found in a similar situation both in the Sherborne district of north Dorset (Buckman, 1893) and in the 'Cole syncline' of Somerset (Richardson, 1916). Although the presence of humphriesianum Zone rocks has been suggested (Richardson, 1919, p.152), physical evidence for this is slight. Specimens of the brachiopod Striirhynchia dundriensis (S. Buckman), which has been recorded from the humphriesianum Zone, Irony bed of north Dorset (Richardson, 1932, p.69), have been found in beds of a slightly later date than the Brown Iron-shot bed, at the North Main-road quarry, Dundry (Tutcher in Kellaway & Wilson, 1941, p.153). This tenuous evidence is supported by the occurrence of the holotype of Oppelia subradiata (J. de C. Sow.), which is purported to have come from Dundry. If this is so, then this ammonite is strong evidence for the occurrence of humphriesianum/subfurcatum rocks on Dundry, as this species in only known in situ from this later horizon (Sturani, 1971, p.114). However, before any positive statements can be



Fig. 2. A stratigraphic distribution of those Aalenian Bajocian ammonite genera and subgenera so far recorded from Dundry Hill.



Fig. 3. A correlation of the four must important exposures of the Dundry Interior Oolite.

made on the presence of rocks of this age on Dundry, further evidence, provided by the <u>in situ</u> collection of ammonites, is needed.

(g) The garantiana Zone

The few ammonites which have been collected from the Maes Knoll Conglomerate, including <u>Garantiana</u> sp., (Buckman & Wilson, 1896, p.685) and <u>Parkinsonia rarecostata</u> S. Buckman (BUGM.3397), indicate a correlation with the upper <u>garantiana</u> Zone, <u>acris</u> Subzone; it is thus the same age as the Astarte bed of south Dorset, (Senior, Parsons & Torrens, 1970).

(h) The parkinsoni Zone

A solitary specimen of <u>Parkinsonia</u> has been collected from the Dundry Freestone, (BUGM.3458, <u>ex</u>. T. Fry collection, from ... "3"-4" above the best beds, Dundry Down"), which is most closely related to the <u>P. parkinsoni</u> (Sow.) group. The most likely correlation for this horizon is with the <u>truellei</u> Subzone of the <u>parkinsoni</u> Zone, as with the <u>truellei</u> bed of south Dorset, (Senior, Parsons & Torrens, 1970, p.115).

5. CORRELATION OF THE DUNDRY INFERIOR OOLITE

The lateral correlation of the exposures described here is shown in a diagrammatic form in Text fig.3. Virtually all the separate lithostratigraphic units described from these sections were found to be isochronous. In this connection the bed, here called the <u>ovalis</u> bed, as previously noticed by Buckman and Wilson (1896, pp.678, 681), was found to be particularly valuable as a marker horizon over all the hill west of East Dundry. Similarly the <u>Pleurotomaria</u> bed, with its very flat, limonite-encrusted top, is another good datum level.

Where ammonites are abundant stratigraphic comparison with other, equivalent beds in Southern England is much simplified; hence all the beds below the Maes Knoll Conglomerate are easily equated with their lateral equivalents. The detailed correlation of these beds with the Dorset succession has largely been given in the previous section, thus only the Cotswold and Bruton districts need further attention.

(a) The Cotswolds

The recent revised dating of much of the Cotswold Inferior Colite (Parsons, 1976a & 1980), provides a basis for the detailed correlation of these beds with the Dundry sequence. It is evident that the Cotswold, Hartley Hill Formation is almost the exact lateral equivalent of the Elton Farm Limestone. The most exact correlation is between the "Witchellia Grit" and the Witchellia bed, both of which have very similar ammonite faunas. However, the Cotswold Lower-Trigonia-grit has yielded discites Zone ammonite faunas, it may thus be correlated with either the base of the Elton Farm Limestone, or with the top of the Grove Farm Limestone. There are greater problems in attempting any detailed correlation of the Cotswold Cheltenham Formation (= pars Lower Inferior Oolite, Mudge, 1978). The Scottsquar Hill Limestone has produced several specimens of Brasilia, which suggests a correlation with the bradfordensis Subzone of the murchisonae Zone, a horizon which does not seem to be represented on Dundry. Ammonites are relatively rare in the Frocester Hill Oolite - Devil's Chimney Oolite sequence, but it does appear to be the lateral equivalent of at least part of the Barns Batch Limestone.

To the south of Dundry all beds of the above age (Aalenian/Lower Bajocian) disappear, and only representatives of the Stroud Formation ("Upper Inferior Oolite") are now preserved. The Maes Knoll Conglomerate is undoubtedly to be correlated with the Upper-Trigonia-grit to the north and part of the Doulting Ragstones to the south (Parsons, 1975). The lack of ammonites from directly above this horizon makes further correlation difficult. The Dundry Freestone is most likely basal <u>parkinsoni</u> Zone in age as is the Doulting Freestone (Parsons, 1975). If this is so, then there is no exact Cotswold equivalent to these beds, as the <u>Clypeus</u>-grit appears to be mainly upper <u>parkinsoni</u> Zone in age (Parsons 1976a, p.62).

(b) The Bruton district

To the south of the Mendips, Lower Bajocian/Aalenian rocks first make their re-appearance in the 'Cole syncline' of the Bruton district, Somerset (Richardson, 1916). This area is the nearest to Dundry to preserve an ammonite rich development of rocks of this age, and the two faunal sequences are very similar. The previous description of the Bruton sections by Richardson (1916), incorporated Buckman's ammonite identifications. The consequent confused dating of these exposures, often with the occurrence in the same bed of ammonites said to be representative of several different hemerae (cf. Richardson, 1916, p.497, bed 10) is a result of the artificial nature of many of Buckman's hemerae, particularly those based on the Sandford Lane "fossil-bed" (Parsons, 1974, pp.162-4; 1976a, p.47). A revision of these important ammonite faunas, based on recent <u>in situ</u> collections from the most complete and presently accessible section; the Bruton railway-cutting (Richardson, 1916, p.495); is thus essential before any detailed correlation can be made with Dundry.

The 'Pecten bed' (Richardson, 1916, p.495, beds 3-4), which has yielded: Emileia (E.) brocchii, E. (E.) contrahens Buckman, E. (E.) polyschides, E. (Otoites) cf. douvillei, E. (O.) aff. fortis (West.), Trilobiticeras (Emileites) liebi, Mollistephanus cf. mollis, Sonninia (S.) aff. propinguans, S. (?S.) aff. carinodisca (Qu.), S. (?S.) straeleni Maubeuge, S. (Euhoploceras) acanthera, Shirbuirnia aff. trigonata, Witchellia (W.) cf. albida, W. (W.) aff.connata, W. (W.) cf. glauca, W. (W.) laeviuscula, W. (Pelekodites) sp., Lissoceras semicostulatum: is of lower laeviuscula Subzone age, and is thus to be correlated with the Dundry Limonitic bed. Beds 4a and 6 (Richardson, 1916, p.495) have yielded: W. (W.) aff. connata and Sonninia (?S.) sp. (bed 4a); and W. (W.) cf. romanoides, S. (S.) sp., S. (Fissilobiceras) aff. fissilobata, S. (F.) cf. ovalis, S. (F.) aff. subtrigonata (Gillet non Buckman), T. (Trilobiticeras) cf. cricki, from bed 6: they are thus of ovalis Subzone age, as with the Dundry ovalis bed. The 'Ammonite bed' (Richardson, 1916, p.495, bed 8) has yielded from the top: S. (Euhoploceras) substratum Buckman, Docidoceras cf. planulatum, Hyperlioceras (H.) cf. subdiscoideum Buckman, H. (H.) cf. liodiscites Buckman: and from the base: Graphoceras (G.) apertum (Buckman), G. (G.) cf. formosum (Buckman), G. (G.) sublineata (Buckman), G. (Ludwigella) compactum (Buckman), G. (L.) compressum (Buckman), H. (H.) subleve Buckman: and thus spans the discites Zone and part of the concavum Zone, as does the Grove Farm Limestone. The basal Conglomerate Bed (Richardson, 1916, p.495, bed 9), is of murchisonae Zone age (Richardson, 1916, p.501) and is thus at least in part the equivalent of the Earns Batch beds.

204

6. THE HORIZON OF/SOWERBYS' AND BUCKMAN'S TYPES

THE

Very few of the ammonite species figured from Dundry have had any accurate stratigraphic information associated with them. Whilst Buckman went to some effort in an attempt to trace the original horizons of the Sowerbys' types, (Buckman & Wilson, 1896, p.701, Table V), he himself made the same mistake of figuring over twenty ammonite species from Dundry, of which only a very small minority were collected <u>in situ</u> from known beds. Thus often very little is known of the type horizon's of many Dundry ammonites, and any information on this point must be garnered from the matrix of the type specimen where this is sufficiently characteristic, and from the horizon of any subsequently collected topotypes.

(a) / Sowerbys' Type specimens

i. <u>Sonninia corrugata</u> (J. de C. Sow.) has a type specimen which is too small for positive identification, (Buckman & Secretary, 1908, Pl. vi, figs.4a,b), although its densely 'iron-shot' matrix points to the Brown Iron-shot bed as its type horizon. Similar specimens to the type have been collected from this latter bed, but the holotype is much too small to be certain if Buckman's larger figured specimen, (Buckman, 1909-30, Pl.412a) is in fact con-specific.

ii. <u>Sonninia browni</u> (J. Sow.) again has a type specimen which is too
small and incomplete for accurate identification, (Buckman & Secretary,
1909, Pl.vi, figs.5a,b), and the lack of adherent matrix makes it
impossible to even guess at the original horizon.

iii. Sonninia sowerbyi (J. Sow.) is more fortunate, since although its

type specimen is again rather fragmentary (Buckman, 1904, Pl.52), the densely 'iron-shot', purple stained matrix points to the Brown Iron-shot as its type horizon. This specimen probably thus represents the inner whorls of a species of <u>Papilliceras</u>, (Parsons, 1974, pp.160-1), and is probably conspecific with <u>S</u>. (<u>P</u>.) <u>mesacantha</u>.

iv. <u>Witchellia laeviuscula</u> (J. de C. Sow.) is perhaps the most easily recognisable of the Sowerbys' species from Dundry, since it has a relatively complete type specimen, which in turn has a distinct matrix. The <u>Witchellia</u> bed is the type horizon for this species, and it has produced numerous topotypes from the South Main-road quarry.

v. <u>Emileia brocchii</u> (J. Sow.) is restricted as a species to the forms closest to the lectotype, (Buckman, in Buckman & Wilson, 1896, p.701), whilst the smaller syntype (J. Sowerby, 1818 in 1812-46) can now be referred to the species <u>Emileites liebi</u> (Maubeuge), specimens of which are common in the <u>ovalis</u> bed, the undoubted source of Sowerby's specimen.

vi. <u>Normannites braikenridgei</u> (J. de C. Sow.) has a fine type specimen, (Buckman, 1909-30, Pl.81), with a densely 'iron-shot' matrix and this together with the topotype specimens recently collected, would indicate the Brown Iron-shot bed as the type horizon.

vii. <u>Otoites contractus</u> (J. de C. Sow.) must be considered a rather dubious species, since owing to the absence of the original type specimen, there is now considerable uncertainty over both its interpretation and type horizon. Westermann's invalid neotype of this species (Westermann, 1954, pp.88-94, Pl.1, figs. 4a,b,c), which is only doubtfully con specific, comes from the wrong locality and is possibly from a different horizon.

 TABLE 4 The probable type horizons of those Dundry ammonite species figured by Buckman (1909-30) in 'Type Ammonites'.

 T = horizon of topotype

 ? = probable type horizon, from the matrix of type.

			Volume	Plate number	Bed at South Main-road Quarry			
					105	10a -	9	8
Sonninia corrugata (Sow.) S. corrugata Dundryites albidus Pelekodites pelekus Spatulites spatians Macerites aurifer Cadomoceras carinatum C. ellipticum Hebetoxyites hebes Leptostrigites languidus Varistrigites compressus Toxamblyites arcifer Emileites malenotatus Emileia subcadiconica Frogdenites profectus Labyrinthoceras amphilaphes L. extensum L. gibberulum Oerostenberus dolibocents			4 6 6 4 7 7 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	412 412A 687 399 765 766 456 455 475 477B 468 473 702 711 430 279 214 278 265	T ? T T ?	T T ? ?		T T T T T

١

viii. <u>Oppelia subradiata</u> (J. de C. Sow.) has a problematic type specimen. The densely 'iron-shot! matrix of this type (Arkell, 1951-9, Text fig.11, 1a,b) would indicate a provenance from the Brown Iron-shot bed, however no topotypes have subsequently been collected of this species, which is more characteristic of the <u>humphriesianum</u> Zone, rather than the <u>sauzei</u> Zone, in the Sherborne area of north Dorset.

(b) Buckman's specimens

S.S. Buckman described and figured a large number of Dundry ammonites in two main publications; his monograph (Buckman, 1887-1907) and 'Type Ammonites', (Buckman, 1909-30). Very few Dundry ammonites in fact appeared in the 'Monograph on the Inferior Oolite Ammonites', and these are discussed individually in the text. The larger number of Dundry specimens figured in 'Type Ammonites', have mostly been traced to their type horizons and the original source of these specimens is shown in Table 4. The question marks (?) in this table refer to type specimens of which either no topotypes have been collected or which have a rather doubtful matrix. An individual discussion of these specimens would take up too much space, and only the more important specimens and groups will be mentioned here.

i. <u>W. (Pelekodites) dundriensis</u> (Buckman) has a type specimen (Buckman, 1887-1907, Pl.23, figs.5 & 6), which Buckman thought came from the 'Iron-shot bed' of Dundry. However this specimen is so close to <u>W. (P.) macra</u> (Buckman), that it would seem unlikely that it originated from any bed other than the <u>Witchellia</u> bed, where this latter species is common. For further details see the subsequent discussion of the <u>W. (Pelekodites</u>) group.

<u>W.</u> (P.) aff. <u>buckmani</u> (Haug) - (Buckman, 1887-1907, Pl.23, figs.
7 & 8), undoubtedly came, as stated, from the Brown Iron-shot bed,
where forms allied to this species are quite common.

iii. <u>Ludwigia obtusa</u> (Buckman, 1887-1907, Sup. Pl.4, fig.10 <u>ex</u>. Qu.), would appear from its matrix to have come from the <u>Pleurotomaria</u> bed of the Barns Batch Limestone. Similar specimens of this species have been found in this and subjacent beds at Barns Batch Spinney, (section c, beds 2 and 3).

iv. ?<u>Brasilia wilsoni</u> (Buckman); the paratype of this species (Buckman, 1887-1907, Sup. Pl.12, fig.7) came from the <u>Pleurotomaria</u> bed, from which horizon several closely related forms have recently been collected.
v. Oedania parvicostata Buckman (1887-1907, Sup. Pl.21, figs.7-9)

v. <u>Oedania parvicostata</u> Buckman (1887-1907, Sup. Pl.21, figs.7-9) and

vi. <u>Braunsina projecta</u> Buckman (<u>op. cit.</u>, Sup. Pl.20, figs.7-9) are forms characteristic of the <u>discites</u> Zone beds of north Dorset, although only a topotype of the latter species has as yet been collected from the upper Grove Farm Limestone.

vii. <u>Frogdenites</u> is a small sphaeroceratid ammonite genus which so far has only been recorded once from Dundry, (<u>F. profectus</u>, Buckman, 1909-30, Pl.430). However a close study of several specimens from Dundry figured by Buckman as new species of <u>Labyrinthoceras</u>, (<u>L. extensum</u>, Buckman, 1909-30, Pl.214; <u>L. gibberulum</u>, op. cit., Pl.278), reveals their true identity as members of the <u>Frogdenites</u> group. Careful collection of the north Dorset exposures of the Inferior Oolite has shown that <u>Frogdenites</u> has a stratigraphic range restricted to the top of the <u>laeviuscula</u> Zone (Parsons, 1974, p.167); no members of this genus have been collected from the <u>sauzei</u> Zone layer of the Sandford Lane "fossil-bed", (Buckman, 1893, p.492, bed 6). Those members of this genus recorded from Dundry are thus most likely to have originated from the <u>Witchellia</u> bed, rather than the Brown Iron-shot. Collection <u>in</u> <u>situ</u> from the <u>Witchellia</u> bed of the South Main-road quarry (Section a, bed 10a) produced one topotype of <u>F. gibberulum</u> (Buckman), thus confirming this bed as the type horizon for Buckman's specimen. Buckman relied heavily upon slight differences in matrix for the placing of his specimens of <u>Labyrinthoceras</u> and <u>Frogdenites</u> in their correct stratigraphic position. Since the differences in matrix between the Brown Iron-shot and <u>Witchellia</u> beds can be slight to non-existent, until more topotypes come to light, the <u>Witchellia</u> bed should be considered the type horizon of all Buckman's specimens of <u>Frogdenites</u>.

viii. <u>Labyrinthoceras</u> is a genus closely related to the preceding, from which it evolved, and which is characteristic of the <u>sauzei</u> Zone in north Dorset. The specimens of this genus from Dundry; <u>L. amphilaphes</u> Buckman (1909-30, Pl.279) and <u>L. meniscum</u> (Waagen), (BUGM.3289); thus undoubtedly came from the Brown Iron-shot bed, a position fully in accord both with their matrix and other specimens of this genus subsequently collected in situ.

ix. <u>Pelekodites</u> is the small microconch subgenus of the genus <u>Witchellia</u>, both of which range from the base of the <u>ovalis</u> Subzone to the top of the <u>sauzei</u> Zone. Five specimens of this subgenus were figured by Buckman from Dundry, of which four were as new species. However, there are only three successive populations to which these different specific names may be applied.

1. The <u>ovalis</u> Subzone is characterised by the <u>Witchellia</u> (<u>W.</u>) <u>romanoides</u> (Douvillé) macroconch group, associated with which there are specimens of <u>W.</u> (<u>Pelekodites</u>) <u>macra</u> (Buckman) and <u>W.</u> (<u>P.</u>) <u>pelekus</u> (Buckman).

2. The <u>laeviuscula</u> Subzone has a more varied <u>Witchellia</u> fauna, which naturally is reflected in its microconch population. <u>W</u>. (P.) <u>macra</u> (including sub. syn. <u>P. dundriensis</u>, <u>P. costulatus</u>, and <u>P.</u> <u>aurifer</u> Buckman) is still present along with the largest member of this subgenus, P. spatians (Buckman).

3. The <u>sauzei</u> Zone has the last members of this subgenus, which are typically inflated and coarsely ribbed as in <u>W</u>. (<u>P</u>.) <u>sulcata</u> Buckman, (including sub. syn. <u>P</u>. <u>buckmani</u> (Haug), <u>P</u>. <u>zurcheri</u> (Douv.) and <u>P</u>. <u>schlumbergeri</u> (Haug)), and which represent the microconch counterparts of the <u>W</u>. (<u>W</u>.) <u>hebridica</u> Morton group.

The horizons of the topotypes from Dundry of the Buckman species are shown in Table 4.

ACKNOWLEDGEMENTS

The main part of this work was undertaken during the tenure of a University of Keele Research Studentship, at that University; this award is gratefully acknowledged. I should also like to thank the various individuals who have given help during the course of this work; Drs. H. Torrens, J. Callomon and J. Whicher; as well as the various museums who have allowed me access to material in their care, notably, the British Museum (N.H.) and Institute of Geological Sciences, London, the Bristol City and Bristol University, Department of Geology, Museums, Bristol and the Oxford University Museum.
REFERENCES

Arkell, W.J., 1933, (Reprinted, 1970), The Jurassic System in Great Britain. Oxford University Press; 681 pp. 41 Pl.

1951-9, A monograph of the English Bathonian Ammonites. Palaeontogr. Soc.; 264 pp. 33 Pl.

Bayer, U., 1969, Die Gattung <u>Hyperlioceras</u> Buckman. Jber. u. Mitt. oberrh. geol. Ver. 51 : 31-70.

Buckman, S.S., 1887-1907,

A Monograph of the Ammonites of the Inferior Oolite Series. Palaeontogr. Soc. : cclxii + 456 pp. 127 Pl.

1889.

The relation of Dundry with the Dorset-Somerset and Cotteswold Areas during part of the Jurassic Period. Proc. Cotts. Nat. F. Club <u>9</u>: 374-87.

1893, The Bajocian of the Sherborne district; its relation to subjacent and superjacent strata. Quart. J. geol. Soc. Lond. <u>49</u>: 479-522.

1901, Excursion to Dundry Hill. Proc. Geol. Assoc. <u>17</u>: 152-8.

1904, <u>Ammonites sowerbyi</u> (Miller m.s.) J. Sowerby 1821. Palaeontologia Universalis Pl. 52.

1909-30, 'Type Ammonites' London & Thame, vols. 1-7; 790 Pls.

______1910, Certain Jurassic (Lias-Oolite) strata of South Dorset. Quart. J. geol. Soc. Lond. 66 : 52-89.

. & 'Secretary', 1908, Illustrations of Type Specimens of Inferior Oolite Ammonites in the Sowerby Collection. Palaeontogr. Soc. ; 7 Pls. Buckman, S.S., & Wilson, E., 1896,

Dundry Hill : its upper portion, or the beds marked as Inferior Oolite in the maps of the Geological Survey. Quart. J. geol. Soc. Lond. 52 : 669-720.

1897,

The geological structure of the upper portion of Dundry hill. Proc. Bristol Nat. Soc. 8: 188-231.

Cleak, D.S., 1930, The lithology of the Inferior Oolite, Dundry, (North Somerset). Geol. Mag. <u>67</u>: 217-21.

Donovan, D.T., 1958, Easter field meeting : the Lower and Middle Jurassic Rocks of the Bristol District. Proc. Geol. Assoc. <u>69</u> : 130-40.

Donovan, D.T., & Hemingway, J.E., 1963, Lexique Stratigraphique International; Fascicule 3a, Part 3aX, Jurassic. Paris : 394 pp.

Douvillé, H., 1885, Sur quelques fossiles de la zone à <u>Ammonites sowerbyi</u> des environs de Toulon. Bul. Soc. géol. Fr. (3) <u>13</u> : 12-44, 3 Pl.

Gatrall, M., Jenkyns, H.C. & Parsons, C.F., 1972, Limonitic concretions from the European Jurassic, with particular reference to the 'snuff-boxes' of southern England. Sedimentology 18 : 79-103.

Harland, W.B. (Chairman) et al., 1972, A concise guide to stratigraphical procedure. Jl. geol. Soc. <u>128</u>: 295-305.

Kellaway, G.A. & Wilson, V., 1941, An outline of the Geology of Yeovil, Sherborne and Sparkford Vale. Proc. Geol. Assoc. <u>52</u>: 131-174.

Macfadyen, W.A., 1970, Geological Highlights of the West Country. Butterworths, London; 296 pp.

Mudge, D.C., 1978, Stratigraphy and sedimentation of the Lower Inferior Colite of the Cotswolds. Jl. geol. Soc. Lond. <u>135</u>, 611-27

Oppel, A., 1856-8, Die Juraformation Englands, Frankreichs und des sudwestlichen Deutschlands. Jb. Ver. vaterland. Naturkde. Wurtt. (Stuttgart), 12-14 : 1-837. Orbigny, A.D., 1842-51, Paleontologie Française; Terrains Jurassiques, I Cephalopodes. Paris, 642 pp. 234 Pls. Parsons, C.F., 1972, A new temporary section in the Inferior Oolite of South Dorset. Proc. Dorset Nat. Hist. Archaeol. Soc. 93: 117-8. 1974. The sauzei and 'so called' sowerbyi Zones of the Lower Bajocian. Newsl. Stratigr. 3 : 153-79. 1975, Ammonites from the Doulting Conglomerate Bed (Upper Bajocian, Middle Jurassic), Doulting, Somerset. Palaeontology 18 : 191-205. 1976a, Ammonite evidence for dating some Inferior Oolite sections in the north Cotswolds. Proc. Geol. Ass. 87, 45-63, 2 Pls. 1976ъ. A stratigraphic revision of the humphriesianum/subfurcatum Zone rocks (Bajocian Stage, Middle Jurassic) of Southern England. Newsl. Stratigr., 5, 114-42. 1977. Two new Bajocian microconch Otoitid ammonites and their significance. Palaeontology, 20, 101-18, 1 Pl. 1980. A stratigraphic revision of the Bajocian (Jurassic) rocks of the Cotswold Hills. In Press. Chapter from Unpublished Ph.D. Thesis University of Keele, 1980, pp.215-254. Reynolds, S.W. et al., 1930, The Geology of the Bristol district. British Assoc. Advance. Science. Bristol : 59 pp. Richardson, L., 1907, The Inferior Oolite and Contiguous Deposits of the Bath-Doulting District. Quart. J. geol. Soc. Lond. 63 : 383-423.

Richardson, L., 1916,

The Inferior Oolite and Contiguous Deposits of the Doulting-Milborne Port district (Somerset). Quart. J. geol. Soc. Lond. 71: 473-520.

1919, The Inferior Oolite and Contiguous Deposits of the Crewkerne District (Somerset). Quart. J. geol. Soc. Lond. <u>74</u>: 145-73, Pl. xiv-xvi.

1932, The Inferior Oolite and Contiguous Deposits of the Sherborne District, Dorset. Proc. Cotts. Nat. Field. Club 24 : 35-85, Pl. vi-xi.

Senior, J.R., Parsons, C.F. & Torrens, H.S., 1970, New sections in the Inferior Oolite of south Dorset. Proc. Dorset Nat. Hist. Archaeol. Soc. <u>91</u>: 114-9.

Sowerby, J. & Sowerby, J. de C., 1812-46, 'The Mineral Conchology of Great Britain' London: 1287 pp. 648 Pl.

Sturani, C., 1971, Ammonites and Stratigraphy of the '<u>Posidonia alpina</u>' beds of the Venetian Alps. Mem. Ist. Geol. Mineral. Padova <u>28</u>: 190 pp. 16 Pl.

Tutcher, J.W., 1903, The Lower Oolites near Bristol. Proc. Bristol Nat. Hist. (N.S.) 10 : 150-168.

1929. Excursion to Dundry and Radstock. Proc. Geol. Assoc. 40: 174-6.

Westermann, G.E.G., 1954, Monographie der Otoitidae. Beih. Geol. Jb. <u>15</u> : 363 pp. 33 Pls.

Wilson, E., 1893,

Excursion to Dundry Hill. Proc. Geol. Assoc. 13: 128-132.

Wright, T., 1860.

On the subdivisions of the Inferior Oolite in the South of England, compared with the Equivalent Beds of that Formation in Yorkshire. Quart. J. geol. Soc. Lond. 16 : 1-48.

2 B. A STRATIGRAPHIC REVISION OF THE BAJOCIAN (JURASSIC) ROCKS

OF THE COTSWOLD HILLS

1. Sugar She

C.F. Parsons

i de la ferrar de la companya de la

승규는 영국 문

的现在分词是是是自己的意思。在这些主义的是是

SUMMARY: A formal scheme of lithostratigraphic subdivisions for the Inferior Oolite 'series' is erected. The Inferior Oolite Group of the mid-Cotswolds is defined as being made up of three Formations; the Cheltenham (= 'Lower Inferior Oolite'), Hartley Hill (= 'Middle Inferior Oolite') and Stroud (= 'Upper Inferior Oolite'); with an additional fourth Formation (the Stanway Hill Formation, with Harford Sands and Snowshill Clay Members), in the north Cotswolds. Three Members of the Hartley Hill Formation are re-defined or newly named; the Lower-Trigoniagrit, Windrush and Notgrove Members; whilst the existing informal units. the Clypeus-grit and Upper-Trigonia-grit are redefined as formal Members of the Stroud Formation. Bajocian ammonite faunas from the Cotswolds are listed (+ 100 specimens) and discussed, whilst the correlation of the ·【本われ語句》(本語文語)(注意)) Upper Bajocian rocks of the south Cotswolds is discussed and compared COMP. B. THERE AND AND A REPORT OF A DESCRIPTION OF with the successions in the Doulting and Dundry districts. Caral and the second states of

的一种分析是在大学的影响。这些是是是是是

the Cold Cold and the American sector and

The Inferior Oolite rocks (Aalenian-Bajocian Stages, Middle Jurassic) of the Cotswold Hills are superbly displayed in a series of natural scarps and artificial exposures created by the exploitation of their famous building stones. This wealth of exposure, together with the abundance of fossils led several generations of geologists to turn their attention to the stratigraphic problems of these rocks. In broad terms the Inferior Oolite of the Cotswolds may be divided into two divisions, an upper series of mainly bioclastic limestones (the Upper and Middle Inferior Oolite sensu Arkell, 1933) of Bajocian age and a lower group of predominantly colitic limestones and marls (Lower Inferior Oolite) of Aalenian age. This simple grouping was recognized at an early stage by workers such as Lycett (1850), who included the upper beds within his 'Ragstones' and the lower beds within the 'Freestones'. The aim of this work is to revise the detailed bio- and litho-stratigraphy of the rocks of Bajocian age (the 'Ragstones'), since the Aalenian beds have been the subject of a recent work (Mudge, 1978).

The existing informal lithostratigraphic subdivisions of the Cotswold Inferior Oolite were introduced haphazardly over a period of more than one hundred years, hence they tend to be both poorly defined and ambiguous in use. A rationalization of these units in order to produce a standardized hierarchical scheme, in line with current practice (Holland <u>et al.</u>, 1978) is thus long overdue. In particular a more rigorous definition of the lithostratigraphic units is required. In the past there has been a tendency to place more emphasis on the biostratigraphic value of the constituent faunas of a unit, such as the <u>Witchellia</u> ammonites of the <u>Witchellia</u> Grit^{*} than on its lithology and value in field mapping. This

ZONE	SUBZONE
Parkinsonia parkinsoni	<u>P. bomfordi</u> Strigoceras truellei
	P.acris
<u>Strenoceras</u> (<u>Garantiana</u>) <u>garantiana</u>	<u>S.(G.) tetragona</u> <u>S.(G.) subgaranti</u>
	S.(Pseudogarantiana) dichotom
<u>S.(Strenoceras)</u>	<u>Caumontisphinctes</u> (<u>C</u> .)
subfurcatum	<u>polygyralis</u> <u>Teloceras</u> b <u>anksi</u>
<u>Stephanoceras</u> (S.) humphriesianum	<u>T.blagdeni</u> <u>S.(S.) humphriesianum</u> <u>Dorsetensia romani</u>
Emileia (Otoites) sauzei	
<u>Witchellia (M.) laeviuscula</u>	<u>W.(W.) laeviuscula</u> Sonninia ovalis
Hyperlioceras (H.) discites	

Table 1.

UPPER BAJOCIAN

LOWER BAJOCIAN

Zones and subzones of the Bajocian Stage (excl. Aalenian), modified after Parsons (1974 & 1976).

has led, particularly in the 'Middle Inferior Oolite', to the use of what in some cases are little more than biostratigraphic units, which do not warrant formal recognition in a hierarchical, lithostratigraphic scheme. If lithological homogeneity and ease of mapping are to be taken as the main criteria for establishing the Cotswold Inferior Oolite Formations and Members (Holland, et al., 1978, p.8), then in many cases only a more limited number of formal units can be recognized. However, this approach does not prevent the continued informal use of lithological units, which have not been incorporated in the formal scheme. A revision of the biostratigraphy has become necessary because of the recent work which has been undertaken on the Bajocian rocks of Dorset and south It was during the course of his work in these latter districts Somerset. that S. Buckman (1893), laid down the biostratigraphic framework, which he was to use to elucidate the stratigraphy of the Cotswolds. It is now evident that Buckman's scheme of hemerae (approx. = subzone in present usage), which he used for this work, contains contradictions and at least one major error, which subsequently became incorporated in the Bajocian Standard Zonal scheme (Spath, 1936; Arkell, 1956). Some new insight. which is now available into the ammonite faunas of Dorset (Parsons, 1974, 1976), has enabled a more accurate correlation to be made of the Cotswold deposits. The previous Zonal scheme for the Lower Bajocian (Middle Bajocian olim) has been found to contain the greatest inconsistencies and hence most attention has been given to the ammonite faunas of this age. The Upper Bajocian rocks of the south Cotswolds have been treated in some detail, as they appear to have been mis-correlated in the past. Since the Inferior Oolite rocks found just to the south of the Mendips, in the



Figure 1.

A sketch map showing the relative positions of the main exposures mentioned in the text.

Doulting district, are of importance for the understanding of the succession in the south Cotswolds, some details of these are also given.

The revised Zonal scheme for the Bajocian used in this work is shown in Table 1, whilst the general area under discussion and the location of some of the more important exposures is shown in Figure 1.

THE LITHOSTRATIGRAPHY OF THE COTSWOLD INFERIOR OOLITE

The following revision of the lithostratigraphy of the Cotswold Inferior Oolite (see Table 2), is an attempt to implement the stratigraphic procedure recommended by the Geological Society of London (Holland <u>et al.</u>, 1978). As far as is possible I have retained existing well proven terminology within the new hierarchical scheme of Group, Formation and Member. However in some cases, where several existing units have been combined, the use of the earliest available name, such as the Gryphite Grit, for the combined <u>L. buckmani</u>/Gryphite Grits, would only create confusion and ambiguity. In these circumstances new formal names have been introduced.

The Inferior Oolite Group

The mid-Cotswold Inferior Oolite is a relatively thick deposit (60 m.+ at Leckhampton) containing various lithologies and facies, which may be divided into three main divisions or Formations. These Formations are broadly similar; they overwhelmingly consist of limestones; and since together they constitute a naturally related association in comparison with the Lias below and the Great Oolite above, it is logical to



Table 2.

A revised lithostratigraphic scheme for the Inferior Oolite Group in the mid and north Cotswolds, showing its relationship to equivalent beds in Southern England.

combine them in a single formal Group. In deference to existing usage, this has been called The Inferior Oolite Group (McKerrow and Kennedy, 1973, p.2).

The existing informal divisions of the Inferior Oolite 'Series' into Upper, Middle and Lower Inferior Oolite (Arkell, 1933), although similar in their boundaries to three of the Formations proposed here, can never have any formal status (Holland, <u>et al.</u>, 1978, pp.10-1). Firstly these terms have been used with a variety of different meanings (see below), thus even after formal re-definition, the possibility of ambiguity would exist. Secondly, it has been recommended, that when an established unit is broken down into two or more formally defined formations, the existing name should only either be raised to group status, or abandoned (Hedberg, 1976, p.44). Since the former course has been taken, new formational names become necessary.

The limits of the currently used divisions derive from Buckman, who was the first to recognize the two major unconformities which naturally subdivide the Cotswold Inferior Oolite (Buckman, 1897, and 1901). However, several other alternative subdivisions of the Inferior Oolite have been popular, including a two-fold division into Upper and Lower Inferior Oolite (Woodward, 1894) and yet another three-fold subdivision with Upper ('Ragstones'), Middle ('Freestones') and Lower ('Pea Grit') Inferior Oolite subdivisions (Murchison, 1834), which were commonly accepted throughout much of the nineteenth century (Lycett, 1850; Witchell, 1882). The revised three-fold division of the Inferior Oolite suggested by Cave and Penn, (1972) utilizes yet a different set of vertical limits, with the inclusion of the Tilestone/Snowshill Clay/Harford Sands in the Middle

rather than the Lower Inferior Colite. For a further discussion of the vicissitudes of these various subdivisions of the Inferior Colite, one should refer both to Cave and Penn (1972) and to the Lexique Stratigraphique (Worssam and Donovan, pp.174-6 in Donovan and Hemingway, 1963). There is thus a strong case for establishing formal, named units to stabilize this inconsistent and informal nomenclature. An obvious and apparently logical move, would be to accredit the status of Formation to the majority of the existing lithostratigraphic subdivisions of the Inferior Colite, as has largely been done with the Great Colite (Sellwood and McKerrow, 1974). A step in this direction has been taken with the use of the term'<u>Clypeus</u>-grit Formation' (Kennedy, Sellwood and McKerrow, p.2 in Ager <u>et al</u>., 1973; see also, Murray, 1969). There are however problems associated with this seemingly straightforward approach.

As already noted there are two prominent unconformities with extensive 'hard-grounds' within the Cotswold Inferior Oolite. These unconformities at the base of the Upper and Lower-Trigonia-grits, constitute the major factors controlling the preservation and present distribution of the 'Middle and Lower' Inferior Oolite (Buckman, 1901, fig.3; Arkell, 1933, fig.35). Any new scheme of lithostratigraphic nomenclature must recognize and reflect the stratigraphic importance of these two erosion surfaces. In this connection Arkell (1933, p.231) made the following pertinent comments concerning the uppermost of the two unconformities "Certainly no better plane for separating two Formations occurs in the Jurassic System, and the failure of our classification to take account of it is one of its worst shortcomings" If the existing lithological subdivisions of the Inferior Oolite were to be

considered as Formations, then the only way in which the revised nomenclature could take account of the importance of these unconformities, would be by arranging the constituent Formations into three or more separate Groups. However, this would totally over-emphasise the relative importance of the Inferior Colite in relation to neighbouring Groups, and this approach must be rejected.

The other alternative is to take the two unconformities as the starting points for the establishment of a more 'natural' classification. The components thus delimited can be considered as Formations, with the existing lithological units being re-defined as constituent Members. These Formations can fulfil all the essential criteria for units of this rank (Holland et al., 1978, p.8) and it is this procedure which is followed here (cf. Mudge, 1978). The Inferior Oolite Group of the Cotswolds is thus here defined as being made up of the Cheltenham, Hartley Hill and Stroud Formations, with an additional fourth Formation; the Stanway Hill Formation; in the north Cotswolds. The mid-Cotswold units are clearly distinguished both by their different lithologies and by the disconformities, which separate them throughout much of Southern England. Thus they are readily recognizable to the east in boreholes in the Hampshire/Weald basin (e.g. Warlingham; Worssam & Ivimey-Cook, 1971, pp.45-6). However, south of Stroud the Hartley Hill and Cheltenham Formations are rapidly over-stepped, so that beyond Chipping Sodbury the Stroud Formation is the only representative of the Inferior Oolite Group. When the Aalenian a. Lower Bajocian rocks reappear from beneath the unconformity at the base of the Stroud Formation, in the Bruton district of Somerset (Richardson, 1916). they have lost their marked differences in facies and are not subdivisible

into separate Formations. Here the Upper Bajocian, "Top beds" (op. cit., p.485) are underlain by a group of thin, 'condensed', highly fossiliferous, glauconitic, lenticular deposits, which further south are very ferruginous and 'iron-shot' (Kellaway & Wilson, 1941, p.148, 'group A beds'). The former are undoubtedly a lateral continuation of the Stroud Formation, whilst the latter are the equivalent of the Hartley Hill, Stanway Hill and Cheltenham Formations of the Cotswolds (Parsons, 1979; see Table 2). The lower beds are informally named the 'Oborne beds' in Table 2, after the excellent sections around the Oborne area of Sherborne, and in the Oborne borehole (Wilson <u>et al.</u>, 1959, pp.96-7; $185\frac{3}{4}$ ' - 217'). No formal definition as a Formation is made here as this must await a detailed revision of the relevant beds.

Since my aim has been, where possible, to re-define the existing informal units, few details are given of the area of preservation and lateral variation of the various Cotswold beds described, as these may be obtained from the earlier literature (Buckman, 1895, 1897, 1901; Arkell, 1933).

The Cheltenham Formation

All those beds in the mid-Cotswolds below the Lower-Trigonia-grit and above the base of the 'scissum beds' (= Leckhampton Limestone, Mudge, 1978); the 'Lower Inferior Oolite' of Arkell (1933); are here named the Cheltenham Formation. This unit thus includes all the members defined by Mudge (1978), bar the Harford Sands, which have here been transferred to the Stanway Hill Formation (see below). Leckhampton Hill (National Grid reference S0952185-949186) is here designated as the type locality, where

the Formation predominantly consists of an alternation of massive oolitic limestones and oolitic and pisolitic marls, some 50m thick (Richardson, 1906, pp.187-9; Murray, 1969, pp.541-5; Mudge, 1978a, log.1).

The Stanway Hill Formation

The beds previously known as the 'Tilestone' (Buckman, 1901), 'Snowshill Clay' (Buckman, 1897), 'Harford Sands' (Buckman, 1888) and 'Naunton Clay! (Richardson, 1929), and jointly included in the 'concava beds! (Arkell, 1933) are here transferred to a single, new unit: the Stanway Hill Formation; named after the type section at Jackdaw Quarry (SP078310). Stanway Hill (Parsons, 1976, pp.55-61, beds 4-14). From their original definition (op. cit.), the classification of these beds has proved to be problematic, although most subsequent authors (e.g. Arkell. 1933; Ager, 1956; Green & Melville, 1956; Mudge, 1978), have included them in the 'Lower Inferior Oolite' (= Cheltenham Formation). Doubts as to the precise relationship of these beds to the basal Lower-Trigoniagrit disconformity prompted Cave & Penn, (1972), to include them in the 'Middle Inferior Oolite' (= Hartley Hill Formation). More detailed information on the north Cotswold Inferior Oolite, which has subsequently come to light (Parsons, 1976; Mudge, 1978), has led to the rejection of Cave & Penn's conclusions. However, these problems of classification only help to confirm that these beds are out of place in either of these informal subdivisions.

These beds are a-typical (in facies, original depositional environment and present pattern of distribution) of most of the rest of the

Inferior Oolite Group. The Cotswold Inferior Oolite as a whole, is fully marine and dominated by shallow water, carbonate facies: open shelf, carbonate sands; colite shoals etc. (Mudge, 1978, pp.622-3). In direct contrast the Stanway Hill Formation is dominated by terrigenous clastic material, with abundant clays, shales and fine quartz sands. There is strong evidence that these horizons are not wholly marine. There is intense bioturbation, with common lignite and plant fragments, together with the presence of strongly depleted faunas, with virtually mono-specific beds of Lucina, Liostrea, Pinna etc. (Parsons, 1976, pp.58-9; Mudge, 1978, p.623); all of which point to deposition in a shallow, marginally marine. coastal plain environment (loc. cit.). The present pattern of preservation of this Formation is also anomalous. The majority of the Inferior Oolite is at its thickest in the centres of the Painswick and Cleeve Hill 'synclines' (Buckman, 1897, pl.46; Arkell, 1933, fig.35; Mudge, 1978, figs.1,3 & 5), with an associated thinning north-east towards the Vale of Moreton 'swell'. By comparison, the Stanway Hill beds are absent from the Painswick 'syncline', extremely attenuated in the Leckhampton-Wistley Hill area, and thicken progressively towards the north-east, where they are thickest in the Blockley-Broadway district (Richardson, 1929), at the northern extremity of the present Inferior Oolite outcrop . Taking into account all the above factors, there can be little doubt that these beds merit the status of Formation, especially since they are readily mappable in relation to adjacent Formations.

With their identical stratigraphic positions (pre-<u>discites</u> Zone/ post-<u>murchisonae</u> Zone) it is likely that the present beds are the lateral equivalent of the Grantham Formation (Kent, 1975), of the East Midlands.



Although now disjunct, due to subsequent periods of intra-Bajocian and Bathonian erosion, it would seem likely that these two once formed a contiguous deposit across the 'Oxfordshire Shallows' (Sylvester-Bradley & Ford, 1968, pp.211-2). Whilst it could be possible to apply the East Midland nomenclature to the north Cotswold beds, the separation of the latter on the periphery of a different depositional basin, merits a distinctive name.

The relationship of the Formation to sub- and suprajacent strata is clear cut. It is overlain throughout the north Cotswolds disconformably by the Lower-Trigonia-grit Member (of proven discites Zone age), of the Hartley Hill Formation (Parsons, 1976, p.62). It is underlain by the Scottsquar Hill Limestone Member of the Cheltenham Formation (= 'Upper Freestone /or its equivalents, and 'Oolite Marl'), usually, as at Jackdaw quarry (Parsons, 1976, p.62), with a gradational contact - see Fig.2. The Brasilia sp. recorded from the top of the Scottsquar Hill Member at Westington Hill quarry (op. cit., p.58), is very similar to specimens recently collected from the top of this same unit at the Frith quarry, Painswick (Buckman, 1895, p.400, bed 20). This would suggest that not only is the Scottsquar Hill Member virtually complete throughout much of the north Cotswolds, but that it forms a relatively isochronous horizon. This undoubtedly refutes the suggestion made by Mudge (1978, pp.620-1), that the Harford Sands represent a lateral facies variant of part of the A.新教授·哈尔萨姆的副主任公司 more southerly Scottsquar Hill Member. The precise age of the Stanway Hill Formation is still rather speculative, although its overall strati-这次有点的"金融就会_会"的"路"的"路 graphic position (see above), suggests a correlation with the concavum Zone: an age which is not out of step with the one recorded ammonite (Parsons, 1976, p.51).

The Harford Sands Member

The present member is here restricted to the Harford Sands <u>sensu</u> <u>stricto</u> (Buckman, 1888), excluding the 'Snowshill Clay - Tilestone', included in it by Mudge (1978, p.618). The existing type locality (Harford Cutting : Parsons, 1976, p.48), is unsatisfactory, in that the base of the unit has never been exposed. Hence, Happylands quarry (SP128357), is here designated as a hypostratotype (cf. Hedberg, 1976, p.38), where the complete thickness of the Harford Sands (c.2.6m of orange, silty sands and 'doggers'), is seen underlying the Snowshill Clay, and resting disconformably on the Scottsquar Hill Member (see Fig.2). The Naunton Clay is here redefined as a formal bed within the Harford Sands Member, with Jackdaw quarry as its type locality, where it consists of 0.70m of grey and brown, fossiliferous clay (Parsons, 1976, p.57, beds 4-5).

The Snowshill Clay Member

The original type locality of the informal unit (Cleeve Hill : Worssam & Donovan, p.324 in Donovan & Hemingway, 1963), is unsatisfactory, as it is uncertain that the thin clays present at this locality are in fact contiguous with the thicker deposits found between Snowshill and Blockley. Hence the nearest existing exposure to Snowshill; Happylands quarry; is here formally defined as the type section of the Snowshill Clay Member. Here 1.4m+ of dark blue, weathering brown, sticky clay, is exposed (including the base); whilst a total thickness of 4.7m have been recorded from a shallow commercial borehole, directly to the west of the present quarry.

The 'Tilestone beds'

It is uncertain that the informal beds, previously recorded from several localities as the 'Tilestone' (Parsons, 1976, p.59), in fact represent a single, homogenous unit. In many cases they may include beds best considered part of the Lower-Trigonia-grit Member. Until further research clarifies this problem, it is thought inadvisable to formally redefine this unit.

The Hartley Hill Formation

Those beds preserved between the unconformities at the base of the Upper and Lower-Trigonia-grit Members (= 'Middle Inferior Oolite', Arkell, 1933), are here named the Hartley Hill Formation. At its type locality; Hartley Hill quarry (S0951180-6); this unit consists of approximately 8.0m of bioclastic limestones (Richardson, 1906; Murray, 1969). Elsewhere, notably Cleeve Hill, this Formation is thicker (up to 21.0m) and contains further beds of bioclastic and 'iron-shot' limestones (Richardson, 1929, pp.46-7). This Formation is here partly divided into three, formal Members - see Table 2.

The Lower-Trigonia-grit Member

The existing informal unit, the 'Lower <u>Trigonia</u> Grit' is here retained as a formal Member of the Hartley Hill Formation. The type locality for the re-defined unit remains the same, that is Leckhampton Hill (selected by Worssam, in Donovan and Hemingway, 1963, p.353), where it is represented by approximately 2.0m of ferruginous, 'iron-shot', bioclastic limestone (Ager in Torrens, 1969, p.B43; Murray, 1969, p.545).

The term Lower-Trigonia-grit should be restricted, where possible. to the iron-shot , ferruginous facies (= mainly obbiomicrites). However, the newly defined base of the Windrush Member (see below), results in the transfer to the present unit of beds previously included in the 'L.buckmani Grit! (e.g., Buckman, 1895, p.398, bed 8; p.399, bed 6). Whilst these beds are less ferruginous than the bulk of the Lower-Trigonia-grit Member. their highly fossiliferous nature, with an abundant, varied fauna, contrasts strongly with the succeeding, more sandy Windrush Member. Internal, informal, lithological subdivision of these beds is possible. The characteristic, 'iron-shot', highly bioturbated obbiomicrite, which makes up the bulk of this member, has previously been known as the 'Lamellibranch bed' (op. cit., p.399, bed 8). Similarly the top bed of this unit, previously included in the 'L. buckmani Grit', has been called the 'Serpula bed' (op. cit., p.409, bed 6; p.413, bed 6). The bulk of this Member has a well documented discites Zone age, whilst in the Painswick 'syncline', the top has locally yielded rare, basal ovalis subzone, laeviuscula Zone ammonites (see below).

The Windrush Member

The bulk of the beds previously known as the 'L. buckmani and Gryphite Grits', are here included in a single formal unit; the Windrush Member of the Hartley Hill Formation. Buckman's main criteria for separating the 'L. buckmani' from the 'Gryphite Grit', was the greater relative abundance of Lobothyris in the former (Buckman, 1895, p.390). This is an inadequate basis for the definition of a formal lithostratigraphic unit, especially since there are no lithological criteria for separating these beds at

either their type locality; Leckhampton Hill (Murray, 1969, p.545); or at many other localities such as Harford Cutting (Parsons, 1976) and in the Stowell Park bore-hole (Green and Melville, 1956, p.15). In these circumstances they should both be combined in a single formal unit, here called the Windrush Member, after the excellent exposures of those beds in the valley of the river Windrush. At its type locality, here designated as Harford Cutting (SP137218 = the "third from Bourton-on-the-Water"), the Windrush limestone consists of 3.7m of bioclastic limestones (= sparsely sandy biomicrites), alternating with sandy marls (Parsons, 1976, beds 4-9).

The base of this unit may be difficult to define, since there is often a gradational boundary between it and the Lower-Trigonia-grit Member. Where it is present, the Buckmani Bed acts as a useful basal 'marker bed' (Hedberg, 1976, pp.39-40), since it has a highly characteristic lithology, marking, as it does, a sharp influx of quartz sand. It also produces a strong anomaly in Beta and Gamma logs., which makes it easily traceable at the subsurface (Ponsford in Green & Melville, 1956, p.63, fig.6). Lithostratigraphic subdivision of the bulk of the unit is difficult, as Thus in the Leckhampton area, the base is relait is highly variable. tively massive, with rare Gryphaea, whilst the top is thinner bedded, with abundant Gryphaea. By contrast, in the Stroud district, the reverse is true, and it is the top which is more massive and less fossiliferous. The single most distinctive bed, or group of beds is the basal Buckmani Bed. The term 'buckmani - marl' was originally introduced by Buckman (1895, p.413, bed 5), for a more restricted horizon, now included in the basal part of the formal bed(s). At its type locality; here designated as the Frith quarry (SO866082); it is 0.30-0.35m thick (op. cit., p.399,

beds 5-6), and is made up at the base, of a grey-brown, soft marl, with crushed shell fragments, whilst the top consists of a fine, orange, quartz sand and silt, which penetrates the irregular base of the bed above. This sequence is typical of much of the Painswick 'syncline' and the southern part of the Cleeve Hill 'syncline', where the basal marl horizon often yields common specimens of <u>Lobothyris buckmani</u> (Davidson). However, at some localities a more complex sequence is found, often with the interdigitation of a sandy limestone between the sand and marl layers (e.g. Cats Brain quarry, Painswick; SO867114).

In the mid-Cotswolds these beds are wholly <u>ovalis</u> subzone, <u>laeviuscula</u> Zone in age. In the north Cotswolds, much of the Lower-Trigonia-grit Member seems to pass laterally into a sandy limestone facies, since here the base of the Windrush Member has yielded <u>discites</u> Zone ammonites (Parsons, 1976).

The Notgrove Member

The 'Notgrove Freestone' (Buckman, 1888) is here formally redefined as the Notgrove Member of the Hartley Hill Formation, with Notgrove Cutting (SP085209) as the type locality (selected by Worssam in Donovan & Hemingway, 1963, p.241), where it is represented by 3.6m of oolitic limestones, mainly comicrites (Richardson, 1929, p.76). Since the basal part of this section is now obscured, it is thought advisable to designate the Harford Cutting as a hypostratotype, where the member is 0.40-2.70m thick (Parsons, 1976, p.49). Although ammonites are very rare, these beds appear to be of <u>laeviuscula</u> Zone and subzone age.

The Cleeve Hill beds!

There are at present no good exposures of the beds found between the Notgrove and Upper-Trigonia-grit Members, since although the 'H. phillipsiana beds' are partly visible at the Rolling Bank quarry, Cleeve Hill (S0987267), there are currently no exposures of the 'Bourguetia beds' and 'Witchellia Grit' (= 'Perna bed', Wright, 1860, p.42). These beds all have a very restricted distribution, centred on a small area adjacent to the Cleeve Hill plateau (i.e. at the centre of the Cleeve Hill 'syncline'). Since their detailed inter-relationship cannot be currently established, no formal lithostratigraphic re-definition of these beds can be attempted. If the presence of a bored 'hard-ground' at the base of the 'Witchellia Grit' (Buckman, 1895, p.417, beds 5/6) could be confirmed, this would suggest the inclusion of this 'Grit', together with the 'H. phillipsiana and Bourguetia beds! in a single formal member, with the redefinition of individual horizons as formal bed(s). The age of these beds is well established, with abundant laeviuscula subzone ('Witchellia Grit') and sauzei Zone ammonite faunas.

The Stroud Formation

The unconformity below the Upper-Trigonia-grit Member, which marks the base of the Stroud Formation, is one of the most clearly defined in the whole Jurassic (Arkell, 1933, p.231). The Chipping Norton Formation (including the Hook Norton Limestone) has been placed within the Great Oolite Group (Sellwood and McKerrow, 1974), the Stroud Formation is thus restricted in the mid and north Cotswolds to the units previously known as the 'Upper Trigonia Grit' and 'Clypeus Grit'. At its type locality,

here designated as Rodborough Hill, Stroud, this Formation consists of +6.Om of predominantly oolitic limestone. The Fort Quarry (SO850041) probably shows the best existing exposures in the district, including a good section across the junction of this Formation, with the subjacent Hartley Hill Formation (Richardson, 1907a, p.76). The Stroud Formation is the most important within the Inferior Oolite Group of the Cotswolds, as it has the widest geographic distribution: it is the only one to be found across the whole Cotswold district, and the Mendips, Dorset and Somerset to the south.

The Upper-Trigonia-grit Member

The existing informal unit, the 'Upper Trigonia Grit', is here defined as a formal Member of the Stroud Formation. At its type locality, Leckhampton Hill (selected by Worssam in Donovan and Hemingway, 1963, p.353), this unit consists of about 1.5m (Ager in Torrens, 1969, p.B42) of bioclastic limestone (highly fossiliferous biomicrites), although at other localities up to 3.5m have been recorded (Richardson, 1907a). The 'Upper coral bed' (Witchell, 1882) is here informally defined as the upper-most bed of the Upper-Trigonia-grit Member, as exposures at its type locality. Rodborough Hill, are now too poor for formal redefinition. It is possible that at some Cotswold localities there may be some remnants of horizons older than the Upper-Trigonia-grit, although still deposited after or during the Upper Bajocian transgression. Just to the south of the Mendips the 'Doulting Conglomerate' Bed, of subfurcatum Zone age (Parsons, 1975a), is found in exactly this stratigraphic position and may thus warrant separation as a full Member of the Stroud Formation. The

Upper-Trigonia-grit Member is of acris subzone, garantiana Zone age.

The Clypeus-grit Member

This unit consists of those beds found below the Stroud, 'White Oolite', Chipping Norton Limestone or other Formations of the Great Oolite Group, and above the Upper-Trigonia-grit. This Member, which coincides in its limits with the informal unit of the same name, consists of 12.0m of rubbly oolitic limestone (= intraclastic, pisolitic biomicrites + oobiomicrites) at its type locality, here designated as the first cutting west of Notgrove (S0085209), (Richardson, 1929, pp.75-6). The Clypeusgrit has a disconformity at its base, which explains its over-stepping of the Upper-Trigonia-grit in the north and east Cotswolds (Richardson, 1929, p.73; Arkell, 1947, pp.28-30). This over-stepping results in the Clypeusgrit being the sole Member of the Stroud Formation over much of the Cotswolds. The status of Formation has been suggested for this unit (Kennedy, Sellwood and McKerrow, in Ager <u>et al.</u>, 1973, p.2), but this is rejected here as it would necessitate a similar, but unacceptable status for the Upper-Trigonia-grit Member.

Lithostratigraphic subdivision of this unit will undoubtedly prove possible. In particular the basal beds are more massive and intraclastic, whilst towards the top there are thin, harder, usually highly fossiliferous horizons, often with abundant nerineid gastropods. All these beds are of <u>parkinsoni</u> Zone, and probably <u>bomfordi</u> subzone age.

医骨髓炎 计算法分词 化合金

The upper boundary of the Stroud Formation

The top of the Stroud Formation is defined by the base of the Great

Oolite Group, which in the north and mid-Cotswolds falls at the base of the Chipping Norton Formation (Sellwood & McKerrow, 1974, pp.190-1). Thus in these areas the Stroud Formation is restricted to the Upper-Trigonia-grit and Clypeus-grit Members. It is because of their widespread and clear recognition, that these latter two units have been formally defined as Members of the Stroud Formation. In the south Cotswolds additional limestone units appear either above or as partial lateral equivalents of the Clypeus-grit (Stroud 'White Oolite', 'Anabacia Limestones' and 'Rubbly beds'; see Fig.5, & Richardson, 1907b), which have been included in the Inferior Oolite (Richardson, 1907b). One of the main criteria for determining the base of a Formation should be its mappability (Holland et al., 1978, p.8). In the south Cotswolds there is a relatively sharp lithological break, producing a well marked topographic feature, between the top of the Inferior Oolite Group, and the basal Formation of the Great Oolite Group; the Lower Fullers Earth. There is thus no doubt that the base of the latter Formation should define the top of the Stroud Formation. The fullonicus Limestone consists of a series of intermittent, argillaceous limestones, interbedded with clays and marls and resting disconformably on the 'Anabacia Limestones' (including the 'Rubbly beds': Torrens, 1969, p.B18). When deeply weathered it is difficult to differentiate between these former beds, and the bulk of the Fullers Earth, thus the main mappable feature is produced by the top of the Anabacia Limestones. Taking this into account, along with their close lithological and palaeontological affinity with the Fullers Earth, it is logical to include the fullonicus Limestone in the latter Formation (cf. Torrens, 1969). The 'Anabacia Limestones' and the 'White Oolite' are thus an integral part of the Stroud Formation.

No attempt is here made to formally define these units, since, far more work is needed on their lateral variation and distribution. A possible result of such work would be to establish the '<u>Anabacia Lime</u>stones' as a formal Member of the Stroud Formation, and to re-define the 'White Oolite' as a formal bed(s), within an extended Clypeus-grit Member. For the present, there is no difficulty in continuing to use these informal units, since a Formation need not be completely divided into Members (Holland <u>et al.</u>, 1978, p.10).

THE BAJOCIAN AMMONITE FAUNAS OF THE COTSWOLDS

Ammonites are not common in the Cotswold Inferior Oolite and our present knowledge of their distribution is the result of over one hundred and fifty years of intensive collecting. In the period 1880-1910 S. Buckman assembled a large collection of Cotswold ammonites, containing the cream of the material collected from the numerous small quarries open during this time. As well as ammonites which he had collected himself, Buckman's collection was swelled by material from his numerous friends, such as C. Upton and L. Richardson, as well as by specimens purchased from professional collectors and quarry men. Most of these ammonites are now preserved in the British Museum, Institute of Geological Sciences, Oxford University and Manchester Museum collections. The faunas which are discussed below are largely founded on this material, with various additions and amendations based on specimens which I and others have collected <u>in</u> <u>situ</u> (over one hundred specimens).

The discites Zone

The horizons which may be correlated with this Zone are very fossiliferous, and they have probably yielded almost as many ammonites as the rest of the Cotswold Bajocian rocks put together. The most prolific source of discites Zone ammonites has undoubtedly been the Lower-Trigonia-grit of the Frith quarry, near Stroud, where a single bed (Buckman, 1895, p.399, bed 8), has yielded large numbers of ammonites. particularly of the genus Hyperlioceras. Ammonites from this locality include: Hyperlioceras cf. liodiscites Buckman, BMNH.C9934; H. subsectum (Buckman), BMNH.C9944, C9955; H. walkeri (Buckman), BMNH.C9929; Darellia alta Buckman, L11236; D. cf. polita Buckman, BMNH.C9941; Reynesella cf. inops Buckman, OUM.J16166; Docidoceras cf. planulatum Buckman, OUM. J2465; Sonninia (Euhoploceras) sp., BMNH.C9933. The preservation of all these specimens (grey shells, yellow, ferruginous, micritic matrix, with common, very fine ferruginous coliths), is typical of bed 8, which has been confirmed by the location of an in situ specimen of H. cf. rudidiscites Buckman, CP1313, some 1.0m above the top of bed 18 (Buckman, 1895).

To the south of Frith quarry, the Lower-Trigonia-grit first appears beneath the "Upper Bajocian transgression" at Rodborough Hill, Stroud (<u>op. cit.</u>, p.394, bed 4). Here it has yielded <u>H. walker</u>i (Buckman, 1887-1907, p.94); <u>H. cf. liodiscites</u>, IGS.25319; <u>Darellia</u> sp., BMNH.C9928; <u>Reynesella</u> sp., BMNH.C9937 and <u>Docidoceras</u> cf. <u>planulatum</u>, BMNH.C9935. To the north the Lower-Trigonia-grit Member still yields <u>discites</u> Zone ammonites, such as <u>H. discoideum</u> (Qu. in SSB), CP3478, Frampton Mansell railway cutting (S0914028); <u>H. aff subsectum</u>, BMNH.C96095 and <u>Darellia</u> cf. <u>coela</u> Buckman, EMNH.C56098, from Cooper's Hill (S0887142; Channon,

1950, p.242); <u>Braunsina</u> cf. <u>projecta</u> Buckman, L11143 and <u>G</u>. (<u>Ludwigella</u>) aff. <u>stigmosum</u> Buckman, BMNH.C79539, Leckhampton Hill; <u>H</u>. <u>subsectum</u> (Buckman, 1887-1907, p.cxxi = ?IGS24751) and <u>Darellia</u> sp., IGS27627, from Wistley Hill (S0975185; Buckman, 1895, p.414) and <u>H</u>. aff. <u>subsectum</u> CP2529 and <u>G</u>. (<u>L</u>.) aff. <u>rudis</u> (Buckman), CP.3677, from bed 15b, Jackdaw quarry (Parsons, 1976). However over much of the north Cotswolds, because of the newly restricted limits of the Lower-Trigonia-grit, the most prolific <u>discites</u> Zone faunas come from the base of the Windrush Member. At Harford cutting this horizon (Parsons, 1976, bed 4), has produced: <u>Darellia</u> aff. <u>toxeres</u> Buckman, <u>H</u>. <u>subsectum</u>, <u>H</u>. <u>rudidiscites</u>, <u>Reynesella</u> sp. and <u>S</u>. (<u>Euhoploceras</u>) sp. (<u>op. cit</u>.).

All the ammonites cited above are typical of the <u>discites</u> Zone, as characterized by the upper part of the 'fossil-bed', Bradford Abbas, near Sherborne, Dorset (Parsons, 1974, p.170). The base of the <u>discites</u> Zone in the Cotswolds would appear to coincide with the unconformity below the Lower-Trigonia-grit, as the 'Tilestone bed' of the Stanway Hill Formation has produced a single ammonite indicating a possible <u>concavum</u> Zone age for this horizon (Parsons, 1976).

The laeviuscula Zone

The ovalis Subzone

The bulk of the Windrush Member of the Hartley HI I Formation has yielded few ammonites, but those which have been found in the mid-Cotswolds would indicate a correlation with the <u>ovalis</u> subzone of the <u>laeviuscula</u> Zone. Specimens of <u>Sonninia</u> (<u>Fissilobiceras</u>) <u>ovalis</u> (Buckman <u>ex</u>. Qu.), BMNH.C8787-8, from Leckhampton Hill (Buckman, 1895, p.512); <u>S</u> (F.) aff. ovalis, Harford cutting (Parsons, 1976, bed 8); <u>S. cf. patella</u> (Waagen), CP1312, from Harford cutting (op. cit. bed 6a); <u>S. (F.) aff. fissilobata</u> (Waagen), CP3338, Cats Brain quarry, Painswick; <u>Trilobiticeras (Emileites)</u> <u>liebi</u> (Maubeuge), EMNH.C9973, from Swifts Hill, Stroud (SO878067) and <u>Witchellia (Witchellia) romanoides</u> (Douvillé) from Swifts Hill, EMNH. C9969 and Stroud Hill, EMNH.C9974 and OUM.J37882, together make up a fauna similar to that recorded from the 'ovalis bed' of Dundry Hill, near Bristol (Parsons, 1974, p.169). The Windrush Member is also the inferred type horizon of <u>Witchellia</u> ('Zugophorites') <u>zugophorus</u> (Buckman, 1909-30, pl.341).

In the Stroud area, the uppermost part of the Lower-Trigonia-grit has also yielded <u>ovalis</u> subzone ammonites. These include <u>S</u>. (<u>F</u>.) aff. <u>fissilobata</u>, CP3343 from the Frith quarry (Buckman, 1895, p.399, bed 7) and <u>S</u>. (<u>?Euhoploceras</u>) aff. <u>acanthera</u> (Buckman), CP3477, from a similar horizon at the Frampton Mansell railway cutting (see Fig.3).

The laeviuscula subzone

Much of the Notgrove Member is virtually unfossiliferous. Only one ammonite has been previously recorded from this deposit, a specimen of <u>S</u>. (<u>F</u>.) aff. <u>ovalis</u> from Whittington (SPO13212; Buckman, 1897, p.608), whilst I have collected <u>S</u>. (?<u>Euhoploceras</u>) sp. CP3663, from the top 10 cm of this unit at the first cutting west of Notgrove. The exact age of the above specimens must remain in doubt, and they may well be of <u>ovalis</u> subzone age. In direct contrast, the thin and highly restricted horizon, the <u>Perna</u> bed or <u>Witchellia</u> Grit of the 'Cleeve Hill beds' has produced extensive ammonite faunas from two localities; Bolling Bank quarry, Cleeve Hill and Cold Comfort Farm. Ammonites from the latter include; <u>Emileia</u> (<u>Emileia</u>) aff. <u>brocchii</u>(Sow.), IGS.8077; <u>E</u>. (<u>Otoites</u>) cf. <u>fortis</u>, CP3664; <u>Witchellia</u> (<u>W</u>.) <u>patefactor</u> Buckman, holotype -IGS.47195 and OUM.J37866-9; <u>W</u>. (<u>W</u>.) <u>glauca</u> Buckman, OUM.J37874 and 37872, <u>W</u>. (<u>W</u>.) cf. <u>sutneri</u> (Branco), RUGC.S3140 and <u>Shirbuirnia</u>, cf. <u>platymorpha</u> (Buckman), OUM.J37876-7; whilst Rolling Bank has produced <u>W</u>. (<u>W</u>.) sp., OUM.J37881 and 37878, <u>W</u>. (<u>Pelekodites</u>) sp. OUM.J37880, <u>Shirbuirna</u> cf. <u>stephani</u> (Buckman), EMNH. C76772; <u>S</u>. aff. <u>superba</u> (Buckman), EMNH. C76771 and <u>S</u>. cf. <u>trigonata</u> (Qu.), OUM.J37879. These faunas are largely identical to that recorded from the 'green-grained marl' of Oborne, Dorset (Buckman, 1893, p.500, bed 9), which is to be correlated with the top of the <u>laeviuscula</u> Zone and subzone (Parsons, 1974, p.176).

The Notgrove Member is probably represented in some parts of the Cotswolds by a thin (-0.50m), highly bored limestone, variously called the 'Pitching' or 'Bored bed' (Buckman, 1893, p.512). This is the type horizon of <u>W</u>. (<u>W</u>.) <u>pavimentarius</u> (Buckman), holotype - IGS.49330, as has been confirmed by the location of <u>in situ</u> topotypes, CP2125, and 2127, from Leckhampton (loc. cit., bed 3).

The sauzei Zone

Only one ammonite is definitely known to have been collected <u>in situ</u> from the upper part of the 'Cleeve Hill beds'. This was the specimen of <u>Skirroceras aff. leptogyrale</u> Buckman, OUM.J10801 collected from bed III.1, Rolling Bank quarry (Buckman, 1897, p.609). However, other ammonites probably from this horizon, including <u>Emileia</u> (<u>Emileia</u>) cf. <u>bulligera</u> Buckman, CP.3665; <u>E. (E.) aff. pseudomultifida</u> Maubeuge, BMNH.C79434, and E. (E.) <u>vagabunda</u> Buckman, holotype - IGS.49313 (cf. Arkell, 1951-9, p.15), would indicate a correlation with the <u>sauzei</u> Zone 'Brown iron-shot' of Dundry Hill, Bristol (Parsons, 1979).

The garantiana Zone, acris subzone

Ammonites are not uncommon in the Upper-Trigonia-grit, and specimens of Strenoceras (Garantiana) spp. have been found at: Cleeve Hill, BMNH. C8822 and C80971; Harford cutting, BMNH.C79330-1, (Parsons, 1976, bed 11); Leckhampton Hill, L11193; Rodborough Hill, BMNH. C8839; Selsey Hill (S0826026), IGS.GSM25134, BUGM.3396; Stantonbury Hill (ST673636), Cb4982 (Tutcher, 1903, p.162) and Stroud Hill (Buckman, 1895, p.395). The latter specimen came from an horizon which Witchell (1882, p.60) correlated with the 'Upper coral bed'. There is no reliable evidence to suggest that the 'Upper coral bed' is any younger than the garantiana Zone. The specimens of Lissoceras psilodiscum (Schloenbach) from Wotton-under-Edge, RUGC.J523, (Richardson, 1910a, pp.103-4) and Stantonbury Hill (Tutcher, 1903, p.163), BUGM.89, which Buckman (in Tutcher, 1903, p.163) considered as indicating a parkinsoni Zone, truellei subzone age for this horizon are not diagnostic. This species ranges throughout the Upper Bajocian from the garantiana Zone (Parsons, 1975b, bed 13) up into the Bathonian (Krystyn, 1972, p.249). Other ammonites from the Upper-Trigonia-grit include Parkinsonia rarecostata (Buckman) from Long Wood, Stroud, BMNH.C9171; P. cf. subarietis Wetzel, CP3244, railway cutting west of Notgrove (SP085209); Leptosphinctes aff. meseres (Buckman), BUGM. 3396 and L. sp. BMNH. C80972 from Nibley Knoll (ST743956), Cadomites (Polyplectites) cf. gracilis (Westermann) from Upton Cheyney (ST694699), BUGM. 3389-3392, (= P. cf. linguiferus, Richardson,

1910a, p.90); <u>Oppelia (0.) aff. subcostata</u> (J. Buckman), from Brimpsfield, (S0931125), RUGC.R3074, (Richardson, 1903, p.387) and <u>Leptosphinctes</u> sp., BMNH.C9178 from Mount Surat, Rodborough. All these ammonites would support a correlation with the <u>acris</u> subzone of the <u>garantiana</u> Zone, as with the <u>Astarte</u> bed of south Dorset (Parsons, 1975a, fig.2). There is thus no evidence for the presence of any representatives of the bulk of the <u>garantiana</u> Zone (= <u>dichotoma</u>, <u>subgaranti</u> and <u>tetragona</u> subzones).

The parkinsoni Zone, bomfordi subzone

Members of the genus <u>Parkinsonia</u> are not uncommon in the Clypeusgrit. Specimens from this horizon include: <u>Parkinsonia</u> (<u>Parkinsonia</u>) cf. <u>crassa</u> Nicolesco (cf. Arkell, 1951-9, text fig.58), from, Harford cutting, BMNH.C8831 and CP2748; Roundhill Fox-covert (SP130219), CP2835, and Fawler (SP372173), OUM.J1187-8: <u>Parkinsonia</u> (P.) <u>eimensis</u> (Buckman, 1909-30, <u>non</u> Wetzel), from Roundhill Fox-covert, P. Hackling collection and Harford cutting, CP2940; <u>P. (P.) bomfordi</u> Arkell, from Fawler, OUM. J1180 and Upper Coberley (probably the Seven Springs quarry, Richardson, 1933, p.26), EMNH.C8828; <u>P. (?Okribites</u>) aff. <u>parkinsoni</u> (Buckman 1909-30, <u>non</u> Sow.), from Harford cutting, CP3134; the first cutting west of Notgrove, CP2837 and Little Tew (SP372 290), IGS.AH37. These ammonites suggest a correlation with the <u>bomfordi</u> subzone of the <u>parkinsoni</u> Zone. However, the distribution of the Parkinsonids is still very badly known, and further work is needed on the stratigraphy of the Dorset <u>parkinsoni</u> Zone, before any reliable, detailed correlations can be made.

11111 Figur 6 Formation scale, Pormation ¢ 2 schematic 1.4 1:0 and 13 Ó the "Tilestone orrelation 0.20 north Cotsvolds, 0 the Jo asped Lower which the Dajecian, Martley Stanway 1111 is approximately



N
CORRELATION OF THE COTSWOLD BAJOCIAN ROCKS

The correlation of the constituent Members of the Hartley Hill Formation is straightforward. Apart from minor changes in nomenclature, the correlation summarised in figure 3 is essentially the same as that put forward by Buckman (1895 and 1901) and Arkell (1933). There are however, more problems with the correlation of parts of the Stroud Formation in the south Cotswolds. A summary of the correlation suggested here is shown in figure 5. This differs in some important respects from that put forward by Richardson (1907b, pp.386-8 and 420-2, Tab.1 and fig.4; 1916, p.485), who suggested a more complex sequence, see figure 4. Richardson considered the 'Dundry and Doulting Freestones' as distinct deposits and hence recognized the following horizons in the Mendip/south Cotswold district:

(Rubbly beds

'Top beds'

(<u>Anabacia</u> Limestones (<u>Doulting Stone</u> (<u>Upper Coral bed</u> (<u>Dundry Freestone</u> (<u>Upper Trigonia Grit</u>

Unfortunately this succession is largely artificial, as it is the result of mis-correlation between three different areas (the Doulting district, the Cotswolds and Dundry Hill). No single locality has been described which has shown convincing exposures of all of the above deposits. Whilst there is an area of over-lap between the Cotswold and Mendip districts, which allows a degree of lithostratigraphic correlation, the isolation of Dundry Hill demands the use of biostratigraphic methods, as

Inferior Colite Text figure (1907b, 1910a S The 11 thostraticraphic * 1916) (stroud Formation) recognized in the hendips subdivisions and south Cotswolds. by dichardson 20 t'10 'Upper'

DOULTING	DUNDRY	COTSWOLDS
ANABACIA LIMESTONES	2 2	WHITE OOLITE
DOULTING STONE		CLYPEUS GRIT
	CORALLINE BEDS =	UPPER CORAL BED
	DUNDRY FREESTONE	7
DOULTING CONGLOMERATE	MAES KNOLL CONGLOMERATE	UPPER TRIGONIA GRIT

it is impossible to trace the lateral variation of these beds. Unhappily Richardson in the latter case relied heavily on the evidence of facies controlled faunas, such as that from the 'Upper coral bed' (Richardson, 1907b, p.387). This led him to make correlations which are now at variance with the ammonite evidence.

The Doulting district

In the Doulting railway cutting (ST51424), the Inferior Oolite is represented by the '<u>Anabacia</u> Limestones', 'Doulting beds' and by a basal conglomerate - the 'Doulting Conglomerate' (Richardson, 1907b, p.390). Richardson (<u>op. cit.</u>, p.386), considered that the 'Conglomerate bed' was to be correlated with the 'Upper <u>Trigonia</u> Grit' (<u>garantiana</u> hemera), the 'Doulting beds' with the '<u>Clypeus</u> Grit' (<u>truellei</u> hemera) and the '<u>Anabacia</u> Limestone' with the 'White Oolite' of Stroud (pre-<u>zigzag</u> hemera), although he had little or no ammonite evidence for these assumptions. These correlations made the presence of an unconformity necessary between the 'Doulting beds' and the 'Conglomerate bed', in order to explain the absence of the 'Dundry Freestone' and the 'Upper coral bed' at Doulting see Fig.4.

Recent work has changed this interpretation. The 'Doulting Conglomerate' is now known to be of <u>subfurcatum</u> Zone age, whilst ammonites indicating the presence of the <u>garantiana</u> Zone have been found in the 'Ragstone' (Parsons, 1975a), that is the basal part of the 'Doulting beds' (Richardson, 1907b, p.396). More is now also known of the ammonite distribution within the '<u>Anabacia</u> Limestones'. It is evident that the <u>Parkinsoni/zigzag</u> Zonal boundary falls within these beds, and that the lower half of the <u>Anabacia Limestones</u> is to be correlated with the <u>bomfordi</u> subzone of the <u>parkinsoni</u> Zone (Arkell, 1951-9, p.155; Torrens, 1969, p.B19). Since they fall between rocks of known <u>garantiana</u> and upper <u>parkinsoni</u> Zone age, it is logical to expect the bulk of the upper 'Doulting beds' to be <u>parkinsoni</u> Zone, <u>truellei</u> subzone in age. This is partly confirmed by the rare ammonites which are thought to be from this horizon: <u>Parkinsonia</u> cf. <u>dorsetensis</u> (Wright), Mells station cutting (ST710515, Richardson, 1907b, p.405); <u>Bigotites</u> cf. <u>lanquinei</u> (Nicolesco), BUGM. 3401, Great Elm, Vallis Vale; <u>Prorsisphinctes</u> sp., BUGM. 13459, Vallis Vale and Parkinsonia cf. <u>pseudoparkinsoni</u>, Wetzel BUGM.87, Holway.

The Dundry district

Richardson never published any detailed account of the Dundry Inferior Oolite, and it is evident from his references to Dundry, that he relied heavily on Buckman's work (Buckman and Wilson, 1896). The succession of beds within the Stroud Formation on Dundry is apparently simple, but it has been confused by poor exposures and a high degree of cambering. The beds in question, the 'Coralline beds', the Dundry Freestone and the basal, Maes Knoll Conglomerate, are poor in ammonites and hence difficult to date. Buckman found sufficient ammonites to indicate a <u>garantiana</u> hemera age for the Maes Knoll Conglomerate, but above this any correlation was pure conjecture (Buckman and Wilson, 1896, Tab.IV). Buckman suggested that the Dundry Freestone and 'Coralline beds' might be the equivalent of the 'Clypeus Grit' and the 'White Oolite' of Stroud respectively (op. cit., Tab.V).

Richardson made several amendments to these suggestions, since he

considered that the micro-fauna from the 'Coralline beds', particularly the micro-morphic brachiopods, indicated a correlation with the 'Upper coral bed! of the Cotswolds (Richardson, 1907b, p.387). If this were so, then in a complete succession, the Dundry Freestone would have to be stratigraphically lower than the 'Doulting Freestone', and separated from it by the 'Upper coral bed', or its equivalents, see figure 4. The facies controlled faunas from these coral beds were thus the only evidence for the duplication of the Freestone beds in Richardson's interpretation of the upper Inferior Oolite succession. Unfortunately only a few additional ammonites have subsequently been collected from these Dundry beds. Apart from the specimens of Garantiana recorded from the Maes Knoll Conglomerate (Buckman and Wilson, 1896, Tab.V), the specimens of Strenoceras (Pseudogarantiana) aff. dichotoma (Bentz), from Maes Knoll (ST598661), Cb4963, and of Parkinsonia rarecostata from the West End of Dundry Hill (ST553670), BUGM.3397, and Rackledown quarry (ST572654), Cb4962, all confirm that this horizon is to be correlated with the acris subzone of the garantiana Zone (Parsons, 1979). Ammonites are extremely uncommon above this horizon and include; Parkinsonia cf. parkinsoni (Wetzel non Sow.) from 1.0m above the best freestone beds on Dundry Down, BUGM. 13458; Parkinsonia parkinsoni (Buckman non Sow.), Cb4961, from Dundry village and Bigotites cf. tuberculatus (Nicolesco), Cb4959, from a freestone quarry at West Dundry. These ammonites suggest a correlation of the Dundry Freestone with the parkinsoni Zone and possibly with the truellei subzone. All that can be said of the 'Coralline beds' is that they are parkinsoni Zone, and probably just post-truellei subzone in age. The two ammonites known from the latter horizon; Leptosphinctes sp., South Main-road quarry (ST567655), Cb4993, and Cadomites (Polyplectites)

aff. gracile (Westermann), Rackledown quarry, Cb4994; are unfortunately too small and badly preserved to give a precise correlation.

The South Cotswolds

The upper Inferior Oolite exhibits typical Cotswold facies as far south as Chipping Sodbury. South of this point these horizons gradually assume their Mendip characteristics, with a decline in the amount of oolitic limestones present, associated with a marked relative increase in bioclastic material. Richardson (1910a), because of the rarity of ammonites, was forced to use lithostratigraphic methods, supported by the closely related facies controlled faunas of bivalves, brachiopods and echinoderms, to correlate these highly variable beds. He considered that the Upper Trigonia Grit persisted as far as the Mendips, whilst the 'Doulting beds' and 'Anabacia limestones' were the equivalent of the Clypeus Grit and Stroud, White Oolite' respectively (Richardson, 1910a, pp.85-7), - see figure 4. Richardson had little or no ammonite evidence to support these correlations, and unfortunately this situation has shown little subsequent improvement. Most of the ammonites which have been found between Bath and the Mendips are either of limited stratigraphic significance, or poorly localized. The most valuable ammonite fauna has come from the brown, crinoidal, ferruginous, slightly de-calcified limestone, which is the undoubted lateral equivalent of the Upper-Trigoniagrit. Specimens from this horizon include: Strenoceras (Garantiana) sp. Dunkerton (ST700589), BUGM.86. (Tutcher and Trueman, 1925, p.624); S. (G.) cf. garantiana, Midford Railway cutting, (Richardson, 1910b, p.98); S. (G.) garantiana, Cb4964, S. (G.) cf. garantiana (Pavia non d'Orb.), Cb4965, S. (G.) sp. Cb4967 and S. (Pseudogarantiana) cf. minima (Wetzel),

Cb4966, Limpley Stoke and Murhill, Bath (ST782612-795607); <u>Parkinsonia</u> <u>rectangularis</u> (Wetzel), Viaduct quarry, Limpley Stoke, Cb4960 and <u>S</u>. (<u>G</u>.) sp., Stantonbury Hill, Cb4982: all of which indicate a correlation with the <u>acris</u> subzone of the <u>garantiana</u> Zone.

Correlation of the Doulting, Dundry and south Cotswold districts

Using the ammonite evidence it is possible to produce a simple interpretation of the Stroud Formation in the south Cotswolds - see figure 5. First the 'Doulting Conglomerate', with its subfurcatum Zone fauna is unique, as it has no known Cotswold or Dundry equivalents. Secondly the garantiana Zone beds are the most easily correlated, since ammonites of this age have been found in the Cotswold Upper-Trigonia-grit. the Dundry Maes Knoll Conglomerate and in the Doulting 'Ragstone'. The Richardson's latter record thus totally discredits interpretation Text figure 4) of the correlation of these (800 beds , since the Dundry Fst. is unquestionably post-garantiana Zone in Such evidence as is available would suggest that much of the age. 'Doulting Freestone' sensu stricto, is truellei subzone in age. Since the 'Doulting' and Dundry Freestones are thus possibly of the same subzonal age, occur in identical stratigraphic positions resting disconformably on garantiana Zone rocks, and since no locality has been convincingly recorded at which they are both preserved, it would seem unnecessary to maintain them as distinct stratigraphic units. As the Dundry Stone or Freestone has priority, it has been re-defined as a formal Member of the Stroud Formation (Parsons, 1979) with Dundry Down quarry (ST552666), Dundry as its type locality, where it consists of 4.6m of massive bioclas-



Figure 5.

A schematic correlation of the Upper Bajocian, Stroud Formation in the Mendips and south Cotswolds. tic limestone.

The Clypeus-grit Member of the Cotswolds can be correlated with a basal part of the "Anabacia Limestones", since they both contain bomfordi subzone faunas. Taking their relative position into account, it is likely that the 'Coralline beds' are also to be correlated with this subzone, although this needs to be confirmed by the location of further in situ ammonites. This latter correlation is partly substantiated by the rare occurrence in the 'Coralline beds' of the characteristic echinoid Clypeus ploti Klein (T. Fry, pers. com., 1974). There is thus probably a disconformity over much of the Cotswolds between the Clypeus-grit and Upper-Trigonia-grit Members, which marks the position of the absent Doulting/Dundry Freestones (Parsons, 1979). The so-called 'Upper coral bed' is of limited stratigraphic significance. It is probably of garantiana Zone age in the Stroud district, but elsewhere, such as Timsbury Sleight (Richardson, 1907b, p.413), it could be of any age. There is no stratigraphic value to the debris produced by the erosion of corals and their associated facies fauna. These correlations are summarized in figure 5. It must however be stressed that this is still only a provisional interpretation, which must be followed up by further in situ ammonite collection.

ACKNOWLEDGEMENTS AND LOCATION OF MATERIAL

Whilst some of the material recorded here is in the author's collection (CP.), other specimens are located in the British Museum, (Nat. Hist.), London (BMNH.); Bristol City Museum (Cb.); Bristol

University, department of Geology (BUGM.); Institute of Geological Sciences, London (IGS.); Manchester City Museum (L.); Oxford University Museum (OUM.) and Reading University, department of Geology (RUGC). I should like to thank the curators of these museums for access to, and the loan of material in their care. As well as the various individual collectors who have helped in this work, notably P. Hackling of Hook Norton and T. Fry of Bristol, I should also like to thank Dr. I.E. Penn and Dr. H.S. Torrens, for their comments on an earlier version of this manuscript.

这些推动的生活的,并且是有的生活的和心理的。这种我们

Section and the sector of the section of the

(特定)用其你没有把这个,不是知道,就会一点

REFERENCES

AGER, D.V. 1956. Field meeting in the Central Cotswolds. Proc. Geol. Ass. 66, 356-65.

, DONOVAN, D.T., KENNEDY, W.J., McKERROW, W.S., MUDGE, D.C. and SELLWOOD, B.W. 1973. The Cotswold Hills. Geologist's Association guide No.36. 34pp.

ARKELL, W.J. 1933. The Jurassic System in Great Britain. Oxford University Press, xii + 681 pp., 41 pls.

, 1947. The Geology of Oxford. Oxford University Press, 267pp.

, 1951-9. A Monograph of the English Bathonian Ammonites. Palaeontograph. Soc. vii + 264pp., 33 pls.

_, 1956. Jurassic Geology of the World. Oliver and Boyd, Edinburgh, xiv + 806pp., 46 pls.

BUCKMAN, S.S. 1887-1907. A Monograph of the Ammonites of the Inferior Oolite Series. Palaeontograph. Soc. cclxii + 455pp., 127 pls.

> , 1893. The Bajocian of the Sherborne district: its relation to subjacent and superjacent strata. <u>Q.Jl geol. Soc. Lond.</u> 49, 479-522.

. 1895. The Bajocian of the Mid-Cotteswolds. Q.Jl geol. Soc. Lond. 51, 388-462.

> , 1897. Deposits of the Bajocian Age in the Northern Cotteswolds; the Cleeve Hill Plateau. Q.Jl geol. Soc. Lond. 53, 607-29.

, 1901. Bajocian and Contiguous Deposits in the North Cotteswolds; the Main Hill-Mass. Q.Jl geol. Soc. Lond. 57, 126,55.

, 1909-30. Type Ammonites. London and Thame, 1-7, 790 pls.

, and WILSON, E. 1896. Dundry Hill: its Upper Portion or the Beds Marked as Inferior Oolite in the Maps of the Geological Survey. Q.Jl geol. Soc. Lond. 52, 669-720.

- CAVE, R. and PENN, I.E. 1972. On the Classification of the Inferior Oolite of the Cotswolds. <u>Bull. Geol. Surv. G.B.</u> 38, 59-65.
- CHANNON, P.J. 1950. New and Enlarged Jurassic Sections in the Cotswolds. Proc. Geol. Ass. 61, 242-60.
- DONOVAN, D.T. and HEMINGWAY, J.E. 1963. Lexique Stratigraphique International, Vol.I, fasc. 3aX. C.N.R.S. Paris, 394pp.
- GREEN, G.W. and MELVILLE, R.V. 1956. The Stratigraphy of the Stowell Park Borehole. Bull. Geol. Surv. G.B. 11, 1-66.
- HEDBERG, H.D. 1976. International Stratigraphic Guide : <u>A guide to</u> stratigraphic classification, terminology and procedure. Wiley, New York, xvii + 200pp.
- HOLLAND, C.H. et al. 1978. A guide to stratigraphical procedure. Geol. Soc. Lond. Spec. Rep. No.10, 18pp.
- KELLAWAY, G.A. and WILSON, V. 1941. An outline of the geology of Yeovil, Sherborne and Sparkford Vale. <u>Proc. Geol. Ass</u>. 52, 131-174, pls. 8-10.
- KENT, P.E. 1975. The Grantham Formation in the East Midlands : Revision of the Middle Jurassic, Lower Estuarine beds. <u>Mercian geol</u>. 5, 305-327.
- KRYSTYN, L. 1972. Die Oberbajocium- und Bathonium-Ammoniten der Klaus-Schichten des Steinbruches Neumühle bei Wien (Österreich). <u>Ann. Naturhistor. Mus. Wien</u>. 76, 195-310, 24 pls.
- LYCETT, J. 1850. Tabular view of Fossil Shells from the middle division of the Inferior Oolite in Gloucestershire. <u>Proc. Cotteswold</u> <u>Nat. Field Club</u> 1, 62-86, 1 pl.
- McKERROW, W.S., AGER, D.V. and DONOVAN, D.T. 1964. Geology of the Cotswold Hills. Geol. Ass. Guide No.36, 26pp.

and KENNEDY, W.J. 1973. The Oxford District. Geol. Ass. Guide No.3, 46pp.

MUDGE, D.C. 1978. Stratigraphy and sedimentation of the Lower Inferior Oolite of the Cotswolds. <u>Jl. geol. Soc. Lond</u>. 135 : 611-627.

> , 1978a. Supplementary Publication No. SUP 18026, British Library, Boston Spa, 32pp.

MURCHISON, R.I. 1834. Outline of the Geology of the Neighbourhood of Cheltenham. Davies, Cheltenham. 40pp.

- MURRAY, J.W. 1969. The Inferior Oolite of the Cotswold Scarp, Wootonunder-Edge to Leckhampton. <u>Proc. Bristol Nat. Soc</u>. 31, 535-49.
- PARSONS, C.F. 1974. The <u>sauzei</u> and 'so called' <u>sowerbyi</u> Zones of the Lower Bajocian. <u>Newsl. Stratigr. 3</u>, 153-80.

, 1975a. Ammonites from the Doulting Conglomerate Bed (Upper Bajocian, Jurassic) of Somerset. <u>Palaeontology</u> 18, 191-205, 1 pl.

, 1975b. The Stratigraphy of the Stony Head Cutting, Bridport, Dorset. Proc. Dorset Nat. Hist. Archaeol. Soc. 96, 8-13.

, 1976. Ammonite evidence for dating some Inferior Oolite sections in the north Cotswolds. Proc. Geol. Ass. 87, 45-64.

, 1979. A stratigraphic revision of the Inferior Oolite of Dundry Hill, Bristol. Proc. Geol. Ass. 90, 133-151.

RICHARDSON, L. 1903. On a section at Cowley, near Cheltenham, and its bearing upon the Interpretation of the Bajocian Denudation. <u>Q. Jl geol. Soc. Lond.</u> 59, 382-9.

> , 1906. Excursion to Leckhampton Hill. Proc. Cotteswold Nat. Field Club, 15, 182-9.

> , 1907a. On the Top-Beds of the Inferior Oolite at Rodborough Hill, near Stroud. <u>Proc. Cotteswold Nat. Field Club</u>, 16, 71-80.

, 1907b. The Inferior Oolite and Contiguous Deposits of the Bath Doulting District. Q.Jl geol. Soc. Lond. 63, 383-423.

, 1910a. The Inferior Oolite and Contiguous Deposits of the south Cotteswolds. Proc. Cotteswold Nat. Field Club, 17, 63-136, 7 pls.

, 1910b. On the sections of Inferior Oolite on the Midford-Camerton section of the Limpley Stoke railway, Somerset. Proc. Geol. Ass. 21, 97-100, 2 pls.

- RICHARDSON, L. 1916. The Inferior Oolite and Contiguous Deposits of the Doulting-Milborne-Port District (Somerset). Q. Jl geol. Soc. Lond. 71, 473-520, 4 pls.
 - , 1928-30. The Inferior Oolite and Contiguous Deposits of the Burton Bradstock - Broadwindsor District. Proc. Cotteswold Nat. Field Club, 23, 35-68, 149-85 and 253-64, 13 pls.

, 1929. The country around Moreton-in-the-Marsh. Mem. Geol. Surv. G.B. 126pp.

, 1932. The Inferior Oolite and Contiguous Deposits of the Sherborne District, Dorset. Proc. Cotteswold Nat. Field Club, 24, 35-85, 6 pls.

, 1933. The Country around Cirencester. <u>Mem. Geol. Surv.</u> <u>G.B.</u> 119pp.

- SELLWOOD, B.W. and McKERROW, W.S. 1974. Depositional Environments in the Lower Part of the Great Oolite Group of Oxfordshire and North Gloucestershire. <u>Proc. Geol. Ass</u>. 85, 189-210.
- SPATH, L.F. 1936. On Bajocian Ammonites and Belmnites from eastern Persia (Iran). <u>Palaeont. Indica</u> (N.S.), 22, 1-21, 1 pl.
- SYLVESTER-BRADLEY, P.C. and FORD, T.D. 1968. The geology of the East Midlands. Leicester Univ. Press. 400pp. 7 pls.
- TORRENS, H.S. (Editor) 1969. International Field Symposium on the British Jurassic. Excursion Guide No.2, North Somerset and Gloucestershire. University of Keele, 47pp.
- TUTCHER, J.W. 1903. The Lower Oolites near Bristol. Proc. Bristol Nat. Soc. (N.S.), 10, 150-68.

and TRUEMAN, A.E. 1925. The Liassic Rocks of the Radstock District, Somerset. Q.Jl geol. Soc. Lond. 81, 595-666, 4 pls.

- WILSON, V., WELCH, F.B.A., ROBBIE, J.A. and GREEN, G.W. 1959. Geology of the country around Bridport and Yeovil. <u>Mem. Geol. Surv. G.B.</u>, xii + 239pp. 7 pls.
- WITCHELL, E. 1880. On a section of Stroud Hill, and the Upper Ragstone Beds of the Cotteswolds. <u>Proc. Cotteswold Nat. Field Club</u>, 7, 118-35.

- WITCHELL, E. 1882. The Geology of Stroud and the area drained by the Frome. James, Stroud, viii + 108pp. 3 pls.
- WOODWARD, H.B. 1894. The Jurassic Rocks of Britain iv Lower Oolitic Rocks of England. <u>Mem. Geol. Surv. G.B</u>. 628pp.
- WORSSAM, B.C. and IVIMEY-COOK, H.C. 1971. The Stratigraphy of the Geological Survey borehole at Warlingham, Surrey. <u>Bull. Geol.</u> <u>Surv. G.B.</u> 36, 1-111.
- WRIGHT, T. 1856. The Palaeontological and Stratigraphical Relations of the so-called 'Sands of the Inferior Oolite'. Q.Jl geol. Soc. Lond. 12, 292-325.
- , 1860. On the Subdivisions of the Inferior Oolite in the South of England, compared with the Equivalent Beds of that Formation in Yorkshire. Q. Jl geol. Soc. Lond. 16, 1-48.

Render and the second

2C.A STRATIGRAPHIC REVISION OF THE SCARBORCUGH FORMATION, MIDDLE

JURASSIC, OF NORTH-EAST YORKSHIRE

CONTENTS

I.	INTRODUCTION	257
II.	HISTORY OF PREVIOUS RESEARCH	258
III.	THE BIOSTRATIGRAPHY OF THE SCARBOROUGH FORMATION	263
	(a) The subzone of <u>Dorsetensia romani</u> (Oppel)	
	(b) The subzone of <u>Stephanoceras humphriesianum</u> (Sow.))
	(c) The subzone of <u>Teloceras</u> <u>blagdeni</u> (Sow.)	
	(d) Stratigraphically unlocalised ammonites	
IV.	THE LITHOSTRATIGRAPHY OF THE SCARBOROUGH FORMATION	271
	(a) The Hundale Shale Nember	
	(b) The Crinoid Grit Member	
	(c) The Spindle-Thorn Limestone Member	
	(d) The Ravenscar Shale Member	
	(e) The Lambfold-Hill Grit Nember	
	(f) The White Nab 'Iron-stone' Member	
	(c) The 'Transition shales'	
۷.	THE DESCRIPTION OF THE COASTAL EXPOSURES	276
	(a) Ravenscar	
	(b) Hundale Point	
an di Africa. Ang ika sa	(c) White Nab and Black Rock, Scarborough	
	(d) Gristhorpe Bay	
VI.	STRATIGRAFHIC SUMMARY	292
VII.	ACKNOWLEDGENERTS 295	
VIII.	REFTRENCES	296

SUMMARY

A resume of previous work on the stratigraphy of the Scarborough Formation (Bajocian Stage, Middle Jurassic) and an analysis of its ammonite faunas are given. It is shown that the upper part of the Scarborough Formation spans the Romani, Humphriesianum and Blagdeni subzones of the Humphriesianum Zone, whilst the lower part is to be correlated with a lower, but as yet unspecifiable Zone. The Lithostratigraphy of this Formation is described, several new, formal lithostratigraphic units are introduced (the Hundale shale, the Ravenscar Shale and the White Nab 'Iron-stone' Members) and the main coastal exposures of this Formation at Gristhorpe, Scarborough, Cloughton and Ravenscar are described.

I. INTRODUCTION

The Jurassic rocks of North-East Yorkshire were some of the first in the World to be the subject of detailed scientific study. In the early part of the Nineteenth Century notable pioneer geologists such as N. Smith, A. Sedgwick, J. Phillips, W. Williamson, all left their mark on the progress of Jurassic Stratigraphy in Yorkshire. Towards the end of that century however, the focus of Jurassic workers had shifted towards the South of England, and to the present day this area of the country has been the more intensely studied, often at the expense of other districts. This picture of relative neglect of the Yorkshire Jurassic, is true of the particular horizon which is the subject of this paper; the Scarborough Formation, (Middle Jurassic, Bajocian Stage) - see Table 1. Our knowledge of the stratigraphy of this unit, as based on well localised ammonite collections, has not improved since Williamson's work of 1840. Whilst it is true to say that ammonites are not common in this marine intercalation within the largely nonmarine 'Deltaic Series' (Ravenscar Group, Hemingway & Knox, 1973), a careful search of the more extensive exposures of the Scarborough beds, will not usually go unrewarded. Since, as I have said, large areas of exposure are needed, in order to 'shorten the odds' of finding an in situ amaonite, only the coastal exposures are of any real value for stratigraphic work. Of these, the exposures at Ravenscar, Hundale Point (Cloughton), White Nab (Scarborough) and Gristhorpe Bay are the most extensive and accessible (see Text

fig. 1) and are described here.

The following account summarises the previous stratigraphic

Scalby Formation

Scarborough Formation

Cloughton Formation

Gristhorpe Member Lebberston Member Sycarham Member

Eller Beck Formation

Saltwick Formation

Ravenscar

Group

Table 1.

The lithostratigraphic subdivisions of the Ravenscar Group in north-east Yorkshire, after Hemingway and Knox (1973).

work on these beds, as well as giving details of the relevant ammonites in existing Museum collections, my own collections and those of M. Bradshaw, G. Farrow, Professor P.C. Sylvester-Bradley and J.K. Wright. These new ammonite records have enabled a more detailed correlation to be made of the Scarborough Formation and have made the task of lithostratigraphic subdivision that much easier.

II. HISTORY OF PREVIOUS RESEARCH

The marine horizons within the Ravenscar Group are relatively thin, and escaped the notice of many of the earlier workers, who grouped all of the non-marine sandstones together. Thus Conybeare and Phillips (1822) correlated all of these beds with the Inferior Oolite - Fuller's Earth series and Sedgwick (1826), who recorded specimens of 'Perna', Triconia and 'Avicula' from what were later to be known as the Scarborough Beds of White Nab, (Sedgwick, 1826, p.357), included them all within his 'Newer Coal Formation', (op. cit. p.353). It was John Phillips who gave the first detailed description of the Scarborough Formation, which he variously called the Bath Colite or Grey Limestones; (Phillips, 1829, p.140), and who recorded the first ammonite from this horizon; 'Ammonites' (Teloceras) blagdeni Sowerby, from White Nab, (op. cit. p.152). In spite of recording this typical Bajocian ammonite species, Phillips was able to state - "the Grey Limestone of the Yorkshire Coast is equaeval with the Great or Middle Oolite of Eath" (loc. cit.). In the second edition of his 'Illustrations of the Geology of Yorkshire', Phillips (1835), gave a few more details of the Scarborough Beds, but recorded no further annonites.





A sketch map of the main outcrops of the Scarborough Formation in the north-east Yorkshire Noors, showing the location of the major localities cited in the text.

Williamson (1840) gave a detailed section of what he called the Great or Eath Oolite, of White Nab, Scarborough, and also gave the first well localised records of ammonites from this horizon: citing 'Ammonites' blagdeni from his bed d. and more rarely from bed f, (Williamson, 1840, pp.231-3). This last account is one of the most important works on the Scarborough beds, and Williamson's section is repeated, in a slightly modified form, by several subsequent authors. Morris and Lycett (1851-5) in their famous monograph continued to include the Scarborough beds within their concept of the Great Colite, in spite of their comment that they "possess few mineral features which serve to connect them with their supposed equivalent in Gloucestershire" (Morris & Lycett, 1851-5. p.VI). Apart from specimens of Belemnites giganteus and A. blagdeni, Morris and Lycett figure one ammonite of real interest from the Scarborough beds, A. braikenridgi, which was forwarded to them under the manuscript name A. triptolemus, by Bean. This latter specific name was rejected by them, but was subsequently resurrected by Buckman, (1912).

Oppel (1856) was able to anticipate Wright's correlation of the Scarborough beds, with the Inferior Colite of Southern England, when he stated (<u>op. cit. p.340</u>) - "I am convinced that Phillip's Cave Colithe (or Bath or Great Colite) is entirely identical with the <u>A</u>. <u>humphriesianus</u> schicten of other areas" and recorded the following from Scarborough to support this conclusion; <u>'Beleanites' ciganteus</u>, <u>A. blagdeni A. humphriesianus</u>, and <u>A. subcoronatus</u> Oppel. Phillips however still stuck to his correlation with the Bath Colite' (Phillips, 1858) and it was not until the publication of Wright's paper (1860), that he was willing to admit his error "Dr. Wright has

lately placed before the Society a proposal to assign to the Inferior Colite a portion of the sandy, shaly, irony and calcareous beds which Professor Morris and myself refer to the Great Oolite of Bath. In the section referred to, between unequivical Lias and unequivical Combrash, there have been gathered very few ammonites In addition to 'Ammonites' blagdeni, which was found at Gristhorpe so long ago, the diligent naturalists of Scarborough have collected A. murchisonae ..., A. humphreysianus and A. parkinsoni which makes the palaeontological evidence in favour of the slightly greater antiquity of these Oolitic beds preponderate" (Phillips, 1860, p.xli). At this time Phillips also recorded a specimen of A. humphriesianus, which he had found at Gristhorpe in 1855, (op. cit. p. xlii). Wright, although finally getting the age of the Scarborough beds correct in fact added very little to our knowledge of these beds, since he merely repeated Williamson's section; with A. humphriesianus and A. braikenridgi in addition to A. blagdeni from White Nab, (Wright, 1860, p.29).

We have now reached a lull in the progress of research on the Scarborough Formation. Hudleston, who introduced the term 'Scarborough limestones', rather than Grey limestones', (Hudleston, 1874, p.292) and Fox-Strangways (1892, 1904 & 1915) described many detailed sections, but recorded no new ammonites, except for "<u>Ammonites laeviusculus</u>"

(= ? Dorsetensia sp.), from Hundale, (Hudleston, 1882, p.147).
The honour of recording the first members of the genus
Dorsetensia from the Scarborough beds must go to Drake (1903), who
Pre-empted Buckman's identification of material from the Herries

collection by four years. In a short, and totally forgotten note in the 'Naturalist', Drake recorded two specimens of <u>Dorsetensia</u> <u>subtecta</u> Buckman from the Scarborough beds of Hundale Point, one from his own collection and the other from that of a friend, G. Sheppard, (Drake, 1908, pp.296-7). In the same paper Drake also mentions two specimens of <u>Normannites braikenridgi macer</u> (Qu.), which he had collected from the fore-shore at Scarborough. It is unfortunate that neither Drake nor Buckman (1912) made any attempt to give an accurate horizon for these specimens of <u>Dorsetensia</u>

, and it has taken to the present day, to locate the faunas in situ.

The most important stratigraphic work on the Scarborough beds is undoubtedly that of Richardson (1912), who, in the company of E.T. Paris, had studied in detail the outcrop of these beds. Richardson introduced several new, named lithostratigraphic units, proposed a new name for the whole series; the Scarborough beds rather than Scarborough limestone, since there is in fact very little limestone present; as well as recording several species of ammonites. Within the main body of his paper Richardson (1912, p.195) recorded Teloceras coronatura (Qu., non Brugière), Stepheoceras subcoronatum (Oppel) and Parkinsonia sp. from White Nab; whilst in an Appendix, Buckman discussed the following species Skirroceras ? triptolemus (Bean m.s.), Teloceras sp. nov. (= T. blagdeniformis (Roche)), T. coronatum (Qu.), Stemmatoceras subcoronatum (Oppel), Stepheoceras of. sieteni (Qu.), S. cf. pyritosum (Qu.), Dorsetensia romani and D. liestraca S.B., which were either re-interpretations of material previously recorded from the Scarborough beds, or new records from the Herries collection, (Buckman, 1912, pp.205-8).

Buckman used the Zonal divisions of the Bajocian established by Mascke in North Germany, and from the previously cited specimens, concluded that the Scarborough beds spanned Mascke's <u>Stemmatoceras</u>, <u>Stephanoceras</u>, <u>Stepheoceras</u> and possibly the basal <u>Teloceras</u> Zones, (<u>op. cit. p.208</u>). These Zones all fall within Buckman's Blagdeni hemera, (Buckman, 1910).

Arkell's well known synopsis of the British Jurassic, (Arkell. 1933), is positively misleading as far as the Scarborough beds are concerned. Arkell quoted Buckman's above mentioned ammonite identifications, and correctly inferred a Blagdeni zone date for this fauna, (op. cit., p.221). However, Arkell made the serious mistake of following Richardson's generalised division of the Scarborough beds into three main sections; an upper shale division, a middle sandstone division and a lower limestone division, and solely on this basis, made the assumption that all the ammonites found had come from the lower limestone division. It was not until 1965, that Bate, pointed out that the opposite is in fact true, and that most ammonites come from the upper portion of the sequence, (Bate, 1965, p.93). In the forty years subsequent to the publication of Richardson's paper, there has been very little progress in elucidating the stratigraphy of these beds. In 1951, Sylvester-Bradley recorded a single specimen of Dorsetensia pulchra S.B. from Monk's Walk Wood, Sneaton, (Hemmingway, 1951, p.119), and since that time there have been various other records of Dorsetensia from the coast. Rare specimens of Dorsetensia have been recorded from Ravenscar, (Henningway, et al., 1969, p.C18), and from Hundale point, (Hemingway et al., 1963, p.19).

In 1953, Sylvester-Bradley gave a useful summary of the best

exposures of the Scarborough beds, and cited <u>Teloceras blagdeniformis</u> (Roche) as a typical ammonite, following Cox and Arkell's (1948-50) re-identification of the <u>Teloceras blagdeni</u> Sow., figured by Morris and Lycett, (1851-5, Pl. xiv. fig 3). Other recent publications, covering the Scarborough Formation include, Farrow's work on the trace-fossils, where several unspecified ammonite horizons are recorded in a text figure, (Farrow, 1966, text fig. 3); Bate's work on the Ostracod faunas, where, as well as giving numerous highly detailed, measured sections, he divided the Scarborough Formation between two Ostracod Zones; an upper Zone of <u>Glyptocythere scitula</u> Eate and a lower of <u>G. polita</u> Bate, (Bate, 1965), and Hemingway's general and sedimentological description of these beds, where he unfortunately assigned them to the Bathonian Stage (Hemingway in Rayner & Hemingway, 1974, p.185).

This then has been the progress of stratigraphic research on the Scarborough Formation, and it is sad to note that there have been no records of ammonites accurately collected <u>in situ</u>, from measured sections, since Williamson's work of 1840.

III. BIO-STRATIGRAPHY OF THE SCARBOROUGH FORMATION

The ammonite faunas to be described below and summarised in text fig. 3, may all be included within the Humphriesianum Zone of the Lower Bajocian. A three-fold subzonal division of this Zone is adopted here -

Humphricsianum Zone

Blagdeni subzone Humphriesianum subzone Romani subzone

This follows a similar subdivision of the Humphriesianum Zona

described from the Basse Alpes of South-East France, (Pavia & Sturani, 1968), but uses pre-existing subzonal indices, rather than any new terminology; thus the Romani subzone (Haug, 1891) = the Cycloides subzone, (Sturani, 1971).

The stratigraphy of the North German Bajocian, of which the Yorkshire rocks are merely a continuation, mirrors these subdivisions.

Subzones used	Zones of Masche	Zones of Kumm.
here	1907	1952
Blagdeni	Teloceras -	Blagdeni
Humphriesianum	Stepheoceras =	Humphriesi
Domoni	(<u>Stephanoceras</u> =	Umbilicum
n Omen L	(<u>Stemmatoceras</u> =	Coronatum

NOTE:

In the following section specimen numbers prefixed by CP. refer to specimens in the author's collection, and by G. to specimens in the Scarborough Natural History Museum. Secondly any bed numbers quoted from the Gristhorpe, White Nab and Hundale Point exposures refer to the sections which are re-described here, whilst bed numbers quoted from any other sections are mainly those used by Bate (1965).

(a) The subzone of Dorsetensia romani (Oppel)

The lowest ammonite collected in situ during the course of this Work was a specimen of <u>Stephanoceras</u> (<u>Skirroceras</u>) sp. from the lower Part of the Spindle-Thorn Limestone (bed 14), Hundale Point (National Grid Reference, TAO25948), some 0.43m. above the top of the Crinoid Grit. This specimen (CP.3094), which is too squashed and fragmentary for positive identification, probably belongs to the <u>S</u>. (<u>S</u>.) <u>nodosum</u>

(Hyatt) - macrum (Qu. emend. Renz) group, which is typical of the middle Humphriesianum Zone in south Germany, but which also occurs as low as the Romani subzone. From further up in the Spindle-thorn limestone a single specimen of Dorsetensia sp. has been observed in bed 18d, but it was too squashed and distorted to facilitate extraction. Directly above this horizon, in the base of the Ravenscar Shale, specimens of Dorsetensia become relatively abundant. In the small cliff in the Ravenscar shale, just to the south of the actual point at Hundale, specimens of Dorsetensia have been found at the following heights above the base of bed 20 : D. cf. deltafalcata (Qu.), + 0.70m. (CP3067); D. cf. liostraca S. Buckman. + 1.7 & + 2.9m. (CP2645); D. cf. romani, + 0.70, + 2.80 & + 2.90m. (CP.2223, 2723, 2728 & 9), and D. spp. + 1.1 + 2.80m. (CP.2730 & 3064). Similarly on the fore-shore, three, large specimens of D. liostraca (CP.2681, 2964 & 2987) were found in the shale reefs at a similar horizon to those specimens found in the cliff.

This locality at Hundale Point has in the past been the source of numerous specimens of <u>Dorsetensic</u>. Apart from those ammonites mentioned by Drake (1908), which were destroyed, along with the Hull Museum in the last war, the Herries collection, now in the York museum, contains several specimens of <u>Dorsetensia</u> from Hundale. The three specimens identified by S. Buckman from this collection, have been located, since they still bear L. Richardson's labels. I have cleaned and developed these specimens and they can now be more confidently identified as <u>D</u>. aff. romani (Oppel), <u>D</u>. of. <u>licetraca</u> and <u>D</u>. cf. <u>subtecta</u> S. Buckman, which is in full accord with Buchman's opinions (S. Buckman, 1912, p.207). There are several other specimens, mainly fragmentary, of <u>Dorsetensia</u> in the Herries collection, which look as though they came from the tidal reefs at Hundale and a single specimen of <u>D. liostraca</u> in the Scarborough museum (G.63) also from this locality.

There are no exposures in the Ravenscar Shale south of Hundale and it is significant that no specimens of Dorsetensia have been found south of this point either. To the north at Ravenscar, G. Farrow informs me (pers. com.) that the ammonite recorded by him (Farrow, 1966, p.127), from the basal 3.0m. of his bed SB 27 at Beast Cliff (0V993006), was probably a specimen of Dorsetensia. Similarly J. Wright has found a calcareous nodule, containing a specimen of Dorsetensia, at Ravenscar some 3-4.0m. above the base of the same bed (Wright, pers. com. in lit.). These records are confirmed by the occurrence of a specimen of \underline{D} . aff. liostraca (CP.2747) immediately above the horizon rich in <u>Isognomon</u> (Farrow, 1966, p.127, bed SB 27), which is 3.5m. above the top of the Crinoid Grit. The specimen of D. pulchra S. Buckman, found by P.C. Sylvester-Bradley is now in the Institute of Geological Sciences' collections (83873). This specimen, in a clay-stone matrix rich in Melezgrinella, was found in a freshly slipped mass of shale at Monk's Walk Wood, Sneaton, (NZ896086), and it was identified by W.J. Arkell. It is perhaps a little fragmentary for positive identification, but it must have come from the thick shale which here, as at Ravenscar, rests on the Crinoid Crit (Fox-Strangways, 1915, p.43).

No other genera of ammonites have been found within the Raven-Scar Shale at present. This contrasts strongly with the Romani subzone faunas of Dorset (Whicher & Falmer, 1972) and the Hebrides (Norton, 1965), where Stephanoceratid ammonites are abundant. However, in the north German successions there are horizons which

are rich in <u>Dorsetensia</u>, often to the exclusion of other ammonite groups (Westermann, 1954, p.23), there is thus no difficulty in correlating this Ravenscar shale fauna with the Romani subzone.

(b) The subzone of <u>Stephanoceras humphriesianum</u> (J. de C. Sow.) The majority of the rest of the ammonites to be recorded here originate from this subzone.

(i) The 'Lower Belemnite Bed' of White Nab

This, one of the lowest beds visible at White Nab (TA059865) during low tides, takes its name from the common specimens of Megateuthis elliptica (Miller) to be found (Hudleston, 1874, p.315). Stephanoceratid ammonites are relatively common at this horizon (bed 3) and include Stephanoceras aff. gibbosum (S. Buckman), (CP.2410); S. aff. crassicostatum (Qu. emend. Renz), (CP.2975, ex. M. Bradshaw, & CP. 2834); S. cf. zieteni (Qu. emend. Renz), (CP. 3095) and S. aff. triptolemus (Bean m.s. emend. S. Buckman), (CP.2215). A single, very large specimen of S. nseudo humphriesi Naubeuge (CP 2725), has been found loose at the very base of the exposure at White Nab. Examination of the matrix of this specimen, including comparisons with the aid of thin sections, shows that this specimen came from a lower horizon still, that is bed 2. More collection is required from this horizon, which is only exposed at the very lowest tides. All of the above ammonites would suggest a correlation with the Humphriesianum subzone s. str., since they are identical to the forms commonly found at the type horizon of S. humphriesianum in north Dorset (Whicher & Palmer, 1971, p.113, middle of bed 4).

The 'Lower Belennits bed' has been correlated in Text figure 2, with bed 3 at Gristhorpe Bay, since the latter is the probable source



Text figure 2

A correlation of the main ammonite horizons recognised within the Scarborough Formation, in the coastal exposures between Gristhorpe and Ravenscar. of a specimen of <u>S</u>. <u>humphriesianum</u>. This specimen, which is one of the closest I have seen to the lectotype, was collected by J. Phillips in 1855 (cited, Phillips, 1860, p.xlii), and it is now in the Oxford University Museum (J.16200). Examination of the matrix of this ammonite, which is a dark grey, calcareous, highly fossiliferous shale, with a yellow bloom on the weathered surface of shell fragments, would suggest a provenance from bed 3, which has an identical lithology.

(ii) The 'Black Rock Nodule Bed'

At Black Rock, Scarborough (TA051870), there are extensive exposures of this layer of siderite rich clay-stone nodules, from which previous authors have recorded (Teloceras) 'Ammonites' blagdeni (Williamson, 1840; Wright, 1860, & Hudleston, 1874). This layer of nodules (bed 7) does yield relatively common specimens of Teloceras. However, none of those which I have collected, or examined in other collections, from this horizon are at all close to T. blagdeni, since they all show finer ribs, a higher primary rib density, a more rounded venter and a thinner whorl section, than is typical of \underline{T} . blagdeni. Specimens from this horizon at Black Rock include; T. cf. acuticostatum Weisert, (CP.1267); T. aff. frechi (Renz), (CP.1268), and T. aff. acuticostatum, (G.73). This bed has also produced specimens of <u>Stephanoceras</u> crassicestatum (CP.2680), which together with the Teloceras, accord with a position high up in the Humphriesianum subzone for this bed, as with the middle part of the Chorne Road-stone of north Dorset (Whicher & Palmer, 1971).

Species of <u>Teleceras</u> with a high primary rib density, similar to those recorded above, have been found at Hundale Point. These include; <u>T. of. lotharingicum Maubeuge</u>, (G.72), from 2.0m. below the

top of the Scarborough Formation, that is approximately bed 22, with which there is a close match in matrix, and \underline{T} . cf. <u>acuticostatum</u> Weisert (G74), which was found loose on the beach, but which shows a similar chocolate-brown, mud-stone matrix, rich in <u>Meleagrinella</u>, to that of the previous specimen.

Specimens of <u>Teloceras</u> have also been found in siderite rich nodules at other localities. The ammonites recorded by Farrow (1966, Text fig. 3), from Gristhorpe Bay and Red Cliff are of this genus (Farrow, pers. com.). J. Wright has recently found a specimen of <u>Teloceras</u> in a siderite nodule, at the bottom of the cliff at Ravenscar (Wright, pers. com., <u>in lit</u>.). This must have come from the top 5.0m. of the Scarborough Formation, where there are layers of siderite rich nodules (Bate, 1965, p.85, bed 14) - see Text fig. 2.

(c) The subzone of Teloceras blagdeni (J. Sow.)

The upper part of the White Nab'Iron-stones' at Scarborough is of this age, as it yields relatively common specimens of <u>Teloceras</u>, including <u>T. blagdeni S. str</u>. An impression of the venter of a specimen of <u>T</u>. aff. <u>blagdeni</u> has been seen in bed 8 at White Nab, whilst in the suprajacent bed 9 at Black Rock, specimens of <u>T</u>. <u>blagdeni</u> (CP.2933) and <u>T</u>. sp. (CP.3133) have been collected. This latter horizon is the undoubted source of the numerous, often squashed, specimens of <u>Teloceras</u> to be seen in various museum collections, including a fine fragment of the body-chamber of <u>T</u>. <u>blagdeni</u>, from the Williamson collection in the Manchester Museum (L9561). A single specimen of <u>T</u>. aff. <u>blagdeni</u> in the Scarborough Museum (G.71) has a hard, rust-red coloured matrix, which would suggest a provenance from bed 8 at White Nab. These ammonites thus substantiate Williamson's

record of 'Ammonites' blandeni from his bed f. (Williamson, 1840, p.232).

(d) Stratigraphically unlocalised Ammonites.

The most important ammonite from the Scarborough Formation, is undoubtedly the recently re-discovered type specimen of <u>Stephanoceras</u> <u>triptolemus</u> (Bean m.s. in Morris & Lycett), (British Museum, - 46553). This specimen will be described and refigured at a later date, and all that need be said of it at present, is that its matrix, a dark Grey mud-stone, weathered to a red colour, and with a Belemnite and several specimens of <u>Melearrinella</u> imbedded; would suggest a provenance from the Black Rock nodule bed'; it is undoubtedly Humphriesianum Subzone in age.

There are large numbers of Scarborough Formation ammonités in Various museums throughout the country. The Sedgwick Museum has an excellent selection of material from this horizon, especially that in the Leckonby collection, which includes a fine specimen of <u>Doraetensia</u> <u>deltafalcata</u> (J 1457), very similar to that recorded here from Hundale. Whilst the Geological Survey have several specimens of <u>Telocaras</u> from the Lycett collection, mainly of the <u>T. lugdenensis</u> group, which probably originated from bed 9 at Black Rock; the best collection of Scarborough beds material is in the Scarborough Natural History Museum. The following specimens are all from the Scarborough district, and most probably from the 'Black Rock nodule Bed'; <u>Shirroceras</u> of. <u>macrum</u> (Qu. emend. Renz.), (G.75); <u>Stenhanoceras</u> of. <u>crassicostatum</u> (Qu. emend. Renz), (G.70), and <u>Teloceras lugdenensis</u> (G.67). Again a Humphriceianum subzone age is indicated for this bed.

A search through the material in the Scarborough Euseum showed



no sign of the specimen of Parkinsonia; recorded by Wright (1860. p. 29) as being in this collection. I have considerable doubts over the identification of this specimen, which was supposedly from White Nab. As noted by Buckman (1912, p.208) the matrix of some specimens from the Kellaways Rock is very similar to that of the Scarborough Formation, and this 'Parkinsonia' may well have been a misplaced specimen from this former horizon. The specimen of Parkinsonia identified from White Nab by Richardson, (1912,p.195). is still in the Herries collection at York, and it proves to be a squashed fragment of Teloceras. There is no evidence for the Parkinsoni Zone in the Scarborough Formation, similarly there is no ammonite evidence for any horizon lower than the Romani subzone of the Humphriesianum Zone. However, as at Hundale Point for example, there is almost 1/2 of the Scarborough Formation below the lowest specimen of Dorsetensia, the occurrence of lower stratigraphic horizons, such as the Sauzei Zone, could not be ruled out. The distribution of ammonites, as known to the present, is summarised in text fig. 3.

IV. THE LITHO-STRATIGRAPHY OF THE SCARBOROUGH FORMATION OF THE NORTH-EAST YORKSHIRE MOORS.

The three-fold lithological subdivision of the Scarborough Formation suggested by Richardson, into shale, sandstone and limestone divisions, is a gross over-simplification. Many of the lithostratigraphic units discussed here (see Table 2) are highly lenticular, and quickly disappear when traced laterally. When attempting a broad interpretation of rapidly changing lithologies, such as those of the Scarborough Formation, it is essential to have a bio-
The Scalby

Formation

	The 'Transition Shales' (0.0-3.5m.)
	The White Nab 'Iron-stone' member (0.0-2.0m.)
The Scarborough	The Lambfold-hill Grit member (0.0-3.0m.)
Formation	The Ravenscar Shale member (0.0-36m.)
	The Spindle-Thorn Limestone member (0.0-4.0m.)
	The Crinoid Grit member (0.0-3.5m.)
	The Hundale Shale member (0.0-5.0m.)

The Gristhorpe

member

Table 2

The constituent members of the Scarborough Formation in the north-east Yorkshire Moors. stratigraphic framework against which to correlate the various beds. A step in this direction was made by Bate, whose two ostracod zones demonstrated the essentially isochronous nature of the 'Crinoid Grit', (Bate, 1965, text figs. 2 & 3). A study of existing ammonite horizons would indicate that there are two other isochronous beds, the units here called the Ravenscar Shale Member and the White Nab 'Ironstone' Member. Using these three horizons it is possible to produce a slightly more accurate picture of the variations in lithofacies - see text fig. 4. The lithostratigraphic divisions described below are only of any value to the **e**ast of Rye Dale and Rievaulx, since elsewhere there is more limited biostratigraphic control and far greater variation in lithology. It is to be noted that the lithostratigraphic scheme suggested here is still only provisional, as more work, particularly on the inland sections, is needed.

The Scarborough beds have recently been elevated to the formal rank of a Formation within the Ravenscar Group (Hemingway & Knox, 1973, p.533). Following the recommendations of the Geological Society of London (Harland <u>et al.</u>, 1972), formal subdivision of these beds is thus here undertaken at the level of 'member'.

(a) The Hundale Shale Member.

The lowest marine horizon within the Scarborough Formation seen on the coast, consists of a series of sandy shales and calcareous mud-stones, rich in <u>Gervillia</u>, lying beneath the 'Crinoid Grit', and best exposed at Hundale point, the type locality for this new lithostratigraphic unit (beds 7-2). The age of these beds is doubtful, since no ammonites have been found, but it must be pre-Humphriesianum Zone, and is most likely Sauzei Zone in age. These beds are often absent inland, but where present, such as at Bogmire Gill (Bate, 1965,

p.91, bed 1), they tend to be more calcareous, and nearer to a pure limestone. This unit is some 5m. thick at Hundale, and has a very similar thickness at Ravenscar, where there are some thin sandstone beds present. The occurrence of these sandstones is not unusual, since around the edge of the basin of deposition of the Scarborough Formation sandstones are predominant at almost all horizons, (Bate, 1965, p.96).

(b) The Crinoid Grit Member.

This term was used by Richardson (1912, p.195) to describe the sandstone, rich in comminuted crinoid debris, which occurs towards the middle of the Scarborough beds. Whilst it is most typically the type focefity, developed inland, as around Helmsley Moor, (Bogmire Gill, Bate, 1965, p.91, bed 2, - 3.5m. thick) there is a good thickness of the Crinoid Grit on the Coast at Ravenscar and Hundale point. At the latter locality it is some 2m. thick, with many crinoid ossicles in the upper part, and with prominent ripple marked surfaces towards the middle (beds 12-8).

(c) The Spindle-Thorn Limestone Member.

This limestone overlies the 'Crinoid Grit' in the main part of the Yorkshire Moors, around Spaunton and East Moors, and it was once well seen in an old quarry at Spindle-Thorn, the type locality, (SE 717928 - Richardson, 1912, p.197), where it is about 2m. thick. Although Richardson introduced this term for the inland limestone occurrences (<u>loc. cit.</u>), there are shaly limestones and highly calcareous mud-stones at the same horizon on the coast, (Hundale, beds 13-19).

(d) The Ravenscar Shale Member

This term is introduced here for the thick shale unit with highly fossiliferous nodules, which is so well seen on the coast at Hundale point and Ravenscar cliffs. Although it is probably best exposed at these two localities, where it has yielded diagnostic ammonites, proving it to be Romani subzone, Humphriesianum Zone in age the Ravenscar Shale is at its thickest in the region of Harland Beck and Farndale, where the survey recorded over 36m. of shale, (Fox-Strangways, Reid & Barrow, 1835, p.40). Although thicker at inland localities, exposures are not good, and the most complete exposures are at Hundale and Ravenscar, and the latter, where there are some 13.5m. of Ravenscar Shale, may be taken as the type locality see Bate 1965, p.85, beds 12 & 13.

(c) The Lambfold-Hill Grit Member

This name was introduced by Richardson (1912, p.197) for a sandstone in the upper part of the Scarborough Formation, at Lambfold Hill, Pockley Moor, (SE 615945), where it forms an important topographic feature. This unit is well exposed in Bogmire Gill, near the type locality, where it is 3.0 m. thick, (Bate, 1965, p.91, bed 5) and may be traced intermittently eastwards as far as Bloody Beck (SE 945981), where beds 13-15 (Bate, 1965) probably represent a 'feather edge'.

(f) The White Nab 'Ironstone' Member

At White Nab and Black Rock Scarborough there are extensive exposures of a series of gritty shales, ironstained, sandy, limestones and siderite rich mud-stone nodules, here called the White Nab 'Ironstone' Member. The term 'iron-stone', in this case is/a mis-nomer, since the siderite content of the beds is low, however, weathering produces a characteristic deep red colour to these rocks. At White Nab, the type locality, this unit is 1.75 m. thick and comprises beds 7-16. The siderite rich nodules of this member, yielding <u>Teloceras</u>, have been traced the full length of the coast, there is thus every reason to believe that this horizon in the upper part of the Scarborough Formation is isochronous over most of the eastern part of this district. The particular bed of siderite rich nodules, yielding most of the specimens of <u>Teloceras</u> at Scarborough is here separated informally as the 'Black Rock Nodule bed' (bed 7).

(g) The 'Transition Shales'

No formal name is given to this unit, since it is impossible to be certain that we are dealing with the same Member at different localities. A series of sandy, ironstained, sulphurous, shales are often present in the upper part of the Scarborough Formation, forming a gradual transition to the non-marine sandstones of the Scalby Formation. Thus at Hundale Point there are approximately 2.0 m. of these shales, which are progressively sandier and less fossiliferous up towards the junction with the over-lying Scalby Formation. These shales are often absent, mainly due to 'wash-outs' coming down from the base of the Moor Grit (= basal Scalby beds). Thus at Gristhorpe, in some parts of the cliffs, up to 1.5 m. of these sandy shales have been cut out. These shales are sometimes sufficiently arenaceous to form a soft, poorly cemented sandstone; as in the top of the cliff at Ravenscar, (Bate, 1965, p.85, bed 16).

V. DESCRIPTION OF COASTAL EXPOSURES

As already noted in the Introduction, the coastal exposures of the Scarborough Formation are the only ones to show a large enough area of outcrop for the collection of ammonites. Thus those at Ravenscar, Hundale, Scarborough and Gristhorpe are described here, starting with the most northerly and working south.

(a) Ravenscar

Sections at Ravenscar have been described by Fox-Strangways (1915, p.39), Bate (1965, p.85) and Farrow (1966, p.116). The most accessible and complete exposures are in an old quarry on the cliffface (NZ991009), but even these are becoming increasingly obscured by talus and vegetation. Thus in the last few years it has become difficult to make out the finer details of this section. Taking this into account, this exposure is only described here in the most general terms and for fuller details earlier works should be consulted (Bate, 1965, p.85).

SCARBOROUGH FORMATION

6/ A series of flaggy sandstones, which are more shaly towards the base.
4.04.5m.

White Nab Ironstones

A series of sandy, calcareous shales, with sideritic, clay-stone nodules.

Teloceras sp. (by matrix) 1.0-1.5m.

Ravenscar Shale

51

41

A series of grey shales, which are more sandy towards the

top. These beds are very fossiliferous, particularly 3.5 m. from the base, where there is a layer of <u>Isognomon</u>, surmounted by a horizon rich in crinoid debris.

Dorsetensia aff. liostraca S. Buckman

10.5 -13.5m.

The Crinoid Grit

3/

A massive, sphaeroidally weathered, sandstone, with comminuted crinoid debris towards its top. 1.4m.

The Hundale Shales (2)

2/ A poorly exposed sequence of inter-bedded limestones, mudstones, sandy shales and sandstones, with a prominent horizon rich in <u>Gervillia</u> towards the base.
3.4-4.5m.

1/ A series of thinly bedded, flaggy sandstones. 3.0-4.5m.

(b) Hundale Point, Cloughton

This is the best exposed coastal section (TA026948), which shows one of the thickest developments of the Scarborough beds. Although it has been previously described, (Hudleston, 1874; Fox-Strangways, 1915; Bate 1965), this section has never been given the attention which its abundant faunas warrant. Apart from the term '<u>Isognomon</u> bed' which is new, all the other informal bed names used here originate from Hudleston (1874).

26/ 'The Moor Grit'

A massive, cross-bedded sandstone, which is more thinly bedded towards the base. seen to 2.0m.

THE SCARBOROUGH FORMATION (25-2)

25/ 'The Transition Shales, (25-24)

A laminated, yellow-grey coloured, sandy shale, with brown, sandy layers some 2.5 cm. thick.

0.46. 0.67m.

0.84m.

0.90m.

24/ A series of grey, sulphurus shales, which are more sandy towards the top.

The White Nab'Ironstones' (23-22)

23/ A series of sulphurous, grey, sandy shales, with two layers of unfossiliferous, very iron rich nodules (c.O.20m. thick), one at the top and the other towards the middle of the bed.

22/ A hard, calcareous, grey-brown mudstone, which weathers to a rusty colour, and which contains numerous well preserved fossils.

> <u>Teloceras</u> cf. <u>acuticostatum</u> (from here by matrix) <u>T. aff. lotharingioum</u> Maubeuge Meleagrinella <u>lycetti</u>

0.15-0.30m.

The Ravenscar Shales (21-20)

21c/ A grey, fissile, relatively unfossiliferous shale.

1.5m.

21b/ A dark grey coloured shale, which is slightly softer than the adjacent beds, hence it tends to be weathered back.

Merateuthis sp.

Liostrea sp.

Pseudomelania lonsdalei

0.23m.

21a/ A very dark grey coloured shale, which is highly

fossiliferous, especially towards the base. There are numerous small, round concretions present, some of which contain uncrushed fossils.

Dorsetensia liostraca S.B. D. subtecta S.B. D. cf. romani (Oppel) D. aff. deltafalcata (Qu.) Megateuthis sp. Cenoceras sp.

Meleagrinella lycetti

6.25m.

20/

19/

A dark grey, very fossiliferous shale.

Meleagrinella lycetti

0.30m.

The Spindle-Thorn Limestone (19-13)

A hard, grey, sandy, impure limestone, which contains the same fossils as the bed below, only more sparsely. Material from this bed has been piped down, via burrows, into the bed below.

Neleaminella lycetti etc. etc.

0.22m.

18d/ A highly fossiliferous, grey, sandy, calcareous shale, with some Thalassinoides style burrows, particularly towards the top. 和新闻的话,并且在这些话的话。 Dorsetensia sp. Camptonectes Sp.

Catinula ampulla (d'Archiac)

Gervillia scarburgensis

Meleagrinella lycetti

Modiola sp.

Liostrea sp.

Lonha marshi

Pinna cuneata Phillips (in an upright position)

Platymya scarburgensis (Phillips)

Pseudolima aff. duolicata

Trigonia costata

T. (Myonhorella) signata (Agass.)

Pentacrinus ossicles

Pseudodiadema depressum (Agass.)

Rhabdocidaris maxima (Munst.)

Serpula aff. plicatilis Goldfuss.

0.82m.

280

18c/ A layer of hard, grey, calcareous nodules, with sparse, but well preserved and uncrushed fossils, similar to those in the bed above.

0.0-0.35m.

18b/ A grey, sandy, calcareous shale, slightly softer than bed 18d, and with sparser fossils.

Cenoceras sp.

Pernostrea sp.

<u>Pleuronya uniformis</u> (J. Sow.) (in life position) 0.28m.

18a/ A grey, gritty, highly bioturbated shale, with common <u>Thalassinoides</u> burrows. <u>Pentacrinus</u> ossicles are common, as are burrowing bivelyes, particularly towards the base of the

bed.

Pleuromya uniformis

Pholadomya lirata

0.50m.

17/ A grey, unfossiliferous, gritty shale, which is distinctly mottled, mainly due to intense bioturbation.

0.36m.

16/ A grey, sandy, calcareous shale, with a similar lithology and fauna, as that of bed 18d.

> <u>Meleagrinella lycetti</u> Lopha <u>marshi</u> etc. etc.

> > 0.43m.

0.46m.

15/ A grey, mottled, sandy, highly bioturbated shale, very similar to bed 17.
0.65m.

'The Isomomon bed'

14/ A grey, sandy, calcareous, highly fossiliferous shale, very similar to bed 18d. Occasional calcareous nodules, containing well preserved fossils are found scattered throughout the middle of the bed. Disarticulated valves of <u>Isognomon</u> are common.

> <u>Stephanoceras</u> (<u>Skirroceras</u>) sp. (0.43m. above 12) <u>Catinula ampulla</u> (d'Archiac) ?<u>Gryphaea</u> sp. <u>Exocyra</u> sp. (attached to the ammonite) <u>Isopnomon isopnomonoides</u> (Stahl.) <u>Lopha marshi</u> <u>Pinna</u> sp. (very large and flat lying). Trigonia costata

13/ A grey, bioturbated and mottled shale, with sparse oysters towards the base. The basal 9 cm. is softer, browner coloured and is often packed with serpulid tubes.

> <u>Lopha marshi</u> Pernostrea sp.

Sernula sp.

0.25-0.30m.

The Crinoid Grit (12-8)

- - - - An oyster-encrusted 'hard-ground'

12c/ A hard, crystalline, sandy limestone, with a very flat top, and containing a large quantity of comminuted crinoid debris.

Rhabdocidaris maxima (Munst.)

Pentacrinus sp.

0.05m.

12b/ A hard, massive, highly bioturbated, calcareous sandstone. Trace fossils, particularly <u>Arenicolites</u>, <u>Teichichnus</u>, <u>Rhizocorallium</u> and <u>Thalassinoides</u> are very common.

0.9m.

12a/ A hard, massive calcareous sandstone, with fine parallel laminations, which is cemented to the base of the bed above. Body fossils are rare, except for a few poorly preserved bivalves towards the base, where there is also some bioturbation.

0.70m.

11/ A dark grey coloured, gritty shale, which grades up into the bed above.

The Upper Iron Scar! (10b)

10b/ A hard, grey mudstone, which weathers to a rust red colour. 0.25m.

10a/ Two, and sometimes three, layers of hard, calcareous sandstones, with ripple marked surfaces.

0.63m.

A hard, calcareous, parallel laminated grit, which grades down into a softer, bioturbated, sandy marl.

Pentacrinus sp. 0.61-0.67m.

8/ A grey coloured, mottled and bioturbated sandy shale. 0.66-0.68m.

The Hundale Shales (7-2)

9/

7/ A dark grey, 'coaly' shale.

0.09m.

'The Lower Iron Scar' (6b)

6b/ A hard, fine grained, dark grey, calcareous siltstone, which weathers to a red colour. Fossils are common towards the top of the bed, but are very difficult to extract, and are best seen on weathered surfaces. This horizon has yielded the most varied Gastropod fauna, of any bed within the Scarborough Formation. Towards the base this bed becomes softer and more shaly, and here fossils are easier to extract.

Cucullaca cancellata

Astarte minima

Gervillia scarburgensis

<u>Meleagrinella lycetti</u> <u>Cloughtonia cincta</u> (Phillips) Pseudomelania laevigata (Mor. & Lyc.)

> 0.33-0.40m.

0.20-0.30m.

A rubbly, ironstained, grey, calcareous shale, which becomes harder towards the top, where it grades into the bed above.

Cloughtonia cincta (Phillips)

5/ A highly fossiliferous, grey, calcareous shale, which is packed full of bivalves, and which has sparse patches of pseudo-coliths.

Pleuromya uniformis

6a/

3/

2/

Pholadomya lirata

'Modiola' leckenbyi Mor. & Lyc.

0.13-0.21m.

4/ A shaly and slightly calcareous, fine sandstone, which has a concentration of <u>Gervillia</u> towards the middle and some sparse and poorly preserved fossils towards the top.

> 0.84-0.95m.

An iron-stained, grey, calcareous, sandy, highly bioturbated shale.

0.41m.

0.66m.

- - - dark shale parting 0.01m. thick - - -

A dark grey, carbonaceous shale, with poorly preserved bivalves and plant and wood fragments. THE CRISTHORPE BEDS

1/

A massive cross-bedded sandstone.

3.5m.

بأربع بيدابا والتراكي متفاركوتي

3.0m. +

0.35m.

(c) White Nab and Black Rock, Scarborough

This section was measured from both the small anticline exposed at Black Rock (TA050870), just to the south of the bathing pool and from the headland of White Nab (TA060865). Whilst the greatest thickness of beds is exposed at the latter locality, the former shows wider areas of outerop, more suitable for the search for ammonites. This section is perhaps the most frequently described of all the Scarborough Formation localities having been the subject of work by Phillips (1829), Williamson (1840), Wright (1860), Hudleston (1874), Fox-Strangways (1904), Bate (1965) and Farrow (1966). However the discovery of several ammonite horizons within these beds made their re-description worthwhile. Two of the informal bed names, 'Black Rock nodule bed' and '<u>Gervillis</u> bed' are new. The rest originate from Hudleston (1874).

18/ 'The Moor Grit'

A massive, cross-bedded sandstone.

THE SCARBOROUGH FORMATION (17-1)

17/ 'The Transition Shales'

A yellow-grey coloured, sandy shale, containing poorly

preserved fossils.

The White Nab Ironstones (16-7) 16/ A dark, brown-grey coloured, shale, with mudstone nodules full of well preserved fossils. 15/

14/

Meleagrinella lycetti (Rollier)

0.23m.

A hard, nodular, argillaceous limestone, weathering a rust red. 0.20m.

A purple-grey coloured shale, with numerous well preserved fossils, which have a white powdery preservation, and with frequent patches of pseudo-coliths.

Meleagrinella lycetti

13/ A brown-grey coloured shale, which grades up into a hard, nodular mudstone.

0.33m.

0.20m.

12/ 'The Gervillia bed'

A hard, crystalline, impure limestone, which weathers to a deep pink colour, and which is packed full of <u>Gervillia</u> scarburgensis Paris.

> 0.15-0.23m.

11/ A thin, gritty shale.

0.0-0.08m.

10/ A hard, pink-grey coloured calcareous mudstone, full of broken shell debris.

0.15-0.26m.

9/ A grey, gritty shale, which weathers to a mottled red/brown - grey colour, mainly due to the presence, particularly towards the top of the bed, of numerous <u>Thalassinoides</u> burrows. These burrows pipe down a darker shale, full of broken shell debris.

Teloceras blagdeni (Sow.)

T. sp.

8/

7/

Megateuthis guinquesulcatus (Blain.)

Pernostrea sp.

0.28m.

A nodular and irregular, grey mudstone, which weathers to a dark red colour.

Teloceras aff. blagdeni

0.15-0.20m.

The Black Rock Nodule bed!

A grey-brown coloured, gritty shale, full of comminuted shell debris and squashed fossils. The basal half of the bed is a darker grey colour and contains common oysters, whilst along the middle of the bed there are scattered irregular claystone nodules. These nodules, up to 0.60m. in diameter, are hard, splintery, dark grey in colour and weather to a light pink-grey colour, which is in direct contrast to the rust red of the beds above. Well exposed at Black Rock, where the nodules are scattered over the fore-shore, this bed is at its thickest, and has yielded most ammonites from this point.

Teloceras cf. acuticostatum Weisert

T. aff. frechi (Renz)

T. aff. <u>blagdeniformis</u> (Roche) Stephanoceras crassicostatum (Qu. emend. Renz)

<u>Gervillia scarburgensis</u> Neleagrinella <u>lycetti</u>

Pernostrea sp.

Pseudomelania lonsdalei Mor. & Lyc.

Megateuthis quinquesulcatus

0.30-0.60m.

? The Lambfold-Hill Grit (6-4)

6/ 'The Chemnitzia Limestone'

A hard, crystalline, grey, sandy limestone, which is iron-stained, massive and divided into two blocks by a parting 1.0m. from the base, the upper course being the more coarsely sandy. <u>Pseudomelania lonsdalei</u> ('<u>Chemnitzia</u>' <u>auctt</u>.) is very common, particularly on the upper surfaces.

1.95m.

0.9-0.30m

5/ 'The Upper Belemnite bed'

A thin, gritty, grey shale, which thickens to the north and which contains sporadic Belemnites.

4b/ A hard, sandy, limestone, similar to bed 4a, which thickens to the north.

0.0-0.25m.

4a/ A hard, sandy, blue 'hearted', crystalline limestone, which weathers rust-red and which contains numerous badly preserved and broken fossils. The top surface of the bed, which forms an extensive shelf at low tide, is covered with <u>Thallassinoides</u>. 0.33m.

3/ 'The Lower Belemnite bed' A grey, fossiliferous shale, with abundant slightly squashed <u>Meleagrinella</u> and large Belemnites. The calcareous mud-stone nodules, up to 0.45m. in diameter, found at the middle of the bed, contain well preserved fossils and weather to a light pink colour, similar to that of the nodules from bed 7, from which they are distinguished by their softer and less splintery nature.

<u>Stephanoceras aff. crassicostatum</u> <u>S. aff. gibbosum</u> <u>S. cf. zieteni</u> (Qu. emend. Renz) <u>S. aff. triptolemus</u> (Bean in Mor. & Lyc.) <u>Gresslya sp.</u> <u>Meleagrinella lycetti</u> <u>Pernostrea sp.</u> Pentacrinus ossicles

> 0.40-0.48m.

2/ A hard, grey, sandy, calcareous, shelly mud-stone, which weathers to a rust red colour and which is full of broken shell debris.

Stephanoceras cf. pseudo humphriesi Maubeuge.

0.20-0.35m.

A blocky, bioturbated, marly shale.

(from here by matrix)

seen to 0.30m.

Section incomplete, perhaps a further 3.0m. + below low tide

line.

1/

(d) Gristhorpe Bay

This section which was taken from the south side of the Yons Nab headland (TAO84841), has been previously described by Phillips (1835), Wright (1860) and Bate (1965).

Base of the SCALBY FORMATION

9/ A hard, yellow-grey coloured sandstone, with carbonaceous layers, and which grades down into the bed below.

seen to 0.30m.

THE SCARBOROUGH FORMATION (8-2)

8b/ A grey, sandy shale, mottled with patches of yellow sandy material.

0.66m.

8a/ A soft, well bedded, grey sandy shale.

0.34-0.43m.

(A channel cuts down from bed 9 to the top of bed 7b).

7b/ A compact, very sandy shale, which is grey-brown in colour, and which has a softer, dark grey coloured shale fill to the numerous burrows present. This bed weathers to a yellow-brown colour.

0.82m.

0.36m.

7a/ A grey, silty shale, which has a gradational contact with the bed above. There are occasional claystone nodules towards the top of the bed, full of shell debris.

Pleuromya uniformis

Megateuthis quinquesulcatus

Rhabdocidaris maxima

A hard, silty, very impure limestone, which has a splintery fracture, weathers to a bright red colour and which forms a small reef on the beach at low tide.

<u>Meleagrinella lycetti</u> <u>Trigonia</u> cf. <u>costata</u> <u>Teloceras</u> sp. (<u>Teste</u> G. Farrow)

6/

5/

4/

3/

0.22-0.38m.

A grey, gritty, ironstained, relatively unfossiliferous shale, with a layer of impersistent clay-stone nodules, towards the top of the bed, full of shell debris.

> 0.28-0.50m.

A hard, grey, silty shale.

0.12-0.15m.

A light grey, calcareous, highly fossiliferous shale, which is divided by a parting into two, approximately equal courses. The bottom course is darker, softer and less fossiliferous than the top. This bed forms a prominent reef, well out into the bay at low tide, the upper surface of which is covered with shells and shell fragments, which have weathered to a yellow colour.

<u>Meleagrinella lycetti</u> Pholadomya lirata

Pseudolima aff. duplicata

Trigonia costata

<u>Lopha marshi</u> Pernostrea sp. Pseudomelania lonsdalei

Stephanoceras humphriesianum (J. Sow.) - Phillip's collection, University of Oxford, from here by matrix. 0.64m.

2c/ A grey, laminated, sandy, highly bioturbated, fossiliferous shale.

0.11m.

2b/ A grey, laminated, sandy shale, with only sparse fossils. 0.31m.

2a/ A laminated sandy shale, with a dark charcoal grey colour, which weathers to a streaky brown and yellow colour. There is a lumachelle of oysters towards the middle of this bed, which has a gradational contact with the bed below.

- gradational contact

Liostrea sp.

0.26m.

THE GRISTHORPE BIDS

1/ A massive sandstone, with laminated carbonaceous layers, and grading up into the above.

seen to 0.5m.

I. STRATIGRAPHIC SUMMARY

The ammonite evidence for the correlation of the Scarborough Formation is summarised in text figs. 2 and 3, and it would indicate that only the top half of the beds, as exposed at Hundale Point, belong to the Humphriesianum Zone. It is probable that the base of this ammonite Zone coincides with the base of Bate's zone of Glyptocythere scitula. The horizons below this are pre-Romani sub-



Text figure 4

The lithostratigraphic subdivisions of the Scarborough Formation between Rye Dale and the coast at Cloughton.

zone in age. and most likely to be correlated with the late Sauzei Zone. These new ammonite records have enabled a more detailed correlation of the coastal exposures of the Scarborough Formation The Ravenscar Shale, where present, would appear to be to be made. isochronous, as is the Black Rock nodule bed of the White Nab Member. All of the beds exposed below this latter horizon at White Nab are lower Humphriesianum subzone in age. These beds include the horizons equated with the Lambfold-Hill Member (beds 6-4). It is thus clear that these sandstones cannot possibly be correlated with the Crinoid Grit, as has been suggested by Bate (1965, p.98) and Farrow (1966. fig. 3), as they have an entirely different stratigraphic position. Allowing for the unfossiliferous nature of the top of the Ravenscar Shale at Hundale, there are strong reasons for suspecting the presence of an unconformity between the base of bed 22 and the top of the shales at that section. This would account for the lack of ammonites at Hundale, which are characteristic of beds 1-6 at White Nab. This idea is supported by the nature of the surface upon which the Black Rock nodule bed' (7) rests at Scarborough; it has all the characteristics of a 'hard-ground'.

The litho-stratigraphic picture is more complex and is summarised in text fig. 4. Bate's concept of a shallow marine basin, opening to the east, and with a peripheral area of sandstone deposition, (Bate, 1965, p.96, fig. 4), would appear to be essentially correct for the Scarborough Formation; thus the thickest deposits of these beds exist along the axis of the basin - the Ravenscar/Blea Wyke district. This totally contradicts Farrow's thesis of a bathometric gradient from Ravenscar to Scarborough, with the deeper water, at the southern end. This contradiction can easily

be explained, since in his work on the trace-fossils, Farrow did not compare stratigraphically equivalent horizons. Thus at White Nab, the siderite rich nodules with relatively common ammonites are not the deep water facies equivalent to all the beds exposed at Ravenscar, as suggested by Farrow, but are the direct equivalent solely to some thin beds at the very top of this latter section. (see fig. 2). Re-interpreting Farrow's work in the light of this evidence, it would seem that the Scarborough Formation was first laid down in a shallow and relatively small basin, centred on the Ravenscar area, where sandstones rich in Thallassinoides and Rhizocorallium were deposited. With the deepening of the basin. marine influences were increased and the greatest abundance of ammonites coincides with the greatest areal extent of these beds. which are represented only by the upper part of the sequence at Ravenscar, and by all of the beds at White Nab and Gristhorpe: the earlier shallow water facies being absent at the two latter localities. Thus, there is not a lateral, but a vertical 'bathymetric gradient'. upwards through the Scarborough Formation, with the progressive overstepping of younger, but deeper water beds.

VII. ACKNOWLEDGEMENTS

a fathana a star star

I should like to thank the various individuals who have either donated specimens or given valuable stratigraphic information, that is: M. Bradshaw, G. Chadwick, G. Farrow, J. Senior, P. Sylvester-Bradley and J. Wright. I must express a particular debt of gratitude to the latter, without whose kind help, this work would not have been completed. At this point I should also like to thank the curators of the following institutions for access to material in their care: The British Museum (N.H.) and Institute of Geological Sciences, London; The Sedgwick Museum, Cambridge; the Oxford University Museum; the Scarborough Natural History Museum and the Yorkshire Museum, York.

VIII. REFERENCES

Arkell, N. J. 1933,

The Jurassic System in Great Britain. Oxford University Press, (Oxford) 681 p.41 pl.

Bate, R.H. 1965,

Middle Jurassic Ostracoda from the Grey Limestone Series, Yorkshire. Bul. Brit. Mus. (N.H.), Geol., <u>11</u> (3) : 75-133, 21 pl.

Buckman, S.S. 1910,

Certain Jurassic (Lias-Oolite) strata of South Dorset. Q. Jl. geol. Soc. Lond. <u>66</u> : 52-89.

1912,

Ammonites from the Scarborough Beds. Appendix 1 to Richardson, 1912. Proc. Yorks. geol. Soc. <u>17</u>: 205-8.

Conybeare, W.D. & Phillips, W. 1822,

Outlines of the Geology of England and Males. Phillips, (London), 470p. 6 figs.

Cox, L.R. & Arkell, W.J. 1948-50,

A survey of the Mollusca of the British Great Colite. Palaeontograph. Soc.

Drake, H.C. 1908,

Unrecorded cephalopods from the Yorkshire Oolites. The Naturalist, No. 619 : 296-7.

Farrow, G. 1966,

Bathymetric zonation of Jurassic Trace fossils from the coast of Yorkshire, England. Palaeogeography, Palacoclimatology & Palaeoecology <u>2</u>: 103-151.

Fox-Strangways, C. 1892,

The Jurassic Rocks of Britain, vol. 1 - Yorkshire. Mem. Gool. Surv. Grt. Brit. : ix + 551p.

1904,

The Geology of the Oolitic and Cretaceous rocks South of Scarborough. Mem. Geol. Surv. Grt. Brit. : 119p. xi pl.

1915,

A Second revised edition of, Fox-Strangways and Barrow. The Geology of the Country between Whitby and Scarborough. Mem. Geol. Surv. Grt. Brit.: IV + 144 P.

> c. _, Reid, & Barrow, G. 1885

The Geology of Eskdale, Rosedale. Mem. Geol. Surv. Grt. Brit. :65p.

Harland, W.B. (Chairman) et al., 1972,

A concise guide to stratigraphic procedure. Jl. geol. Soc. <u>128</u>: 295-305.

Haug, E. 1891,

Les Chaines Subalpines entre Gap et Digne. Bull. Serv. Carte geol. Fr., III; 21: 199pp.

Hemingway, J.E. 1951,

The report of the Whitby Field meeting. Proc. Yorks. geol. Soc. 28 : 118-122.

, & Knox, R.W. O'B. 1973,

Lithostratigraphical nomenclature of the Middle Jurassio Strata of the Yorkshire Basin of North-East England. Proc. Yorks. geol. Soc. <u>39</u> : 527-535. Hemingway, J.E., Wilson, V., & Wright, C.W. 1963, (2nd. edit. 1968) Geology of the Yorkshire Coast. Geol. Assoc. Guide No. 34.

, Wright, J.K., & Torrens, H.S. (Editors), 1969, International Field symposium on the British Jurassic. Excursion Guide 3 - North-East Yorkshire. University of Keele (Keele). : 47 p.

Hudleston, W.H. 1874,

The Yorkshire Oolites, pt. 1 - The lower Oolites. Proc. Geol. Assoc. <u>3</u>: 283-333.

1882,

Contributions to the Palaeontology of the Yorkshire Oolites. Geol. Mag. Decade II, 2: 145-151.

Kumm, A. 1952,

Das Mesozoikum in Niedersachsen : Der Dogger. Geol. u. Lagerstätten Niedersachsen, <u>2</u>, 186 p., 21 tab., 16 figs.

Mascke, E. 1907,

Die Stephanoceras-Verwandten in den Coronaten Schichten von Norddeutschland. Dissert. Univ. Gottingen, (Gottingen) : 38p.

Morris, J. & Lycett, J. 1850-3,

A Monograph of the Mollusca from the Great Oolite, Chiefly from Minchinhampton and the Yorkshire coast. Palaeontograph. Soc.

Morton, N. 1965,

The Bearreraig Sandstone Series (Middle Jurassic) of Skye and Raasay. Scot. J. Geol. <u>1</u>: 189-216. Die Juraformation Englands, Frankreichs und des südwestlichen Deutschland. Jb. Ver. vaterländ Naturkde. Württ. <u>12-14</u>: 837 p.

Parsons, C.F. 1974,

Towards a revision of the Lithostratigraphic Nomenclature of the Middle Jurassic of North-East Yorkshire. Chapter , Unpublished Ph.D. thesis, Univ. Keele.

Pavia, G. & Sturani, C. 1968,

Etude biostratigraphique du Bajocien des Chaines Sub-Alpines aux Environs de Digne. Boll. Soc. Geol. Italy <u>87</u> : 305-16.

Phillips, J. 1829,

<u>Illustrations of the Geology of the Yorkshire Coast</u>. (York): xvi + 192 p. xiv pl.

1835,

Illustrations of the Geology of the Yorkshire Coast. 2nd edition.

1858,

On some comparative sections in the Oolitic and Ironstone series of Yorkshire. Q. Jl. geol. Soc. Lond. <u>14</u> : 84-98.

1860,

Presidential address to the London Geological Society. Q. Jl. geol. Soc. Lond. <u>16</u> : xxvii-lv.

Rayner, D.H. & Hemingway, J.E. 1974, (Editors),

The Geology and Mineral resources of Yorkshire. The Yorks. geol.

Soc. pp.1-405 + ix, 7 Pl. 79 figs.

Richardson, L. 1912,

The Lower Oolitic Rocks of Yorkshire. Proc. Yorks. Geol. Soc. 17 : 184-215, pls. xxii-xxiv.

Sedgwick, A. 1826,

On the Classification of the strata which appear on the Yorkshire coast. Ann. Philosophy (N.S.) 11 : 339-362.

Sturani, C. 1971,

Ammonites and Stratigraphy of the <u>Posidonia alpina</u> beds of the Venetian Alps. Mem. 1st. Geol. Min. Univ. Padova 28 : 1-190p. 16pl.

Sylvester-Bradley, P.C. 1953,

A stratigraphical guide to the fossil localities of the Scarborough district, pp.19-48, in - The Natural History of the Scarborough District, Vol.1 - Geology and Botany. The Scarborough Field Naturalists Society (Scarborough) : xii + 296 p.8 pl.

Westermann, G.E. 1954,

Monographie der Otoitidae. Beih. Geol. Jb. 15 : 364p. 33 pl.

Whicher, J. & Palmer, C.P. 1971,

Preliminary descriptions of a temporary exposure in the Middle Inferior Oolite, near Sherborne, Dorset. Proc. Dorset Nat. Hist. Archaeol. Soc. <u>92</u> : 110-119.

Williamson, W.C. 1840,

On the distribution of fossil remains on the Yorkshire coast. Trans. Geol. Soc. Lond. (2nd. ser.) 5 : 223-242.

Wright, T. 1860.

On the subdivisions of the Inferior Oolite in the South of England, compared with the equivalent beds of that formation on the Yorkshire coast. Q. Jl. geol. Soc. Lond. 16 : 1-48.